

Customer No. 20350
TOWNSEND and TOWNSEND and CREW LLP
Two Embarcadero Center, 8th Floor
San Francisco, California 94111-3834
(415) 576-0200

ASSISTANT COMMISSIONER FOR PATENTS
BOX PATENT APPLICATION
Washington, D.C. 20231

11/28/00

10960 U.S. PRO

Attorney Docket No. 023070-114700US
Client Ref No. 2001-055-1
"Express Mail" Label No. EL394879473US
Date of Deposit: November 28, 2000

JC929 U.S. PRO
09/724857

11/28/00

Sir:

Transmitted herewith for filing under 37 CFR 1.53(b) is the

- ☒ [X] patent application of
☐ [] continuation patent application of
☐ [] divisional patent application of
☐ [] continuation-in-part patent application of

Inventor(s)/Applicant Identifier: Koen Weterings

Nestor R. Apuya

Robert B. Goldberg

For: POLYNUCLEOTIDES USEFUL FOR MODULATING TRANSCRIPTION

Enclosed are:

- ☒ [X] 69 page(s) of specification
☒ [X] 8 page(s) of claims
☒ [X] 1 page of Abstract
☒ [X] 20 sheet(s) of [] formal [X] informal drawing(s).
☒ [X] 11 page of sequence listing
☒ [X] A [] signed [X] unsigned Declaration.

11/28/00

11/28/00

11/28/00

11/28/00

11/28/00

11/28/00

11/28/00

11/28/00

11/28/00

11/28/00

11/28/00

11/28/00

11/28/00

11/28/00

11/28/00

11/28/00

11/28/00

11/28/00

11/28/00

11/28/00

11/28/00

11/28/00

11/28/00

11/28/00

11/28/00

11/28/00

11/28/00

11/28/00

11/28/00

11/28/00

11/28/00

11/28/00

11/28/00

11/28/00

11/28/00

11/28/00

11/28/00

11/28/00

11/28/00

11/28/00

11/28/00

11/28/00

11/28/00

11/28/00

11/28/00

11/28/00

11/28/00

11/28/00

11/28/00

11/28/00

11/28/00

11/28/00

(Col. 1) (Col. 2)

FOR:	NO. FILED	NO. EXTRA
BASIC FEE		
TOTAL CLAIMS	80 - 20	= *60
INDEP. CLAIMS	21 - 3	= *18
[X] MULTIPLE DEPENDENT CLAIM PRESENTED		

*If the difference in Col. 1 is less than 0, enter "0" in Col. 2.

SMALL ENTITY

RATE	FEE
x \$9.00 =	\$355.00
x \$40.00 =	\$720.00
+ \$135.00 =	
TOTAL	\$1,615.00

OTHER THAN SMALL ENTITY

RATE	FEE
x \$18.00 =	\$710.00
x \$80.00 =	
+ \$270.00 =	
TOTAL	

Please charge Deposit Account No. 20-1430 as follows:

- ☒ [X] Filing fee \$ \$1,615.00
☒ [X] Any additional fees associated with this paper or during the pendency of this application.
☐ [] The issue fee set in 37 CFR 1.18 at or before mailing of the Notice of Allowance, pursuant to 37 CFR 1.311(b)

- ☐ [] A check for \$ is enclosed.
☒ [X] 2 extra copies of this sheet are enclosed.

Respectfully submitted,
TOWNSEND and TOWNSEND and CREW LLP

Kevin Bastian
Reg No.: 34,774
Attorneys for Applicant

Telephone:
(415) 576-0200

Facsimile:
(415) 576-0300

PATENT APPLICATION

**POLYNUCLEOTIDES USEFUL FOR MODULATING
TRANSCRIPTION**

Inventor(s):

Koen Weterings, a citizen of the Netherlands, residing at Heidevenstraat 44
6533 TS, Nijmegen, Netherlands

Nestor R. Apuya, a citizen of the Phillipines, residing at 5396 Dobson Way
Culver City, California, 90230

Robert B. Goldberg, a citizen of the United States, residing at 21907 Canon
Drive, Topanga, California, 90290

Assignee:

THE REGENTS OF THE UNIVERSITY OF CALIFORNIA
1111 Franklin Street, 5th Floor
Oakland, CA 94607-5200

Entity: Small

POLYNUCLEOTIDES USEFUL FOR MODULATING TRANSCRIPTION

INTRODUCTION

5 In most higher plants, the first division of the zygote is asymmetric giving rise to two daughter cells differing in size and developmental fate (Goldberg, R. B., *et al.*, *Science*, 266:605-614 (1994); EMBRYOLOGY OF ANGIOSPERMS (Johri, B. M., ed., 1984); Kaplan, D. R., *et al.*, *Plant Cell*, 9:1903-1919 (1997); Laux, T., *et al.*, *Plant Cell*, 9:898-1000 (1997); EMBRYOGENESIS IN ANGIOSPERMS: A DEVELOPMENTAL AND EXPERIMENTAL STUDY (Raghavan, V., ed. 1986); West, M. A. L., *et al.*, *Plant Cell*, 5:1361-1369 (1993)). The small
10 terminal, or apical cell, is cytoplasmically dense and differentiates into the embryo proper containing one or two cotyledons and an axis with shoot and root meristems. By contrast, the large, highly-vacuolate basal cell differentiates into the hypophysis and suspensor. The hypophysis contributes to the formation of the root meristem within the embryo proper (van
15 Den Berg, C., *et al.*, *Planta Berlin*, 205:483-491 (1998)). The suspensor, on the other hand, is a terminally-differentiated embryonic region that anchors the embryo proper to the surrounding maternal tissue, serves as conduit for nutrients and growth regulators supporting embryo-proper development, and degenerates by the end of embryogenesis (Natesh, S., *et al.*, EMBRYOLOGY OF ANGIOSPERMS, (B. M. Johri, ed., 1984) 377-444; Schwartz, B. W., *et al.*,
20 CELLULAR AND MOLECULAR BIOLOGY OF PLANT SEED DEVELOPMENT, (B. Vasil, ed. 1997) 53-72.; Walthall, E. D., *et al.*, *Cell Differentiation*, 18:37-44 (1986); Yeung, E. C., *et al.*, *Can. J. Bot.*, 57:120-136 (1979); Yeung, E. C., *et al.*, *Plant Cell*, 5:1371-1381 (1993)).

The suspensor provides a novel opportunity to use molecular biology in order to understand how the zygote gives rise to daughter cells with distinct developmental fates. It
25 is highly differentiated and contains cells that are direct clonal descendents of the basal cell and, ultimately the basal region of the egg (Goldberg, R. B., *et al.*, *Science*, 266:605-614 (1994); Schwartz, B. W., *et al.*, CELLULAR AND MOLECULAR BIOLOGY OF PLANT SEED DEVELOPMENT, (B. Vasil, ed. 1997) 53-72; Yeung, E. C., *et al.*, *Plant Cell*, 5:1371-1381 (1993)). Fully developed *Arabidopsis* and tobacco suspensors, for example, are only three to
30 four cell divisions removed from the basal cell (Mansfield, S. G., *et al.*, *Canadian Journal of Botany*, 69:461-476 (1991); Soueges, R., *Compt. Rend. Acad. Sci. Paris*, 170:1125-1127 (1920)). It is possible, therefore, that the mechanisms regulating suspensor-specific gene expression are linked directly to the processes specifying the developmental fate of the basal

cell. An understanding how suspensor gene expression is regulated should provide insight into the molecular mechanisms specifying the fate of the basal cell.

Scarlet Runner Bean (*Phaseolus coccineus*) suspensors are approximately 100 times larger than the suspensors of either *Arabidopsis* or tobacco (Yeung, E. C., *et al.*, *Plant Cell*, 5:1371-1381 (1993)). Because of their large size, Scarlet Runner Bean suspensors can be microdissected from embryos during the early stages of embryogenesis (e.g., globular stage) and used for cDNA cloning, transcript profiling, and EST sequencing studies in order to identify and investigate suspensor-specific gene sets.

Control of the expression of genes in suspensor cells in plants is useful in the production of plants with a range of desired traits. For example, control of gene expression in suspensor cells can be used to make seedless fruit or to regulate embryo size or shape. These and other advantages are provided by the present application.

SUMMARY OF THE INVENTION

The present invention provides polynucleotides comprising a promoter control element, which comprises 1) a nucleotide sequence at least 50% identical to nucleotides 3324 to 3580 of SEQ ID NO:1, or 2) a nucleotide sequence that hybridizes to nucleotides 3324 to 3580 of SEQ ID NO:1 under a condition establishing a T_m of 20°C. In some embodiments, the isolated polynucleotides of the invention comprise a polynucleotide comprising 1) a nucleotide sequence at least 50% identical to SEQ ID NO:1, or 2) a nucleotide sequence that hybridizes to SEQ ID NO:1 under a condition establishing a T_m of 20°C. In some embodiments, the polynucleotides of the invention comprise nucleotides 3324 to 3580 of SEQ ID NO:1. In some embodiments, the polynucleotides of the invention modulate transcription in a cell. In some embodiments, the polynucleotides of the invention specifically modulate transcription in a plant suspensor cell and/or basal region of a plant embryo.

The present invention also provides expression cassettes comprising a promoter sequence comprising a nucleotide sequence at least 50% identical to nucleotides 3324 to 3580 of SEQ ID NO:1 and a promoter polynucleotide with at least basal promoter activity, which promoter polynucleotide is operably linked to a heterologous polynucleotide, wherein when the expression cassette is inserted into a plant, the heterologous polynucleotide is specifically expressed in a suspensor cell and/or basal region of a plant embryo.

The present invention also provides polynucleotides comprising 1) a nucleotide sequence at least 50% identical to SEQ ID NO:1 or nucleotides 1-3154 or SEQ ID

NO:6, or 2) a nucleotide sequence that hybridizes to SEQ ID NO:1 or nucleotides 1-3154 or SEQ ID NO:6 under a condition establishing a T_m of 20°C. In some embodiments, the isolated polynucleotides further comprise a G654 or C541 polynucleotide operably linked to the promoter. Examples of such polynucleotides include SEQ ID NO:2 and SEQ ID NO:6.

Alternatively, the invention provides for a heterologous polynucleotide operably linked to a promoter. In some embodiments, the polynucleotides of the invention comprise a promoter that modulates transcription in a cell. In some embodiments, the polynucleotides of the invention specifically modulate transcription in a plant suspensor cell and/or basal region of a plant embryo.

The present invention also provides for vectors comprising the above-referenced promoter operably linked to a heterologous polynucleotide. For instance, in some embodiments, the promoter is SEQ ID NO:1 or nucleotides 1 to 3154 of SEQ ID NO:6.

The present invention also provides for a host cell comprising the above-referenced promoters. For instance, in some embodiments, the promoter is SEQ ID NO:1 or nucleotides 1 to 3154 of SEQ ID NO:6. In some embodiments, the host cell comprises a vector comprising the promoters of the invention operably linked to a heterologous nucleic acid.

The invention also provides for plants comprising a promoter comprising 1) a nucleotide sequence at least 50% identical to SEQ ID NO:1 or nucleotides 1-3154 or SEQ ID NO:6, or 2) a nucleotide sequence that hybridizes to SEQ ID NO:1 or nucleotides 1-3154 or SEQ ID NO:6 under a condition establishing a T_m of 20°C, wherein the promoter is operably linked to a heterologous polynucleotide. For instance, in some embodiments, the promoter is SEQ ID NO:1 or nucleotides 1 to 3154 of SEQ ID NO:6. In some embodiments, the plant comprises a vector comprising the promoters of the invention operably linked to a heterologous nucleic acid.

The invention also provides methods of modulating transcription in a suspensor cell comprising introducing into the plant an expression cassette comprising a promoter comprising 1) a nucleotide sequence at least 50% identical to SEQ ID NO:1 or nucleotides 1-3154 or SEQ ID NO:6, or 2) a nucleotide sequence that hybridizes to SEQ ID NO:1 or nucleotides 1-3154 or SEQ ID NO:6 under a condition establishing a T_m of 20°C. For instance, in some embodiments, the promoter is SEQ ID NO:1 or nucleotides 1 to 3154 of SEQ ID NO:6. In some embodiments, a G654 or C541 polynucleotide is operably linked to the promoter. In some embodiments, the promoter is operably linked to a heterologous

polynucleotide. In some embodiments, the promoter is operably linked to the heterologous polynucleotide in an antisense orientation.

The present invention also provides isolated nucleic acids comprising a polynucleotide sequence, or complement thereof, encoding a G654 polypeptide at least 50% identical to SEQ ID NO:3 or a C541 polypeptide at least 50% identical to SEQ ID NO:7. In some embodiments, the G654 polypeptide is SEQ ID NO:3. In some embodiments, the C541 polypeptide is SEQ ID NO:7. In some embodiments, the polynucleotide is operably linked to a promoter. For example, the promoter can be a constitutive promoter. In some embodiments, the polynucleotide is linked to the promoter in an antisense orientation.

The invention also provides an expression cassette comprising a promoter operably linked to a heterologous polynucleotide, or complement thereof, encoding a G654 or C541 polypeptide at least 50% identical to SEQ ID NO:3 or SEQ ID NO:7, respectively. In some embodiments, the G654 polynucleotide comprises nucleotides 4242 to 4901 of SEQ ID NO:2. In some embodiments, the C541 polynucleotide comprises nucleotides 3155 to 3552 of SEQ ID NO:6. In some embodiments, the polynucleotide is operably linked to a promoter. For example, the promoter can be a constitutive promoter. In some embodiments, the polynucleotide is linked to the promoter in an antisense orientation.

The present invention also provides for host cells and transgenic plants comprising an exogenous nucleic acid comprising a polynucleotide, or complement thereof, encoding a G654 polypeptide at least 50% identical to SEQ ID NO:3 or a C541 polypeptide at least 50% identical to SEQ ID NO:7.

The present invention also provides for isolated polypeptides comprising an amino acid sequence at least 50% identical to SEQ ID NO:3 or SEQ ID NO:7. The invention also provides for antibodies capable of binding the isolated polypeptides.

The invention also provides methods of introducing an isolated polynucleotide into a host cell. The method comprises providing an isolated polynucleotide that comprises 1) a nucleotide sequence at least 50% identical to SEQ ID NO:1 or nucleotides 1-3154 or SEQ ID NO:6, or 2) a nucleotide sequence that hybridizes to SEQ ID NO:1 or nucleotides 1-3154 or SEQ ID NO:6 under a condition establishing a T_m of 20°C. The method also provides contacting the polynucleotide with the host cell under conditions that permit insertion of the polynucleotide into the host cell.

The invention also provides methods of detecting a polynucleotide in a sample. The methods comprise providing a polynucleotide that comprises 1) a nucleotide sequence at least 50% identical to SEQ ID NO:1 or nucleotides 1-3154 or SEQ ID NO:6, or

2) a nucleotide sequence that hybridizes to SEQ ID NO:1 or nucleotides 1-3154 or SEQ ID NO:6 under a condition establishing a T_m of 20°C. The method also comprises contacting the polynucleotide with a sample under conditions that permit a comparison of the sequence the polynucleotide with a sequence of DNA in the sample and analyzing the result of the comparison. In some embodiments, the polynucleotide and the sample are contacted under conditions that permit formation of a duplex between complementary nucleic acid sequences.

The present invention also provides polynucleotides comprising SEQ ID NO:10 or SEQ ID NO:11. In some embodiments, the polynucleotides of the invention comprise an expression cassette comprising a promoter sequence comprising SEQ ID NO:10 or SEQ ID NO:11 and a promoter polynucleotide with at least basal promoter activity, which promoter polynucleotide is operably linked to a heterologous polynucleotide, wherein when the expression cassette is inserted into a plant, the heterologous polynucleotide is specifically expressed in a suspensor cell and/or basal region of a plant embryo.

The invention also provides methods of constructing a promoter that specifically induces transcription in a plant suspensor cell and/or basal region of a plant embryo, the method comprising (i) providing a promoter polynucleotide capable of at least basal promoter activity in a plant; (ii) inserting a nucleic acid comprising SEQ ID NO: 10 or SEQ ID NO:11 within or adjoining the promoter polynucleotide, thereby constructing a test promoter; and (iii) assaying the test promoter to determine whether the test promoter specifically initiates transcription in a suspensor cell and/or basal region of a plant embryo. In some embodiments, the nucleic acid is SEQ ID NO:10 or SEQ ID NO:11.

DEFINITIONS

The term “basal promoter activity” refers to the ability of a polynucleotide sequence to initiate transcription of an operably linked polynucleotide. Typically, basal activity will provide a low level of constitutive expression that is not inducible under most conditions or that is not cell-specific under most conditions. A basal promoter typically comprises a TATA box and transcriptional start sequence, but does not contain additional stimulatory and repressive elements. An exemplary plant minimal promoter is positions -50 to +8 of the 35S CaMV promoter.

The term “basal region of a plant embryo” refers to the basal cell, i.e., the cell of a two-celled embryo that contacts the suspensor cell. The “basal region” also encompasses derivative or descendent cells of the basal cell.

The term "chimeric" is used to describe polynucleotides or genes, as defined *supra*, or constructs wherein at least two of the elements of the polynucleotide or gene or construct, such as the promoter and the polynucleotide to be transcribed and/or other regulatory sequences and/or filler sequences and/or complements thereof, are heterologous to each other.

Promoters referred to herein as "constitutive promoters" actively promote transcription under most, but not necessarily all, environmental conditions and states of development or cell differentiation. Examples of constitutive promoters include the cauliflower mosaic virus (CaMV) 35S transcript initiation region and the 1' or 2' promoter derived from T-DNA of *Agrobacterium tumefaciens*, and other transcription initiation regions from various plant genes, such as the maize ubiquitin-1 promoter, known to those of skill.

"Domains" are fingerprints or signatures that can be used to characterize protein families and/or parts of proteins. Such fingerprints or signatures can comprise conserved (1) primary sequence, (2) secondary structure, and/or (3) three-dimensional conformation. A similar analysis can be applied to polynucleotides. Generally, each domain has been associated with either a conserved primary sequence or a sequence motif. Generally these conserved primary sequence motifs have been correlated with specific *in vitro* and/or *in vivo* activities. A domain can be any length, including the entirety of the polynucleotide to be transcribed. Examples of domains include, without limitation, AP2, helicase, homeobox, zinc finger, etc.

The term "endogenous," within the context of the current invention refers to any polynucleotide, polypeptide or protein sequence which is a natural part of a cell or organisms regenerated from said cell.

An "enhancer" is a DNA regulatory element that can increase the steady state level of a transcript, usually by increasing the rate of transcription initiation. Enhancers usually exert their effect regardless of the distance, upstream or downstream location, or orientation of the enhancer relative to the start site of transcription. In contrast, a "suppressor" is a corresponding DNA regulatory element that decreases the steady state level of a transcript, again usually by affecting the rate of transcription initiation. The essential activity of enhancer and suppressor elements is to bind a protein factor(s). Such binding can be assayed, for example, by methods described below. The binding is typically in a manner that influences the steady state level of a transcript in a cell or in an *in vitro* transcription extract.

As referred to within, "exogenous" is any polynucleotide, polypeptide or protein sequence, whether chimeric or not, that is introduced into the genome of a host cell or organism regenerated from said host cell by any means other than by a sexual cross.

Examples of means by which this can be accomplished are described below, and include

- 5 Agrobacterium-mediated transformation (of dicots - e.g. Salomon *et al.* *EMBO J.* 3:141 (1984); Herrera-Estrella *et al.* *EMBO J.* 2:987 (1983); of monocots, representative papers are those by Escudero *et al.*, *Plant J.* 10:355 (1996), Ishida *et al.*, *Nature Biotechnology* 14:745 (1996), May *et al.*, *Bio/Technology* 13:486 (1995)), biolistic methods (Armaleo *et al.*, *Current Genetics* 17:97 1990)), electroporation, in planta techniques, and the like. Such a
- 10 plant containing the exogenous nucleic acid is referred to here as a T_0 for the primary transgenic plant and T_1 for the first generation. The term "exogenous" as used herein is also intended to encompass inserting a naturally found element into a non-naturally found location.

- 15 An "expression cassette" refers to a nucleic acid construct, which when introduced into a host cell, results in transcription and/or translation of an RNA or polypeptide, respectively. Antisense or sense constructs that are not or cannot be translated are expressly included by this definition.

- The term "gene," as used in the context of the current invention, encompasses all regulatory and coding sequence contiguously associated with a single hereditary unit with a genetic function (see Figure 1). Genes can include non-coding sequences that modulate the genetic function that include, but are not limited to, those that specify polyadenylation, transcriptional regulation, DNA conformation, chromatin conformation, extent and position of base methylation and binding sites of proteins that control all of these. Genes encoding proteins are comprised of "exons" (coding sequences), which may be interrupted by "introns"
- 25 (non-coding sequences). In some instances complexes of a plurality of protein or nucleic acids or other molecules, or of any two of the above, may be required for a gene's function. On the other hand, a gene's genetic function may require only RNA expression or protein production, or may only require binding of proteins and/or nucleic acids without associated expression. In certain cases, genes adjacent to one another may share sequence in such a
- 30 way that one gene will overlap the other. A gene can be found within the genome of an organism, in an artificial chromosome, in a plasmid, in any other sort of vector, or as a separate isolated entity.

A "G564 polynucleotide" is a nucleic acid sequence or subsequence that encodes a polypeptide with substantial identity (as defined below) to SEQ ID NO:3 or SEQ

ID NO:5. Alternatively, a G564 polynucleotide includes polynucleotide sequences that are substantially identical to SEQ ID NO:1, SEQ ID NO:2, or SEQ ID NO:4 or that hybridize to SEQ ID NO:1, SEQ ID NO:2, or SEQ ID NO:4 under defined conditions.

A "promoter from a G564 gene" or "G564 promoter" will typically be about 500 to about 5000 nucleotides in length, usually from about 2500 to 4000. Exemplary promoter sequences are shown as SEQ ID NO:1 or nucleotides 1-4242 of SEQ ID NO:2. A G564 promoter can also be identified by its ability to direct expression in suspensor cells.

"Increased or enhanced G564 activity or expression of the *G564* gene" refers to an augmented change in G564 activity. Examples of such increased activity or expression include the following. G564 activity or expression of the *G564* gene is increased above the level of that in wild-type, non-transgenic control plants (i.e. the quantity of G564 activity or expression of the *G564* gene is increased). G564 activity or expression of the *G564* gene is in an organ, tissue or cell where it is not normally detected in wild-type, non-transgenic control plants (i.e. spatial distribution of G564 activity or expression of the *G564* gene is increased). G564 activity or expression is increased when G564 activity or expression of the *G564* gene is present in an organ, tissue or cell for a longer period than in a wild-type, non-transgenic controls (i.e. duration of G564 activity or expression of the *G564* gene is increased).

A "C541 polynucleotide" is a nucleic acid sequence or subsequence that encodes a polypeptide with substantial identity (as defined below) to SEQ ID NO:7 or SEQ ID NO:9. Alternatively, a C541 polynucleotide includes polynucleotide sequences that are substantially identical to SEQ ID NO:6, or SEQ ID NO:8 or that hybridize to SEQ ID NO:6 or SEQ ID NO:8 under defined conditions.

A "promoter from a C541 gene" or "C541 promoter" will typically be about 500 to about 5000 nucleotides in length, usually from about 2500 to 4000. Exemplary promoter sequences are shown as nucleotides 1-3154 of SEQ ID NO:6 or nucleotides 1-1609 of SEQ ID NO:8. A C541 promoter can also be identified by its ability to direct expression in suspensor cells.

"Increased or enhanced C541 activity or expression of the *C541* gene" refers to an augmented change in C541 activity. Examples of such increased activity or expression include the following. C541 activity or expression of the *C541* gene is increased above the level of that in wild-type, non-transgenic control plants (i.e. the quantity of C541 activity or expression of the *C541* gene is increased). C541 activity or expression of the *C541* gene is in an organ, tissue or cell where it is not normally detected in wild-type, non-transgenic control

plants (i.e. spatial distribution of C541 activity or expression of the *C541* gene is increased). C541 activity or expression is increased when C541 activity or expression of the *C541* gene is present in an organ, tissue or cell for a longer period than in a wild-type, non-transgenic controls (i.e. duration of C541 activity or expression of the *C541* gene is increased).

“Inserting a first polynucleotide within or adjoining” a second polynucleotide is discussed below. “Inserting a first polynucleotide within a second polynucleotide” refers to manipulating or constructing a first and second polynucleotide such that the first polynucleotide interrupts the second polynucleotide (e.g., the first polynucleotide is inserted between the 5’ end and the 3’ end of the second polynucleotide). “Inserting a first polynucleotide adjoining a second polynucleotide” refers to manipulating or constructing a polynucleotide such that the first and second polynucleotides are linked, i.e., the first polynucleotide is adjacent to the second polynucleotide. Of course, one of skill in the art will recognize that the first and the second polynucleotide can be linked in either orientations (e.g., 1→2 or 2→1) or can be linked via a polynucleotide spacer. In the context of promoter sequences, polynucleotides comprising TATA boxes and other basal promoter elements are typically at the 3’ end of a promoter and can be operably linked at their 3’ end to a polynucleotide that is to be transcribed. Moreover, in some embodiments, promoter sequences comprise fewer than 10,000 base pairs, more typically fewer than 5,000 base pairs, sometimes fewer than 3,000, 1,000 or 500 base pairs. However, as noted elsewhere within this application, enhancer elements can function independently of their distance from a basal promoter. Therefore, in some embodiments, the active elements of a promoter can be separated by more than 10,000 base pairs.

“Heterologous sequences” are those that are not operatively linked or are not contiguous to each other in nature. For example, a promoter from corn is considered heterologous to an Arabidopsis coding region sequence. Also, a promoter from a gene encoding a growth factor from maize is considered heterologous to a sequence encoding the maize receptor for the growth factor. Regulatory element sequences, such as UTRs or 3’ end termination sequences that do not originate in nature from the same gene as the coding sequence originates from, are considered heterologous to said coding sequence. Elements operatively linked in nature and contiguous to each other are not heterologous to each other.

In the current invention, a “homologous” gene or polynucleotide or polypeptide refers to a gene or polynucleotide or polypeptide that shares sequence similarity with the gene or polynucleotide or polypeptide of interest. This similarity may be in only a fragment of the sequence and often represents a functional domain such as, examples

including without limitation a DNA binding domain or a domain with tyrosine kinase activity. The functional activities of homologous polynucleotide are not necessarily the same.

An "inducible promoter" in the context of the current invention refers to a promoter, the activity of which is influenced by certain conditions, such as light, temperature, chemical concentration, protein concentration, conditions in an organism, cell, or organelle, etc. A typical example of an inducible promoter, which can be utilized with the polynucleotides of the present invention, is PARSK1, the promoter from an Arabidopsis gene encoding a serine-threonine kinase enzyme, and which promoter is induced by dehydration, abscisic acid and sodium chloride (Wang and Goodman, Plant J. 8:37 (1995)). Examples of environmental conditions that may affect transcription by inducible promoters include anaerobic conditions, elevated temperature, the presence or absence of a nutrient or other chemical compound or the presence of light.

As used herein, the phrase "modulate transcription" describes the biological activity of a promoter sequence or promoter control element. Such modulation includes, without limitation, includes up- and down-regulation of initiation of transcription, rate of transcription, and/or transcription levels.

In the current invention, "mutant" refers to a heritable change in nucleotide sequence at a specific location. Mutant genes of the current invention may or may not have an associated identifiable phenotype.

An "operable linkage" is a linkage in which a promoter sequence or promoter control element is connected to a polynucleotide sequence (or sequences) in such a way as to place transcription of the polynucleotide sequence under the influence or control of the promoter or promoter control element. Two DNA sequences (such as a polynucleotide to be transcribed and a promoter sequence linked to the 5' end of the polynucleotide to be transcribed) are said to be operably linked if induction of promoter function results in the transcription of mRNA encoding the polynucleotide and if the nature of the linkage between the two DNA sequences does not (1) result in the introduction of a frame-shift mutation, (2) interfere with the ability of the promoter sequence to direct the expression of the protein, antisense RNA or ribozyme, or (3) interfere with the ability of the DNA template to be transcribed. Thus, a promoter sequence would be operably linked to a polynucleotide sequence if the promoter was capable of effecting transcription of that polynucleotide sequence.

"Orthologous" is a term used herein to describe a relationship between two or more polynucleotides or proteins. Two polynucleotides or proteins are "orthologous" to one

another if they serve a similar function in different organisms. In general, orthologous polynucleotides or proteins will have similar catalytic functions (when they encode enzymes) or will serve similar structural functions (when they encode proteins or RNA that form part of the ultrastructure of a cell).

"Percentage of sequence identity," as used herein, is determined by comparing two optimally aligned sequences over a comparison window, where the fragment of the polynucleotide or amino acid sequence in the comparison window may comprise additions or deletions (e.g., gaps or overhangs) as compared to the reference sequence (which does not comprise additions or deletions) for optimal alignment of the two sequences. The percentage is calculated by determining the number of positions at which the identical nucleic acid base or amino acid residue occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the window of comparison and multiplying the result by 100 to yield the percentage of sequence identity. Optimal alignment of sequences for comparison may be conducted by the local homology algorithm of Smith and Waterman *Add. APL. Math.* 2:482 (1981), by the homology alignment algorithm of Needleman and Wunsch *J. Mol. Biol.* 48:443 (1970), by the search for similarity method of Pearson and Lipman *Proc. Natl. Acad. Sci. (USA)* 85: 2444 (1988), by computerized implementations of these algorithms (GAP, BESTFIT, BLAST, PASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group (GCG), 575 Science Dr., Madison, WI), or by inspection. Given that two sequences have been identified for comparison, GAP and BESTFIT are preferably employed to determine their optimal alignment. Typically, the default values of 5.00 for gap weight and 0.30 for gap weight length are used.

A "plant promoter" is a promoter capable of initiating transcription in plant cells and can modulate transcription of a polynucleotide. Such promoters need not be of plant origin. For example, promoters derived from plant viruses, such as the CaMV35S promoter or from *Agrobacterium tumefaciens* such as the T-DNA promoters, can be plant promoters. A typical example of a plant promoter of plant origin is the maize ubiquitin-1 (ubi-1) promoter known to those of skill.

The term "plant tissue" includes differentiated and undifferentiated tissues or plants, including but not limited to roots, stems, shoots, cotyledons, epicotyl, hypocotyl, leaves, pollen, seeds, tumor tissue and various forms of cells and culture such as single cells, protoplast, embryos, basal and apical cells, suspensor cells and callus tissue. The plant tissue may be in plants or in organ, tissue or cell culture.

“Preferential transcription” is defined as transcription that occurs in a particular pattern of cell types or developmental times or in response to specific stimuli or combination thereof. Non-limiting examples of preferential transcription include: high transcript levels of a desired sequence in suspensor cells; detectable transcript levels of a desired sequence in certain cell types during embryogenesis; and low transcript levels of a desired sequence under drought conditions. Such preferential transcription can be determined by measuring initiation, rate, and/or levels of transcription.

A “promoter” is a DNA sequence that directs the transcription of a polynucleotide. Typically a promoter is located in the 5' region of a polynucleotide to be transcribed, proximal to the transcriptional start site of such polynucleotide. More typically, promoters are defined as the region upstream of the first exon; more typically, as a region upstream of the first of multiple transcription start sites; more typically, as the region downstream of the preceding gene and upstream of the first of multiple transcription start sites; more typically, the region downstream of the polyA signal and upstream of the first of multiple transcription start sites; even more typically, about 3,000 nucleotides upstream of the ATG of the first exon; even more typically, 2,000 nucleotides upstream of the first of multiple transcription start sites. The promoters of the invention comprise at least a core promoter as defined below. Additionally, the promoter may also include at least one control element such as an upstream element. Such elements include UARs and optionally, other DNA sequences that affect transcription of a polynucleotide such as a synthetic upstream element.

The term “promoter control element” as used herein describes elements that influence the activity of the promoter. Promoter control elements include transcriptional regulatory sequence determinants such as, but not limited to, enhancers, scaffold/matrix attachment regions, TATA boxes, transcription start locus control regions, UARs, URRs, other transcription factor binding sites and inverted repeats. Exemplary promoter control elements include, e.g., SEQ ID NO:10 and SEQ ID NO:11.

The term “public sequence,” as used in the context of the instant application, refers to any sequence that has been deposited in a publicly accessible database prior to the filing date of the present application. This term encompasses both amino acid and nucleotide sequences. Such sequences are publicly accessible, for example, on the BLAST databases on the NCBI FTP web site (accessible at ncbi.nlm.gov/blast). The database at the NCBI GTP site utilizes “gi” numbers assigned by NCBI as a unique identifier for each sequence in the databases, thereby providing a non-redundant database for sequence from various databases,

including GenBank, EMBL, DBBJ, (DNA Database of Japan) and PDB (Brookhaven Protein Data Bank).

The term "regulatory sequence," as used in the current invention, refers to any nucleotide sequence that influences transcription or translation initiation and rate, or stability and/or mobility of a transcript or polypeptide product. Regulatory sequences include, but are not limited to, promoters, promoter control elements, protein binding sequences, 5' and 3' UTRs, transcriptional start sites, termination sequences, polyadenylation sequences, introns, certain sequences within amino acid coding sequences such as secretory signals, protease cleavage sites, etc.

"Related sequences" refer to either a polypeptide or a nucleotide sequence that exhibits some degree of sequence similarity with a reference sequence.

The term "substantial identity" of polynucleotide sequences means that a polynucleotide comprises a sequence that has at least 25% sequence identity. Alternatively, percent identity can be any integer from 25% to 100%. More preferred embodiments include at least: 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, or 99%. compared to a reference sequence using the programs described herein; preferably BLAST using standard parameters, as described below. For instance, promoter sequences of the invention sequences of the invention include nucleic acid sequences that have substantial identity to SEQ ID NO:1 or other sequences of the invention such as nucleotides 1-4582 of SEQ ID NO:4, nucleotides 1-3154 of SEQ ID NO:6 or nucleotides 1-1609 of SEQ ID NO:8. One of skill will recognize that these values can be appropriately adjusted to determine corresponding identity of proteins encoded by two nucleotide sequences by taking into account codon degeneracy, amino acid similarity, reading frame positioning and the like. Substantial identity of amino acid sequences for these purposes normally means sequence identity of at least 40%. Preferred percent identity of polypeptides can be any integer from 40% to 100%. More preferred embodiments include at least 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, or 99%. Most preferred embodiments include 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74% and 75%. Polypeptides which are "substantially similar" share sequences as noted above except that residue positions which are not identical may differ by conservative amino acid changes. Conservative amino acid substitutions refer to the interchangeability of residues having similar side chains. For example, a group of amino acids having aliphatic side chains is glycine, alanine, valine, leucine, and isoleucine; a group of amino acids having aliphatic-hydroxyl side chains is serine and threonine; a group of amino acids having amide-containing side chains is asparagine and glutamine; a group of

amino acids having aromatic side chains is phenylalanine, tyrosine, and tryptophan; a group of amino acids having basic side chains is lysine, arginine, and histidine; and a group of amino acids having sulfur-containing side chains is cysteine and methionine. Preferred conservative amino acids substitution groups are: valine-leucine-isoleucine, phenylalanine-tyrosine, lysine-arginine, alanine-valine, aspartic acid-glutamic acid, and asparagine-glutamine.

In the context of the current invention, "specific promoters" refers to a subset of promoters that have a high preference for modulating transcript levels in a specific tissue or organ or cell and/or at a specific time during development of an organism, i.e., that are "specifically initiated" or "specifically modulated" in a specific tissue or at a specific developmental time. By "high preference" is meant at least 3-fold, preferably 5-fold, more preferably at least 10-fold still more preferably at least 20-fold, 50-fold or 100-fold increase in transcript levels under the specific condition and/or a specific tissue over the transcription under any other reference condition and/or in any other reference tissue considered. Examples of tissue-specific promoters under developmental control include promoters that initiate transcription only in certain tissues or organs, such as suspensor cell, root, ovule, fruit, seeds, or flowers. See also "Preferential transcription".

"Stringency" as used herein is a function of probe length, probe composition (G + C content), and salt concentration, organic solvent concentration, and temperature of hybridization or wash conditions. Stringency is typically compared by the parameter T_m , which is the temperature at which 50% of the complementary molecules in the hybridization are hybridized, in terms of a temperature differential from T_m . High stringency conditions are those providing a condition of T_m minus 5°C to T_m minus 10°C. Medium or moderate stringency conditions are those providing T_m -minus 20°C to T_m minus 29°C. Low stringency conditions are those providing a condition of T_m minus 40°C to T_m minus 48°C. The relationship of hybridization conditions to T_m (in °C) is expressed in the mathematical equation

$$T_m = 81.5 - 16.6(\log_{10}[Na^+]) + 0.41(\%G+C) - (600/N) \quad (1)$$

where N is the length of the probe. This equation works well for probes 14 to 70 nucleotides in length that are identical to the target sequence. The equation below for T_m of DNA-DNA hybrids is useful for probes in the range of 50 to greater than 500 nucleotides, and for conditions that include an organic solvent (formamide).

$$T_m = 81.5 + 16.6 \log \{[Na^+]/(1 + 0.7[Na^+])\} + 0.41(\%G+C) - 500/L - 0.63(\%\text{formamide}) \quad (2)$$

where L is the length of the probe in the hybrid. (P. Tijessen, "Hybridization with Nucleic Acid Probes" in LABORATORY TECHNIQUES IN BIOCHEMISTRY AND MOLECULAR BIOLOGY, (P.C. van der Vliet, ed. 1993)). The T_m of equation (2) is affected by the nature of the hybrid; for DNA-RNA hybrids T_m is 10-15°C higher than calculated, for RNA-RNA hybrids T_m is 20-25°C higher. Because the T_m decreases about 1°C for each 1% decrease in homology when a long probe is used (Bonner *et al.*, *J. Mol. Biol.* 81:123 (1973)), stringency conditions can be adjusted to favor detection of identical genes or related family members.

Equation (2) is derived assuming equilibrium and therefore, hybridizations according to the present invention are most preferably performed under conditions of probe excess and for sufficient time to achieve equilibrium. The time required to reach equilibrium can be shortened by inclusion of a hybridization accelerator such as dextran sulfate or another high volume polymer in the hybridization buffer.

Stringency can be controlled during the hybridization reaction or after hybridization has occurred by altering the salt and temperature conditions of the wash solutions used. The formulas shown above are equally valid when used to compute the stringency of a wash solution. Preferred wash solution stringencies lie within the ranges stated above; high stringency is 5-8°C below T_m , medium or moderate stringency is 26-29°C below T_m and low stringency is 45-48°C below T_m . Hybridization conditions include those in which the salt concentration is less than about 1.0 M sodium ion, typically about 0.1 to 1.0 M sodium ion concentration (or other salts) at pH 7.0 to 8.3 and the temperature is at least about 65°C or about 60°C, more preferably 55°C and more preferably 50°C.

A composition containing A is "substantially free of" B when at least 85% by weight of the total A+B in the composition is A. Preferably, A comprises at least about 90% by weight of the total of A+B in the composition, more preferably at least about 95% or even 99% by weight. For example, a plant gene can be substantially free of other plant genes. Other examples include, but are not limited to, ligands substantially free of receptors (and vice versa), a growth factor substantially free of other growth factors and a transcription binding factor substantially free of nucleic acids.

"TATA to start" shall mean the distance, in number of nucleotides, between the primary TATA motif and the start of transcription.

A “transgenic plant” is a plant having one or more plant cells that contain at least one exogenous polynucleotide introduced by recombinant nucleic acid methods.

In the context of the present invention, a “translational start site” is usually an ATG or AUG in a transcript, often the first ATG or AUG. A single protein encoding transcript, however, may have multiple translational start sites.

“Transcription start site” is used in the current invention to describe the point at which transcription is initiated. This point is typically located about 25 nucleotides downstream from a TFIID binding site, such as a TATA box. Transcription can initiate at one or more sites within the gene, and a single polynucleotide to be transcribed may have multiple transcriptional start sites, some of which may be specific for transcription in a particular cell-type or tissue or organ. “+1” is stated relative to the transcription start site and indicates the first nucleotide in a transcript.

An “Upstream Activating Region” or “UAR” is a position or orientation dependent nucleic acid element that primarily directs tissue, organ, cell type, or environmental regulation of transcript level, usually by affecting the rate of transcription initiation. Corresponding DNA elements that have a transcription inhibitory effect are called herein “Upstream Repressor Regions” or “URR”s. The essential activity of these elements is to bind a protein factor. Such binding can be assayed by methods described below. The binding is typically in a manner that influences the steady state level of a transcript in a cell or in vitro transcription extract.

An “untranslated region” or “UTR” is any contiguous series of nucleotide bases that is transcribed, but is not translated. A 5’ UTR lies between the start site of the transcript and the translation initiation codon and includes the +1 nucleotide. A 3’ UTR lies between the translation termination codon and the end of the transcript. UTRs can have particular functions such as increasing mRNA message stability or translation attenuation. Examples of 3’ UTRs include, but are not limited to polyadenylation signals and transcription termination sequences.

The term “variant” is used herein to denote a polypeptide or protein or polynucleotide molecule that differs from others of its kind in some way. For example, polypeptide and protein variants can consist of changes in amino acid sequence and/or charge and/or post-translational modifications (such as glycosylation, etc). It will be understood that there may be sequence variations within sequence or fragments used or disclosed in this application. Preferably, variants will be such that the sequences have at least 80%, preferably

at least 90%, 95, 97, 98, or 99% sequence identity. Variants preferably measure the primary biological function of the native polypeptide or protein or polynucleotide.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 displays a schematic representation of a gene.

Figure 2 displays the nucleotide sequence of genomic DNA comprising the G564 coding sequence and promoter region from Scarlet Runner Bean (*Phaseolus coccineus*). The ATG start codon is displayed in bold and underlined nucleotides indicates intron sequences.

Figure 3 displays the nucleotide sequence of genomic DNA comprising the G564 coding sequence and promoter region from *Arabidopsis thaliana*. The ATG start codon is displayed in bold and underlined nucleotides indicates intron sequences.

Figure 4 displays the nucleotide sequence of genomic DNA comprising the C541 coding sequence and promoter region from Scarlet Runner Bean (*Phaseolus coccineus*). The ATG start codon is displayed in bold and underlined nucleotides indicates intron sequences.

Figure 5 displays the nucleotide sequence of genomic DNA comprising the C541 coding sequence and promoter region from *Arabidopsis thaliana*. The ATG start codon is displayed in bold and underlined nucleotides indicates intron sequences.

Figure 6 is a schematic representation of a deletion analysis of the Scarlet Runner Bean G564 promoter. Suspensor-specific GUS expression was observed in all constructs except the shortest (deleted from the 5' end to position -662). This figure demonstrates that a suspensor-specific cis-acting sequence is located between positions -921 and -662 (corresponding to nucleotides 3324-3580 of SEQ ID NO:2).

Figure 7 is a schematic representation of a series of promoter fragments from the Scarlet Runner Bean G564 promoter region fused to a minimal 35S promoter and GUS gene.

Figure 8 identifies a number of promoter control elements found within sequences -921 to -662 of Figure 1.

DETAILED DESCRIPTION OF THE INVENTION

A. INTRODUCTION

The present invention provides the identification of two Scarlet Runner Bean mRNAs, designated as C541 and G564, that accumulate specifically within the suspensor of

globular-stage embryos. At the pre-globular, or four-cell stage, both C541 and G564 mRNAs are present in the two basal cells, but are absent from the two embryo-proper cells. Expression analysis of a chimeric *G564/GUS* gene in transgenic tobacco embryos showed that the *G564* promoter is active specifically within the suspensor during early embryo development.

The present invention provides polynucleotides comprising promoters and promoter control elements which are capable of modulating transcription.

Such promoters and promoter control elements can be used in combination with native or heterologous promoter fragments, control elements or other regulatory sequences to modulate transcription and/or translation.

Specifically, promoters and control elements of the invention can be used to modulate transcription of a desired polynucleotide, which includes without limitation:

- (a) antisense;
- (b) ribozymes;
- (c) coding sequences; or
- (d) fragments thereof.

The promoter also can modulate transcription in a host genome in cis- or in trans-.

In an organism, such as a plant, the promoters and promoter control elements of the instant invention are useful to produce preferential transcription which results in a desired pattern of transcript levels in a particular cells, tissues, or organs, or under particular conditions.

The present invention also provides new suspensor-specific genes useful in genetically engineering plants. Suspensor-specific promoter sequences from the genes of the invention can be used, for instance, to ablate embryos to make seedless fruit, e.g., by expressing gene products toxic to the suspensor and/or surrounding cells such as the embryo itself. The suspensor-specific promoters can also be operably linked to growth regulator genes, such as gene products regulating gibberellin production, thereby modulating embryo size, shape and/or rate of development.

B. Identifying and Isolating Promoter Sequences or Structural Polynucleotides of the Invention

The exemplary promoters and promoter control elements of the present invention (e.g., SEQ ID NO:1 and fragments thereof) were identified from Scarlet Runner

bean (*Phaseolus coccineus*). Additional promoter sequences can be identified as described below. SEQ ID NO:1 and SEQ ID NO:2 includes a promoter region of approximately 4200 base pairs upstream of the ATG start codon.

In addition, the coding sequence of a suspensor-specific gene, designated G564, was identified (e.g., nucleotides 4242 to 4349 and 4513 to 4901 of SEQ ID NO:2). The genus of G564 nucleic acid sequences of the invention includes genes and gene products identified and characterized by analysis using the sequences nucleic acid sequences, nucleotides 4242 to 4349 and 4513 to 4901 of SEQ ID NO:2, as well as nucleotides 4242 to 6986 of SEQ ID NO:2, and protein sequences, including SEQ ID NO:3. G564 sequences of the invention include polypeptide sequences having substantial identity to SEQ ID NO:3. The orthologous *Arabidopsis* G564 polynucleotide was also identified (SEQ ID NO:4).

In addition, a polynucleotide designated C541 was also isolated from Scarlet Runner Bean (SEQ ID NO:6). The orthologous *Arabidopsis* C541 sequence is displayed as SEQ ID NO:8. The respective amino acid sequences encoded by the bean and *Arabidopsis* polynucleotides are SEQ ID NO:7 and SEQ ID NO:9.

The promoter sequences of the invention are useful to modulate transcription of polynucleotides. For example, promoter sequences can be operably linked to a polynucleotide of interest to modulate expression of that polynucleotide in desired tissues. Desired tissues for polynucleotide expression include, e.g., suspensor cells and/or the basal region of a plant embryo, the embryo root meristem as well as the plant root tip and plant root meristem.

Alternatively, promoter sequences of the invention, e.g., SEQ ID NO:1, are useful to modulate expression of polynucleotides in desired plant tissues. In addition, the promoter sequences of the invention can also be introduced into a cell in multiple copies, thereby competing with endogenous promoter sequences for transcription factors. By removing some or all of the transcription factors available for a particular promoter, transcription from those endogenous promoters is modulated.

(1) Cloning Methods

Isolation from genomic libraries of polynucleotides comprising the sequences of the genes, promoters and promoter control elements described in SEQ ID NO:1 and SEQ ID NO:2 or other polynucleotides of the present invention is possible using known techniques.

For example, polymerase chain reaction (PCR) can amplify the desired polynucleotides utilizing primers designed from sequences in SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6 or SEQ ID NO:8. Polynucleotide libraries comprising genomic sequences can be constructed according to Sambrook *et al.*, MOLECULAR CLONING: A LABORATORY MANUAL, 2nd Ed. (1989), for example.

Other procedures for isolating polynucleotides comprising the polynucleotide sequences of the invention include, without limitation, tail-PCR, and 5' rapid amplification of cDNA ends (RACE). For tail-PCR, *see, e.g.*, Liu *et al.*, *Plant J* 8(3): 457-463 (1995); Liu *et al.*, *Genomics* 25: 674-681 (1995); Liu *et al.*, *Nucl. Acids Res.* 21(14): 3333-3334 (1993); and Zoe *et al.*, *BioTechniques* 27(2): 240-248 (1999); for RACE, *see, e.g.*, PCR Protocols: A Guide to Methods and Applications, (1990) Academic Press, Inc.

(2) Chemical Synthesis

In addition, the genes, promoters and promoter control elements of the invention can be chemically synthesized according to techniques in common use. *See, e.g.*, Beaucage *et al.*, *Tet. Lett.* 22: 1859 (1981) and U.S. Pat. No. 4,668,777.

Such chemical oligonucleotide synthesis can be carried out using commercially available devices, such as, Biosearch 4600 or 8600 DNA synthesizer, by Applied Biosystems, a division of Perkin-Elmer Corp., Foster City, California, USA; and Expedite by Perceptive Biosystems, Framingham, Massachusetts, USA.

Synthetic RNA, including natural and/or analog building blocks, can be synthesized on the Biosearch 8600 machines, see above.

Oligonucleotides can be synthesized and then ligated together to construct the desired polynucleotide.

C. Isolating Related Polynucleotide Sequences

Included in the present invention are genes, promoters and promoter control elements which are related to those described in SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6 or SEQ ID NO:8. Such related sequence can be isolated utilizing nucleotide sequence identity; coding sequence identity; or common function or gene products.

Relatives can include both naturally occurring genes and promoters and non-natural gene and promoter sequences. Non-natural related gene or promoters include

nucleotide substitutions, insertions or deletions of naturally-occurring gene or promoter sequences that do not substantially affect activity of the polynucleotides (e.g., activity of coding sequences or transcription modulation). For example, the binding of relevant DNA binding proteins can still occur with the non-natural promoter sequences and promoter control elements of the present invention.

According to current knowledge, promoter sequences and promoter control elements exist as functionally important regions, such as protein binding sites, and spacer regions. These spacer regions are apparently required for proper positioning of the protein binding sites. Thus, nucleotide substitutions, insertions and deletions can be tolerated in these spacer regions to a certain degree without loss of function.

In contrast, less variation is permissible in the functionally important regions, since changes in the sequence can interfere with protein binding. Nonetheless, some variation in the functionally important regions is permissible so long as function is conserved. In some embodiments, functionally important regions can include nucleotides 3324 to 3580 of SEQ ID NO:1. As described below, nucleotides 3324 to 3580 of SEQ ID NO:2 are useful for modulating transcriptional activity in suspensor cells and/or basal regions of plant embryos.

The effects of substitutions, insertions and deletions to the promoter sequences or promoter control elements may be to increase or decrease the binding of relevant DNA binding proteins to modulate transcript levels of a polynucleotide to be transcribed. Effects may include tissue-specific or condition-specific modulation of transcript levels of the polypeptide to be transcribed. Polynucleotides representing changes to the nucleotide sequence of the DNA-protein contact region by insertion of additional nucleotides, changes to identity of relevant nucleotides, including use of chemically-modified bases, or deletion of one or more nucleotides are considered encompassed by the present invention.

(1) Relatives Based on Nucleotide Sequence Identity

Included in the present invention are polynucleotides comprising genes or promoters exhibiting nucleotide sequence identity to SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6 or SEQ ID NO:8.

Definition

Typically, such related genes or promoters exhibit at least 50%, sometimes at least 60% or at least 70% or at least 80% sequence identity, preferably at least 85%, more

preferably at least 90%, and most preferably at least 95%, even more preferably, at least 96%, 97%, 98% or 99% sequence identity compared to SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6 or SEQ ID NO:8. Indeed, any percent identity represented by an integer between 50-99 is contemplated for the invention. Such sequence identity can be calculated by the algorithms and computers programs described above.

Usually, such sequence identity is exhibited in an alignment region that is at least 75%, usually at least 80%; more usually, at least 85%, more usually at least 90%, and most usually at least 95%, even more usually, at least 96%, 97%, 98% or 99% of the length of a sequence shown in SEQ ID NO:1.

The percentage of the alignment length is calculated by counting the number of residues of the sequence in region of strongest alignment, e.g., a continuous region of the sequence that contains the greatest number of residues that are identical to the residues between two sequences that are being aligned. The number of residues in the region of strongest alignment is divided by the total residue length of a sequence in SEQ ID NO:1.

These related promoters may exhibit similar preferential transcription as SEQ ID NO:1 or other sequences of the invention such as nucleotides 1-4582 of SEQ ID NO:4, nucleotides 1-3154 of SEQ ID NO:6 or nucleotides 1-1609 of SEQ ID NO:8.

Construction of Polynucleotides

Naturally occurring promoters that exhibit nucleotide sequence identity to those shown in SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6 or SEQ ID NO:8 can be isolated using the techniques as described above. More specifically, such related promoters can be identified by varying stringencies, as defined above, in typical hybridization procedures such as, Southern or probing of polynucleotide libraries, for example.

Non-natural promoter variants of those shown in SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6 or SEQ ID NO:8 can be constructed using cloning methods that incorporate the desired nucleotide variation. See, for example, Ho, S. N., *et al.* *Gene* 77:51-59 (1989), describing a procedure site directed mutagenesis using PCR.

Any related promoter showing sequence identity to those shown in SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6 or SEQ ID NO:8 can be chemically synthesized as described above.

Also, the present invention includes non-natural promoters that exhibit the above-sequence identity to those in SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6 or SEQ ID NO:8.

The promoters and promoter control elements of the present invention may also be synthesized with 5' or 3' extensions, to facilitate additional manipulation, for instance.

(2) Relatives Based on Coding Sequence Identity

In addition, the present invention includes promoters of genes that comprise exons that encode polypeptide sequences that show sequence identity to the amino acid sequence displayed in SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, or SEQ ID NO:9.

Definition

Typically, the amino acid sequence of the genes comprising these related polynucleotides exhibit at least that exhibit at least 50%, at least 60%, at least 70% or at least 80% sequence identity to SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, or SEQ ID NO:9, preferably at least 85%, more preferably at least 90%, and most preferably at least 95%, even more preferably, at least 96%, 97%, 98% or 99% sequence identity to SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, or SEQ ID NO:9. Such sequence identity can be calculated by the algorithms and computers programs described above.

Usually, such sequence identity is exhibited in an alignment region that is at least 75% of the length of a sequence encoded by SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6 or SEQ ID NO:8 or corresponding full-length sequence; more usually at least 80%; more usually, at least 85%, more usually at least 90%, and most usually at least 95%, even more usually, at least 96%, 97%, 98% or 99% of the length of a sequence encoded by SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6 or SEQ ID NO:8.

Construction of Polynucleotides

The isolation of sequences from the genes of the invention may be accomplished by a number of techniques. For instance, oligonucleotide probes based on the sequences disclosed here can be used to identify the desired gene in a cDNA or genomic DNA library from a desired plant species. To construct genomic libraries, large segments of genomic DNA are generated by random fragmentation, e.g. using restriction endonucleases, and are ligated with vector DNA to form concatemers that can be packaged into the appropriate vector. To prepare a library of embryo-specific cDNAs, mRNA is isolated from embryos and a cDNA library that contains the gene transcripts is prepared from the mRNA.

The cDNA or genomic library can then be screened using a probe based upon the sequence of a cloned embryo-specific gene such as the polynucleotides disclosed here.

Probes may be used to hybridize with genomic DNA or cDNA sequences to isolate homologous genes in the same or different plant species.

Alternatively, the nucleic acids of interest can be amplified from nucleic acid samples using amplification techniques. For instance, polymerase chain reaction (PCR) technology to amplify the sequences of the genes directly from mRNA, from cDNA, from genomic libraries or cDNA libraries. PCR and other in vitro amplification methods may also be useful, for example, to clone nucleic acid sequences that code for proteins to be expressed, to make nucleic acids to use as probes for detecting the presence of the desired mRNA in samples, for nucleic acid sequencing, or for other purposes. Appropriate primers and probes for identifying embryo-specific genes from plant tissues are generated from comparisons of the sequences provided herein. For a general overview of PCR see PCR Protocols: A Guide to Methods and Applications. (Innis, M, Gelfand, D., Sninsky, J. and White, T., eds.), Academic Press, San Diego (1990).

Polynucleotides may also be synthesized by well-known techniques as described in the technical literature. See, e.g., Carruthers *et al.*, Cold Spring Harbor Symp. *Quant. Biol.* 47:411-418 (1982), and Adams *et al.*, *J. Am. Chem. Soc.* 105:661 (1983). Double stranded DNA fragments may then be obtained either by synthesizing the complementary strand and annealing the strands together under appropriate conditions, or by adding the complementary strand using DNA polymerase with an appropriate primer sequence.

Identified cDNA sequences can be aligned to the genomic sequences to identify the promoter region and sequences, which are located upstream of the 5'UTR and downstream of the preceding gene.

cDNA Isolation

The cDNAs can be isolated by various cloning methods described above. For example, probes and/or primer can be designed utilizing the sequences in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6 or SEQ ID NO:8. See, e.g., Ausubel *et al.* (1992); and Sambrook *et al.* (1989).

Such probes and primers can be used to identify cDNAs with a comprising at least one transcription start site. Full-length cDNA libraries are useful to identify cDNAs with at least one transcription start site. Such libraries can be constructed as described in the above-captioned applications in the Related Applications Section. Alternatively, tail-PCR or RACE can be used to isolated the 5' end of a cDNA.

Genomic Polynucleotide Isolation

Genomic sequences can be isolated with the sequence from the cDNA also found in the 5' UTR, exons or 3' UTR for probes and/or primers.

Alternatively, the promoter sequences upstream of the transcription start site or translation start site can be isolated using single primers designed having the portions of cDNA sequences 3' of the start codon of a sequence (e.g., SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6 or SEQ ID NO:8) and used with random primers to isolate the corresponding upstream portion of genomic DNA.

Alternatively the promoters and promoter control elements of the invention can be identified by "walking" upstream from 5'-most portions of cDNA sequences in a genomic DNA library.

The promoter sequences will those 5' of the transcription start site which can be located using the 5' end of the corresponding cDNA. Alternatively, the start sites of a transcript can be assessed using primer extension assays (King *et al.*, *Gene* 242:125 (2000)).

In addition, the 5' end of the promoter can be identified by either locating the upstream polyA signal or by identifying the cDNA corresponding to the preceding gene using the techniques described above.

D. Identifying Control Elements

(1) Types of Transcription Control Elements

Promoter sequences comprise a number of promoter control elements that are capable of initiating transcription, regulating transcription rates and levels, etc. Promoter control elements modulate transcription when such control elements exhibit their transcription related activities, such as hybridizing to target polynucleotides; binding to repressor proteins, transcription factors, proteins or components of the nuclear matrix; able to act as a methylation site, etc. Promoter control elements include cis acting elements such as enhancers, scaffold/matrix attachment regions (S/MARs), locus control regions (LCRs).

Other promoter control elements include, without limitation:
core or basal promoters,
TATA boxes,
initiator sites,

transcription factor binding sites,
repressor binding sites;
and inverted repeats.

See, e.g., T. Bouliskas, J. Cell Biochem., 60, 297-316 (1996).

Promoter Control Elements of the Invention

The promoter control elements of the present invention include those that comprise SEQ ID NO:1, nucleotides 1-4582 of SEQ ID NO:4, nucleotides 1-3154 of SEQ ID NO:6 or nucleotides 1-1609 of SEQ ID NO:8, and fragments thereof. A particularly preferred fragment comprises nucleotides 3329 to 3475 of SEQ ID NO:1. As discussed below, this fragment confers suspensor-specific activity to a promoter. Additional promoter control elements include SEQ ID NO:10 and SEQ ID NO:11. Control elements of the invention alone, or as part of a heterologous promoter, are useful for modulation of transcription.

The size of the fragments of SEQ ID NO:1, nucleotides 1-4582 of SEQ ID NO:4, nucleotides 1-3154 of SEQ ID NO:6 or nucleotides 1-1609 of SEQ ID NO:8 can range from 5 bases to about 5 kilobases (kb). Typically, the fragment size is no smaller than 8 bases; more typically, no smaller than 10 or 12; more typically, no smaller than 15 bases; more typically, no smaller than 20 bases; more typically, no smaller than 25 bases; even more typically, no more than 30, 35, 40 or 50 bases.

Usually, the fragment size is no larger than 2 kb bases; more usually, no larger than 1 kb; more usually, no larger than 800 bases; more usually, no larger than 500 bases; even more usually, no more than 250, 200, 150 or 100 bases.

Relatives Based on Nucleotide Sequence Identity

Included in the present invention are promoter control elements exhibiting nucleotide sequence identity to those in SEQ ID NO:1, nucleotides 1-4582 of SEQ ID NO:4, nucleotides 1-3154 of SEQ ID NO:6 or nucleotides 1-1609 of SEQ ID NO:8.

Typically, such related promoters exhibit at least 80% sequence identity, preferably at least 85%, more preferably at least 90%, and most preferably at least 95%, even more preferably, at least 96%, 97%, 98% or 99% sequence identity compared to those shown in SEQ ID NO:1, nucleotides 1-4582 of SEQ ID NO:4, nucleotides 1-3154 of SEQ ID NO:6 or nucleotides 1-1609 of SEQ ID NO:8. Such sequence identity can be calculated by the algorithms and computer programs described above.

Relatives Based on Coding Sequence Identity

In addition, the present invention includes promoter control elements of genes that comprise exons that encode polypeptide sequences that show sequence identity to SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7 or SEQ ID NO:9.

Typically, the amino acid sequence of the genes comprising these related promoters exhibit at least 80% sequence identity to those shown in SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7 or SEQ ID NO:9, preferably at least 85%, more preferably at least 90%, and most preferably at least 95%, even more preferably, at least 96%, 97%, 98% or 99% sequence identity to SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7 or SEQ ID NO:9. Such sequence identity can be calculated by the algorithms and computers programs described above.

Usually, such sequence identity is exhibited in an alignment region that is at least 75% of the length of SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7 or SEQ ID NO:9; more usually at least 80%; more usually, at least 85%, more usually at least 90%, and most usually at least 95%, even more usually, at least 96%, 97%, 98% or 99% of the length of SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7 or SEQ ID NO:9.

Promoter Control Element Configuration

A common configuration of the promoter control elements in RNA polymerase II promoters is shown in FIGURE 1.

For more description, *see, e.g.*, T. Werner, *Mammalian Genome*, 10, 168-175 (1999).

Promoters are generally modular in nature. Promoters can consist of a basal promoter that functions as a site for assembly of a transcription complex comprising an RNA polymerase, for example RNA polymerase II. A typical transcription complex will include additional factors such as TF_{II}B, TF_{II}D, and TF_{II}E. Of these, TF_{II}D appears to be the only one to bind DNA directly. The promoter might also contain one or more promoter control elements such as the elements discussed above. These additional control elements may function as binding sites for additional transcription factors that have the function of modulating the level of transcription with respect to tissue specificity and of transcriptional responses to particular environmental or nutritional factors, and the like.

One type of promoter control elements are polynucleotide sequences representing binding sites for proteins. Typically, within a particular functional module,

protein binding sites constitute regions of 5 to 60, preferably 10 to 30, more preferably 10 to 20 nucleotides. Within such binding sites, there are typically 2 to 6 nucleotides that specifically contact amino acids of the nucleic acid binding protein.

The protein binding sites are usually separated from each other by 10 to several hundred nucleotides, typically by 15 to 150 nucleotides, often by 20 to 50 nucleotides.

Further, protein binding sites in promoter control elements often display dyad symmetry in their sequence. Such elements can bind several different proteins, and/or a plurality of sites can bind the same protein. Both types of elements may be combined in a region of 50 to 1,000 base pairs.

Binding sites for any specific factor have been known to occur almost anywhere in a promoter. For example, functional AP-1 binding sites can be located far upstream, as in the rat bone sialoprotein gene, where an AP-1 site located about 900 nucleotides upstream of the transcription start site suppresses expression. Yamauchi *et al.*, *Matrix Biol.*, 15, 119-130 (1996). Alternatively, an AP-1 site located close to the transcription start site plays an important role in the expression of Moloney murine leukemia virus. Sap *et al.*, *Nature*, 340, 242-244 (1989).

(2) Those Identifiable by Bioinformatics

Promoter control elements from the promoters of the instant invention can be identified utilizing bioinformatic or computer driven techniques.

One method uses a computer program AlignACE to identify regulatory motifs in genes that exhibit common preferential transcription across a number of time points. The program identifies common sequence motifs in such genes. See, Roth *et al.*, *Nature Biotechnol.* 16: 949-945 (1998); Tavazoie *et al.*, *Nat Genet* 22(3):281-5 (1999).

Genomatix, also makes available a GEMS Launcher program and other programs to identify promoter control elements and configuration of such elements. Genomatix is located in Munich, Germany.

Other references also describe detection of promoter modules by models independent of overall nucleotide sequence similarity. See, e.g., Klingenhoff *et al.*, *Bioinformatics* 15, 180-186 (1999).

Protein binding sites of promoters can be identified as reported in Frech, *et al.*, *Nucleic Acids Research*, Vol. 21, No. 7, 1655-1664 (1993).

Other programs used to identify protein binding sites include, for example, Signal Scan, Prestidge *et al.*, *Comput. Appl. Biosci.* 12: 157-160 (1996); Matrix Search, Chen *et al.*, *Comput. Appl. Biosci.* 11: 563-566 (1995), available as part of Signal Scan 4.0; MatInspector, Ghosh *et al.*, *Nucl. Acid Res.* 21: 3117-3118 (1993) available
 5 <http://www.gsf.de/cgi-bin/matsearch.pl>; ConsInspector, Frech *et al.*, *Nucl. Acids Res.* 21: 1655-1664 (1993), available at <ftp://ariane.gsf.de/pub/dos>; TFSearch; and TESS.

Frech *et al.*, "Software for the analysis of DNA sequence elements of transcription" in *BIOINFORMATICS & SEQUENCE ANALYSIS*, Vol. 13, no. 1, 89-97 (1997) is a review of different software for analysis of promoter control elements. This paper also
 10 reports the usefulness of matrix-based approaches to yield more specific results.

For other procedures, *see*, Fickett *et al.*, *Curr. Op. Biotechnol.* 11: 19-24 (2000); and Quandt *et al.*, *Nucleic Acids Res.* 23, 4878-4884 (1995).

(3) Those Identifiable by In-Vitro and In-Vivo Assays

Promoter control elements also can be identified with in-vitro assays, such as transcription detection methods; and with in-vivo assays, such as enhancer trapping protocols.

In-Vitro Assays

Examples of *in vitro* assays include detection of binding of protein factors that bind promoter control elements. Fragments of the instant promoters can be used to identify the location of promoter control elements. Another option for obtaining a promoter control element with desired properties is to modify known promoter sequences. This is based on the fact that the function of a promoter is dependent on the interplay of regulatory proteins that
 25 bind to specific, discrete nucleotide sequences in the promoter, termed motifs. Such interplay subsequently affects the general transcription machinery and regulates transcription efficiency. These proteins are positive regulators or negative regulators (repressors), and one protein can have a dual role depending on the context (Johnson, P. F. and McKnight, S. L. *Annu. Rev. Biochem.* 58:799-839 (1989)).

One type of in-vitro assay utilizes a known DNA binding factor to isolate DNA fragments that bind. If a fragment or promoter variant does not bind, then a promoter control element has been removed or disrupted. For specific assays, *see, e.g.*, B. Luo *et al.*, *J. Mol. Biol.* 266:470 (1997), S. Chusacultachai *et al.*, *J. Biol. Chem.* 274:23591 (1999), D. Fabbro *et al.*, *Biochem. Biophys. Res. Comm.* 213:781 (1995)).

Alternatively, a fragment of DNA suspected of conferring a particular pattern of specificity can be examined for activity in binding transcription factors involved in that specificity by methods such as DNA footprinting (e.g. D.J. Cousins *et al.*, *Immunology* 99:101 (2000); V. Kolla *et al.*, *Biochem. Biophys. Res. Comm.* 266:5 (1999)) or “mobility-shift” assays (E.D. Fabiani *et al.*, *J. Biochem.* 347:147 (2000); N. Sugiura *et al.*, *J. Biochem* 347:155 (2000)) or fluorescence polarization (e.g. Royer *et al.*, U.S. Patent 5,445,935). Both mobility shift and DNA footprinting assays can also be used to identify portions of large DNA fragments that are bound by proteins in unpurified transcription extracts prepared from tissues or organs of interest.

Cell-free transcription extracts can be prepared and used to directly assay in a reconstitutable system (Narayan *et al.*, *Biochemistry* 39:818 (2000)).

In-Vivo Assays

Promoter control elements can be identified with reporter genes in in-vivo assays with the use of fragments of the instant promoters or variants of the instant promoter polynucleotides.

For example, various fragments can be inserted into a vector, comprising a basal promoter, for example, operably linked to a reporter sequence, which, when transcribed, can produce a detectable label. Examples of reporter genes include those encoding luciferase, green fluorescent protein, GUS, neo, cat and bar. Alternatively, reporter sequence can be detected utilizing AFLP and microarray techniques.

In promoter probe vector systems, genomic DNA fragments are inserted upstream of the coding sequence of a reporter gene that is expressed only when the cloned fragment contains DNA having transcription modulation activity (Neve, R. L. *et al.*, *Nature* 277:324-325 (1979)). Control elements are disrupted when fragments or variants lacking any transcription modulation activity. Probe vectors have been designed for assaying transcription modulation in *E. coli* (An, G. *et al.*, *J. Bact.* 140:400-407 (1979)) and other bacterial hosts (Band, L. *et al.*, *Gene* 26:313-315 (1983); Achen, M. G., *Gene* 45:45-49 (1986)), yeast (Goodey, A. R. *et al.*, *Mol. Gen. Genet.* 204:505-511 (1986)) and mammalian cells (Pater, M. M. *et al.*, *J. Mol. App. Gen.* 2:363-371 (1984)).

A different design of a promoter/control element trap includes packaging into retroviruses for more efficient delivery into cells. One type of retroviral enhancer trap was described by von Melchner *et al.* (*Genes Dev.* 6(6):919-27 (1992); U.S. Pat. No. 5,364,783). The basic design of this vector includes a reporter protein coding sequence engineered into

the U3 portion of the 3' LTR. No splice acceptor consensus sequences are included, limiting its utility to work as an enhancer trap only. A different approach to a gene trap using retroviral vectors was pursued by Friedrich and Soriano (*Genes Dev.* 5(9):1513-23 (1991)), who engineered a lacZ-neo fusion protein linked to a splicing acceptor. LacZ-neo fusion protein expression from trapped loci allows not only for drug selection, but also for visualization of β -galactosidase expression using the chromogenic substrate, X-gal.

A general review of tools for identifying transcriptional regulatory regions of genomic DNA is provided by J.W. Fickett *et al. Curr. Opin. Biotechnol.* 11:19 (2000).

(4) Non-Natural Control Elements

Non-natural control elements can be constructed by inserting, deleting or substituting nucleotides into the promoter control elements described above. Such control elements are capable of transcription modulation which can be determined using any of the assays described above.

E. Constructing Promoters with Control Elements

(1) Combining Promoters and Promoter Control Elements

The promoter polynucleotides and promoter control elements of the present invention, both naturally occurring and synthetic, can be combined with each other to produce the desired preferential transcription. Also, the polynucleotides of the invention can be combined with other known sequences to obtain other useful promoters to modulate, for example, tissue transcription specific or transcription specific to certain conditions. Such preferential transcription can be determined using the techniques or assays described above.

Fragments, variants, as well as full-length sequences such as those shown in SEQ ID NO:1, nucleotides 1-4582 of SEQ ID NO:4, nucleotides 1-3154 of SEQ ID NO:6 or nucleotides 1-1609 of SEQ ID NO:8 and relatives are useful alone or in combination.

The location and relation of promoter control elements within a promoter can affect the ability of the promoter to modulate transcription. The order and spacing of control elements is a factor when constructing promoters.

(2) Number of Promoter Control Elements

Promoters can contain any number of control elements. For example, a promoter can contain multiple transcription binding sites or other control elements. One element may confer tissue or organ specificity; another element may limit transcription to

specific time periods, etc. Typically, promoters will contain at least a basal or core promoter as described above. Any additional element can be included as desired. For example, a fragment comprising a basal promoter can be fused with another fragment with any number of additional control elements.

(3) Spacing Between Control Elements

Spacing between control elements or the configuration or control elements can be determined or optimized to permit the desired protein-polynucleotide or polynucleotide interactions to occur.

For example, if two transcription factors bind to a promoter simultaneously or relatively close in time, the binding sites are spaced to allow each factor to bind without steric hindrance. The spacing between two such hybridizing control elements can be as small as a profile of a protein bound to a control element. In some cases, two protein binding sites can be adjacent to each other when the proteins bind at different times during the transcription process.

Further, when two control elements hybridize the spacing between such elements will be sufficient to allow the promoter polynucleotide to hairpin or loop to permit the two elements to bind. The spacing between two such hybridizing control elements can be as small as a t-RNA loop, to as large as 10 kb.

Typically, the spacing is no smaller than 5 bases; more typically, no smaller than 8; more typically, no smaller than 15 bases; more typically, no smaller than 20 bases; more typically, no smaller than 25 bases; even more typically, no more than 30, 35, 40 or 50 bases.

Usually, the fragment size is no larger than 5 kb bases; more usually, no larger than 2 kb; more usually, no larger than 1 kb; more usually, no larger than 800 bases; more usually, no larger than 500 bases; even more usually, no more than 250, 200, 150 or 100 bases.

Such spacing between promoter control elements can be determined using the techniques and assays described above.

F. Control of G564 or C541 Activity of Gene Expression

(1) Use Of Nucleic Acids of the Invention to Inhibit Gene Expression

The isolated sequences prepared as described herein, can be used to prepare expression cassettes useful in a number of techniques. For example, expression cassettes of the invention can be used to suppress endogenous G564 or C541 gene expression. Inhibiting expression can be useful, for instance, to modulate or prevent suspensor cell development

and/or embryo size, shape and/or rate of development. Inhibition of expression is also useful for modulating fertility of a plant.

A number of methods can be used to inhibit gene expression in plants. For instance, antisense technology can be conveniently used. To accomplish this, a nucleic acid segment from the desired gene is cloned and operably linked to a promoter such that the antisense strand of RNA will be transcribed. The expression cassette is then transformed into plants and the antisense strand of RNA is produced. In plant cells, it has been suggested that antisense RNA inhibits gene expression by preventing the accumulation of mRNA which encodes the enzyme of interest, see, e.g., Sheehy *et al.*, *Proc. Nat. Acad. Sci. USA*, 85:8805-8809 (1988), and Hiatt *et al.*, U.S. Patent No. 4,801,340.

The antisense nucleic acid sequence transformed into plants will be substantially identical to at least a portion of the endogenous suspensor-specific gene or genes to be repressed. The sequence, however, does not have to be perfectly identical to inhibit expression. The vectors of the present invention can be designed such that the inhibitory effect applies to other proteins within a family of genes exhibiting homology or substantial homology to the target gene.

For antisense suppression, the introduced sequence also need not be full length relative to either the primary transcription product or fully processed mRNA. Generally, higher homology can be used to compensate for the use of a shorter sequence. Furthermore, the introduced sequence need not have the same intron or exon pattern, and homology of non-coding segments may be equally effective. Normally, a sequence of between about 30 or 40 nucleotides and about full length nucleotides should be used, though a sequence of at least about 100 nucleotides is preferred, a sequence of at least about 200 nucleotides is more preferred, and a sequence of at least about 500 nucleotides is especially preferred.

Catalytic RNA molecules or ribozymes can also be used to inhibit expression of embryo-specific genes. It is possible to design ribozymes that specifically pair with virtually any target RNA and cleave the phosphodiester backbone at a specific location, thereby functionally inactivating the target RNA. In carrying out this cleavage, the ribozyme is not itself altered, and is thus capable of recycling and cleaving other molecules, making it a true enzyme. The inclusion of ribozyme sequences within antisense RNAs confers RNA-cleaving activity upon them, thereby increasing the activity of the constructs.

A number of classes of ribozymes have been identified. One class of ribozymes is derived from a number of small circular RNAs that are capable of self-cleavage and replication in plants. The RNAs replicate either alone (viroid RNAs) or with a helper

virus (satellite RNAs). Examples include RNAs from avocado sunblotch viroid and the satellite RNAs from tobacco ringspot virus, lucerne transient streak virus, velvet tobacco mottle virus, solanum nodiflorum mottle virus and subterranean clover mottle virus. The design and use of target RNA-specific ribozymes is described in Haseloff et al. *Nature*, 334:585-591 (1988).

Another method of suppression is sense suppression. Introduction of expression cassettes in which a nucleic acid is configured in the sense orientation with respect to the promoter has been shown to be an effective means by which to block the transcription of target genes. For an example of the use of this method to modulate expression of endogenous genes see, Napoli et al., *The Plant Cell* 2:279-289 (1990), and U.S. Patents Nos. 5,034,323, 5,231,020, and 5,283,184.

Generally, where inhibition of expression is desired, some transcription of the introduced sequence occurs. The effect may occur where the introduced sequence contains no coding sequence per se, but only intron or untranslated sequences homologous to sequences present in the primary transcript of the endogenous sequence. The introduced sequence generally will be substantially identical to the endogenous sequence intended to be repressed. This minimal identity will typically be greater than about 65%, but a higher identity might exert a more effective repression of expression of the endogenous sequences. Substantially greater identity of more than about 80% is preferred, though about 95% to absolute identity would be most preferred. As with antisense regulation, the effect should apply to any other proteins within a similar family of genes exhibiting homology or substantial homology.

For sense suppression, the introduced sequence in the expression cassette, needing less than absolute identity, also need not be full length, relative to either the primary transcription product or fully processed mRNA. This may be preferred to avoid concurrent production of some plants that are overexpressers. A higher identity in a shorter than full-length sequence compensates for a longer, less identical sequence. Furthermore, the introduced sequence need not have the same intron or exon pattern, and identity of non-coding segments will be equally effective. Normally, a sequence of the size ranges noted above for antisense regulation is used.

One of skill in the art will recognize that using technology based on specific nucleotide sequences (e.g., antisense or sense suppression technology), families of homologous genes can be suppressed with a single sense or antisense transcript. For instance, if a sense or antisense transcript is designed to have a sequence that is conserved

among a family of genes, then multiple members of a gene family can be suppressed. Conversely, if the goal is to only suppress one member of a homologous gene family, then the sense or antisense transcript should be targeted to sequences with the most variance between family members.

Another means of inhibiting G564 or C541 function in a plant is by creation of dominant negative mutations. In this approach, non-functional, mutant G564 or C541 polypeptides, which retain the ability to interact with wild-type subunits are introduced into a plant.

(2) Use of Nucleic Acids of the Invention to Enhance Gene Expression

Isolated sequences prepared as described herein can also be used to prepare expression cassettes that enhance or increase endogenous G564 or C5541 gene expression. Where overexpression of a gene is desired, the desired gene from a different species may be used to decrease potential sense suppression effects. Enhanced expression of G564 or C541 polynucleotides is useful, for example, to modulate suspensor cell and/or embryo size, shape and/or rate of development. Enhanced expression is also useful for modulating plant fertility.

Any of a number of means well known in the art can be used to increase G564 or C541 activity in plants. Any organ can be targeted, such as shoot vegetative organs/structures (*e.g.* leaves, stems and tubers), roots, flowers and floral organs/structures (*e.g.* bracts, sepals, petals, stamens, carpels, anthers and ovules), seed (including apical or basal cells, suspensor, embryo, endosperm, and seed coat) and fruit. Alternatively, one or several G564 or C541 genes can be expressed constitutively (*e.g.*, using the CaMV 35S promoter).

One of skill will recognize that the polypeptides encoded by the genes of the invention, like other proteins, have different domains that perform different functions. Thus, the gene sequences need not be full length, so long as the desired functional domain of the protein is expressed.

(3) Modification of endogenous G564 or C541 genes

Methods for introducing genetic mutations into plant genes and selecting plants with desired traits are well known. For instance, seeds or other plant material can be treated with a mutagenic chemical substance, according to standard techniques. Such chemical substances include, but are not limited to, the following: diethyl sulfate, ethylene

imine, ethyl methanesulfonate and N-nitroso-N-ethylurea. Alternatively, ionizing radiation from sources such as, X-rays or gamma rays can be used.

Modified protein chains can also be readily designed utilizing various recombinant DNA techniques well known to those skilled in the art and described for instance, in Sambrook et al., supra. Hydroxylamine can also be used to introduce single base mutations into the coding region of the gene (Sikorski, *et al.*, (1991). *Meth. Enzymol.* 194: 302-318). For example, the chains can vary from the naturally occurring sequence at the primary structure level by amino acid substitutions, additions, deletions, and the like. These modifications can be used in a number of combinations to produce the final modified protein chain.

Alternatively, homologous recombination can be used to induce targeted gene modifications by specifically targeting the *G564* or *C541* gene *in vivo* (see, generally, Grewal and Klar, *Genetics* 146: 1221-1238 (1997) and Xu *et al.*, *Genes Dev.* 10: 2411-2422 (1996)). Homologous recombination has been demonstrated in plants (Puchta *et al.*, *Experientia* 50: 277-284 (1994), Swoboda *et al.*, *EMBO J.* 13: 484-489 (1994); Offringa *et al.*, *Proc. Natl. Acad. Sci. USA* 90: 7346-7350 (1993); and Kempin *et al.* *Nature* 389:802-803 (1997)).

In applying homologous recombination technology to the genes of the invention, mutations in selected portions of an *G564* or *C541* gene sequences (including 5' upstream, 3' downstream, and intragenic regions) such as those disclosed here are made *in vitro* and then introduced into the desired plant using standard techniques. Since the efficiency of homologous recombination is known to be dependent on the vectors used, use of dicistronic gene targeting vectors as described by Mountford *et al.*, *Proc. Natl. Acad. Sci. USA* 91: 4303-4307 (1994); and Vaulont *et al.*, *Transgenic Res.* 4: 247-255 (1995) are conveniently used to increase the efficiency of selecting for altered *G564* or *C541* gene expression in transgenic plants. The mutated gene will interact with the target wild-type gene in such a way that homologous recombination and targeted replacement of the wild-type gene will occur in transgenic plant cells, resulting in suppression of *G564* or *C541* activity.

Alternatively, oligonucleotides composed of a contiguous stretch of RNA and DNA residues in a duplex conformation with double hairpin caps on the ends can be used. The RNA/DNA sequence is designed to align with the sequence of the target *G564* or *C541* gene and to contain the desired nucleotide change. Introduction of the chimeric oligonucleotide on an extrachromosomal T-DNA plasmid results in efficient and specific *G564* or *C541* gene conversion directed by chimeric molecules in a small number of

transformed plant cells. This method is described in Cole-Strauss *et al.*, *Science* 273:1386-1389 (1996) and Yoon *et al.* *Proc. Natl. Acad. Sci. USA* 93: 2071-2076 (1996).

G. Heterologous Expression of the G564 or C541 Polynucleotides of the Invention

A DNA sequence coding for the desired polypeptide, for example a cDNA sequence encoding a full length protein, will preferably be combined with transcriptional and translational initiation regulatory sequences which will direct the transcription of the sequence from the gene in the intended tissues of the transformed plant.

For example, for overexpression, a plant promoter fragment may be employed which will direct expression of the gene in all tissues of a regenerated plant. Such promoters are referred to herein as "constitutive" promoters and are active under most environmental conditions and states of development or cell differentiation. Examples of constitutive promoters include the cauliflower mosaic virus (CaMV) 35S transcription initiation region, the 1'- or 2'- promoter derived from T-DNA of *Agrobacterium tumefaciens*, and other transcription initiation regions from various plant genes known to those of skill.

Alternatively, the plant promoter may direct expression of the polynucleotide of the invention in a specific tissue (tissue-specific promoters) or may be otherwise under more precise environmental control (inducible promoters). Examples of tissue-specific promoters under developmental control include promoters that initiate transcription only in certain tissues, such as fruit, seeds, or flowers. As noted above, the promoters from the *G564* or *C541* genes described here are particularly useful for directing gene expression so that a desired gene product is located in suspensor cells. Examples of environmental conditions that may affect transcription by inducible promoters include anaerobic conditions, elevated temperature, or the presence of light.

If proper polypeptide expression is desired, a polyadenylation region at the 3'-end of the coding region should be included. The polyadenylation region can be derived from the natural gene, from a variety of other plant genes, or from T-DNA.

The vector comprising the sequences (e.g., promoters or coding regions) from genes of the invention will typically comprise a marker gene which confers a selectable phenotype on plant cells. For example, the marker may encode biocide resistance, particularly antibiotic resistance, such as resistance to kanamycin, G418, bleomycin, hygromycin, or herbicide resistance, such as resistance to chlorosulfuron or Basta.

G564 or C541 nucleic acid sequences of the invention are expressed recombinantly in plant cells to enhance and increase levels of endogenous G564 or C541

polypeptides. Alternatively, antisense or other G564 or C541 constructs (described above) are used to suppress G564 or C541 levels of expression. A DNA sequence coding for a G564 or C541 polypeptide, e.g., a cDNA sequence encoding a full length protein, can be combined with cis-acting (promoter) and trans-acting (enhancer) transcriptional regulatory sequences to direct the timing, tissue type and levels of transcription in the intended tissues of the transformed plant. Translational control elements can also be used.

The invention provides a G564 or C541 nucleic acid operably linked to a promoter that, in a preferred embodiment, is capable of driving the transcription of the G564 or C541 coding sequence in plants. The promoter can be, e.g., derived from plant or viral sources. The promoter can be, e.g., constitutively active, inducible, or tissue specific. In construction of recombinant expression cassettes, vectors, transgenics, of the invention, a different promoters can be chosen and employed to differentially direct gene expression, e.g., in some or all tissues of a plant or animal.

Typically, desired promoters are identified by analyzing the 5' sequences of a genomic clone corresponding to the suspensor-specific genes described here. Sequences characteristic of promoter sequences can be used to identify the promoter. Sequences controlling eukaryotic gene expression have been extensively studied. For instance, promoter sequence elements include the TATA box consensus sequence (TATAAT), which is usually 20 to 30 base pairs upstream of the transcription start site. In most instances the TATA box is required for accurate transcription initiation. In plants, further upstream from the TATA box, at positions -80 to -100, there is typically a promoter element with a series of adenines surrounding the trinucleotide G (or T) N G. J. Messing *et al.*, in GENETIC ENGINEERING IN PLANTS, pp.221-227 (Kosage, Meredith and Hollaender, eds. (1983)). A number of methods are known to those of skill in the art for identifying and characterizing promoter regions in plant genomic DNA (see, e.g., Jordano, *et al.*, *Plant Cell*, 1: 855-866 (1989); Bustos, *et al.*, *Plant Cell*, 1:839-854 (1989); Green, *et al.*, *EMBO J.* 7, 4035-4044 (1988); Meier, *et al.*, *Plant Cell*, 3, 309-316 (1991); and Zhang (1996) *Plant Physiology* 110:1069-1079).

Constitutive Promoters

A promoter fragment can be employed which will direct expression of G564 or C541 nucleic acid in all transformed cells or tissues, e.g. as those of a regenerated plant. Such promoters are referred to herein as "constitutive" promoters and are active under most environmental conditions and states of development or cell differentiation. Promoters that drive expression continuously under physiological conditions are referred to as "constitutive"

promoters and are active under most environmental conditions and states of development or cell differentiation. Examples of constitutive promoters include those from viruses which infect plants, such as the cauliflower mosaic virus (CaMV) 35S transcription initiation region (see, e.g., Dagless (1997) *Arch. Virol.* 142:183-191); the 1'- or 2'- promoter derived from T-DNA of *Agrobacterium tumefaciens* (see, e.g., Mengiste (1997) *supra*; O'Grady (1995) *Plant Mol. Biol.* 29:99-108); the promoter of the tobacco mosaic virus; the promoter of Figwort mosaic virus (see, e.g., Maiti (1997) *Transgenic Res.* 6:143-156); actin promoters, such as the *Arabidopsis* actin gene promoter (see, e.g., Huang (1997) *Plant Mol. Biol.* 1997 33:125-139); alcohol dehydrogenase (Adh) gene promoters (see, e.g., Millar (1996) *Plant Mol. Biol.* 31:897-904); *ACT11* from *Arabidopsis* (Huang *et al.* *Plant Mol. Biol.* 33:125-139 (1996)), *Cat3* from *Arabidopsis* (GenBank No. U43147, Zhong *et al.*, *Mol. Gen. Genet.* 251:196-203 (1996)), the gene encoding stearoyl-acyl carrier protein desaturase from *Brassica napus* (Genbank No. X74782, Solocombe *et al.* *Plant Physiol.* 104:1167-1176 (1994)), *Gpc1* from maize (GenBank No. X15596, Martinez *et al.* *J. Mol. Biol.* 208:551-565 (1989)), *Gpc2* from maize (GenBank No. U45855, Manjunath *et al.*, *Plant Mol. Biol.* 33:97-112 (1997)), other transcription initiation regions from various plant genes known to those of skill. See also Holtorf (1995) "Comparison of different constitutive and inducible promoters for the overexpression of transgenes in *Arabidopsis thaliana*," *Plant Mol. Biol.* 29:637-646.

Inducible Promoters

Alternatively, a plant promoter may direct expression of the G564 or C541 nucleic acids of the invention under the influence of changing environmental conditions or developmental conditions. Examples of environmental conditions that may effect transcription by inducible promoters include anaerobic conditions, elevated temperature, drought, or the presence of light. Such promoters are referred to herein as "inducible" promoters. For example, the invention incorporates the drought-inducible promoter of maize (Busk (1997) *supra*); the cold, drought, and high salt inducible promoter from potato (Kirch (1997) *Plant Mol. Biol.* 33:897-909).

Alternatively, plant promoters which are inducible upon exposure to plant hormones, such as auxins, are used to express the nucleic acids of the invention. For example, the invention can use the auxin-response elements E1 promoter fragment (AuxREs) in the soybean (*Glycine max* L.) (Liu (1997) *Plant Physiol.* 115:397-407); the auxin-responsive *Arabidopsis* GST6 promoter (also responsive to salicylic acid and hydrogen peroxide) (Chen (1996) *Plant J.* 10: 955-966); the auxin-inducible *parC* promoter from

tobacco (Sakai (1996) 37:906-913); a plant biotin response element (Streit (1997) *Mol. Plant Microbe Interact.* 10:933-937); and, the promoter responsive to the stress hormone abscisic acid (Sheen (1996) *Science* 274:1900-1902).

Plant promoters which are inducible upon exposure to chemicals reagents which can be applied to the plant, such as herbicides or antibiotics, are also used to express the nucleic acids of the invention. For example, the maize In2-2 promoter, activated by benzenesulfonamide herbicide safeners, can be used (De Veylder (1997) *Plant Cell Physiol.* 38:568-577); application of different herbicide safeners induces distinct gene expression patterns, including expression in the root, hydathodes, and the shoot apical meristem. The G564 or C541 coding sequences can also be under the control of, e.g., a tetracycline-inducible promoter, e.g., as described with transgenic tobacco plants containing the *Avena sativa* L. (oat) arginine decarboxylase gene (Masgrau (1997) *Plant J.* 11:465-473); or, a salicylic acid-responsive element (Stange (1997) *Plant J.* 11:1315-1324).

The following are promoters that are induced under stress conditions and can be combined with those of the present invention: ldh1 (oxygen stress; tomato; see Germain and Ricard *Plant Mol Biol* 35:949-54 (1997)), GPx and CAT (oxygen stress; mouse; see Franco *et al. Free Radic Biol Med* 27:1122-32 (1999)), ci7 (cold stress; potato; see Kirch *et al. Plant Mol Biol.* 33:897-909 (1997)), Bz2 (heavy metals; maize; see Marrs and Walbot. *Plant Physiol* 113:93-102 (1997)), HSP32 (hyperthermia; rat; see Raju and Maines. *Biochim Biophys Acta* 1217:273-80 (1994)); MAPKAPK-2 (heat shock; *Drosophila*; see Larochelle and Suter *Gene* 163:209-14 (1995)).

In addition, the following examples of promoters are induced by the presence or absence of light can be used in combination with those of the present invention: Topoisomerase II (pea; see Reddy *et al. Plant Mol Biol* 41:125-37 (1999)), chalcone synthase (soybean; see Wingender *et al. Mol Gen Genet* 218:315-22 (1989)) mdm2 gene (human tumor; see Saucedo *et al. Cell Growth Differ* 9:119-30 (1998)), Clock and BMAL1 (rat; see Namihira *et al. Neurosci Lett* 271:1-4 (1998)), PHYA (Arabidopsis; see Canton and Quail *Plant Physiol* 121:1207-16 (1999)), PRB-1b (tobacco; see Sessa *et al. Plant Mol Biol* 28:537-47 (1995)) and Ypr10 (common bean; see Walter *et al. Eur J Biochem* 239:281-93 (1996)).

Tissue-Specific Promoters

Alternatively, the plant promoter may direct expression of the polynucleotide of the invention in a specific tissue (tissue-specific promoters). Tissue specific promoters are

transcriptional control elements that are only active in particular cells or tissues at specific times during plant development, such as in vegetative tissues or reproductive tissues.

Promoters from the G564 or C541 genes of the invention are particularly useful for tissue-specific direction of gene expression so that a desired gene product is generated only or preferentially in suspensors, as described below.

Examples of tissue-specific promoters under developmental control include promoters that initiate transcription only (or primarily only) in certain tissues, such as vegetative tissues, e.g., roots or leaves, or reproductive tissues, such as fruit, ovules, seeds, pollen, pistils, flowers, or any embryonic tissue. Reproductive tissue-specific promoters may be, e.g., ovule-specific, embryo-specific, endosperm-specific, integument-specific, seed and seed coat-specific, pollen-specific, petal-specific, sepal-specific, or some combination thereof.

Suitable seed-specific promoters are derived from the following genes: MAC1 from maize, Sheridan (1996) *Genetics* 142:1009-1020; Cat3 from maize, GenBank No. L05934, Abler (1993) *Plant Mol. Biol.* 22:10131-1038; viviparous-1 from Arabidopsis, Genbank No. U93215; atmyc1 from Arabidopsis, Urao (1996) *Plant Mol. Biol.* 32:571-57; Conceicao (1994) *Plant* 5:493-505; napA from Brassica napus, GenBank No. J02798, Josefsson (1987) *JBL* 26:12196-1301; the napin gene family from Brassica napus, Sjobahl (1995) *Planta* 197:264-271.

The ovule-specific BEL1 gene described in Reiser (1995) *Cell* 83:735-742, GenBank No. U39944, can also be used. See also Ray (1994) *Proc. Natl. Acad. Sci. USA* 91:5761-5765. The egg and central cell specific FIE1 promoter is also a useful reproductive tissue-specific promoter.

Sepal and petal specific promoters are also used to express G564 nucleic acids in a reproductive tissue-specific manner. For example, the Arabidopsis floral homeotic gene APETALA1 (AP1) encodes a putative transcription factor that is expressed in young flower primordia, and later becomes localized to sepals and petals (see, e.g., Gustafson-Brown (1994) *Cell* 76:131-143; Mandel (1992) *Nature* 360:273-277). A related promoter, for AP2, a floral homeotic gene that is necessary for the normal development of sepals and petals in floral whorls, is also useful (see, e.g., Drews (1991) *Cell* 65:991-1002; Bowman (1991) *Plant Cell* 3:749-758). Another useful promoter is that controlling the expression of the unusual floral organs (ufo) gene of Arabidopsis, whose expression is restricted to the junction between sepal and petal primordia (Bossinger (1996) *Development* 122:1093-1102).

A maize pollen-specific promoter has been identified in maize (Guerrero (1990) *Mol. Gen. Genet.* 224:161-168). Other genes specifically expressed in pollen are described, e.g., by Wakeley (1998) *Plant Mol. Biol.* 37:187-192; Ficker (1998) *Mol. Gen. Genet.* 257:132-142; Kulikauskas (1997) *Plant Mol. Biol.* 34:809-814; Treacy (1997) *Plant Mol. Biol.* 34:603-611.

Other suitable promoters include those from genes encoding embryonic storage proteins. For example, the gene encoding the 2S storage protein from Brassica napus, Dasgupta (1993) *Gene* 133:301-302; the 2s seed storage protein gene family from Arabidopsis; the gene encoding oleosin 20kD from Brassica napus, GenBank No. M63985; the genes encoding oleosin A, Genbank No. U09118, and, oleosin B, Genbank No. U09119, from soybean; the gene encoding oleosin from Arabidopsis, Genbank No. Z17657; the gene encoding oleosin 18kD from maize, GenBank No. J05212, Lee (1994) *Plant Mol. Biol.* 26:1981-1987; and, the gene encoding low molecular weight sulphur rich protein from soybean, Choi (1995) *Mol Gen, Genet.* 246:266-268, can be used. The tissue specific E8 promoter from tomato is particularly useful for directing gene expression so that a desired gene product is located in fruits.

A tomato promoter active during fruit ripening, senescence and abscission of leaves and, to a lesser extent, of flowers can be used (Blume (1997) *Plant J.* 12:731-746). Other exemplary promoters include the pistil specific promoter in the potato (*Solanum tuberosum* L.) SK2 gene, encoding a pistil-specific basic endochitinase (Ficker (1997) *Plant Mol. Biol.* 35:425-431); the Blec4 gene from pea (*Pisum sativum* cv. Alaska), active in epidermal tissue of vegetative and floral shoot apices of transgenic alfalfa. This makes it a useful tool to target the expression of foreign genes to the epidermal layer of actively growing shoots.

A variety of promoters specifically active in vegetative tissues, such as leaves, stems, roots and tubers, can also be used to express the G564 or C541 nucleic acids of the invention. For example, promoters controlling patatin, the major storage protein of the potato tuber, can be used, see, e.g., Kim (1994) *Plant Mol. Biol.* 26:603-615; Martin (1997) *Plant J.* 11:53-62. The ORF13 promoter from *Agrobacterium rhizogenes* which exhibits high activity in roots can also be used (Hansen (1997) *Mol. Gen. Genet.* 254:337-343). Other useful vegetative tissue-specific promoters include: the tarin promoter of the gene encoding a globulin from a major taro (*Colocasia esculenta* L. Schott) corm protein family, tarin (Bezerra (1995) *Plant Mol. Biol.* 28:137-144); the curculin promoter active during taro corm development (de Castro (1992) *Plant Cell* 4:1549-1559) and the promoter for the tobacco

root-specific gene TobRB7, whose expression is localized to root meristem and immature central cylinder regions (Yamamoto (1991) *Plant Cell* 3:371-382).

Leaf-specific promoters, such as the ribulose biphosphate carboxylase (RBCS) promoters can be used. For example, the tomato RBCS1, RBCS2 and RBCS3A genes are expressed in leaves and light-grown seedlings, only RBCS1 and RBCS2 are expressed in developing tomato fruits (Meier (1997) *FEBS Lett.* 415:91-95). A ribulose biphosphate carboxylase promoters expressed almost exclusively in mesophyll cells in leaf blades and leaf sheaths at high levels, described by Matsuoka (1994) *Plant J.* 6:311-319, can be used.

Another leaf-specific promoter is the light harvesting chlorophyll a/b binding protein gene promoter, see, e.g., Shiina (1997) *Plant Physiol.* 115:477-483; Casal (1998) *Plant Physiol.* 116:1533-1538. The *Arabidopsis thaliana* myb-related gene promoter (Atmyb5) described by Li (1996) *FEBS Lett.* 379:117-121, is leaf-specific. The Atmyb5 promoter is expressed in developing leaf trichomes, stipules, and epidermal cells on the margins of young rosette and cauline leaves, and in immature seeds. Atmyb5 mRNA appears between fertilization and the 16 cell stage of embryo development and persists beyond the heart stage. A leaf promoter identified in maize by Busk (1997) *Plant J.* 11:1285-1295, can also be used.

Another class of useful vegetative tissue-specific promoters are meristematic (root tip and shoot apex) promoters. For example, the "SHOOTMERISTEMLESS" and "SCARECROW" promoters, which are active in the developing shoot or root apical meristems, described by Di Lorenzo (1996) *Cell* 86:423-433; and, Long (1996) *Nature* 379:66-69; can be used. Another useful promoter is that which controls the expression of 3-hydroxy-3- methylglutaryl coenzyme A reductase HMG2 gene, whose expression is restricted to meristematic and floral (secretory zone of the stigma, mature pollen grains, gynoecium vascular tissue, and fertilized ovules) tissues (see, e.g., Enjuto (1995) *Plant Cell.* 7:517-527). Also useful are kn1-related genes from maize and other species which show meristem-specific expression, see, e.g., Granger (1996) *Plant Mol. Biol.* 31:373-378; Kerstetter (1994) *Plant Cell* 6:1877-1887; Hake (1995) *Philos. Trans. R. Soc. Lond. B. Biol. Sci.* 350:45-51. For example, the *Arabidopsis thaliana* KNAT1 promoter. In the shoot apex, KNAT1 transcript is localized primarily to the shoot apical meristem; the expression of KNAT1 in the shoot meristem decreases during the floral transition and is restricted to the cortex of the inflorescence stem (see, e.g., Lincoln (1994) *Plant Cell* 6:1859-1876).

One of skill will recognize that a tissue-specific promoter may drive expression of operably linked sequences in tissues other than the target tissue. Thus, as used

herein a tissue-specific promoter is one that drives expression preferentially in the target tissue, but may also lead to some expression in other tissues as well.

In another embodiment, a G564 nucleic acid is expressed through a transposable element. This allows for constitutive, yet periodic and infrequent expression of the constitutively active polypeptide. The invention also provides for use of tissue-specific promoters derived from viruses which can include, e.g., the tobamovirus subgenomic promoter (Kumagai (1995) *Proc. Natl. Acad. Sci. USA* 92:1679-1683; the rice tungro bacilliform virus (RTBV), which replicates only in phloem cells in infected rice plants, with its promoter which drives strong phloem-specific reporter gene expression; the cassava vein mosaic virus (CVMV) promoter, with highest activity in vascular elements, in leaf mesophyll cells, and in root tips (Verdaguer (1996) *Plant Mol. Biol.* 31:1129-1139).

The promoters and control elements of the following genes can also be used in combination with the present invention to confer tissue specificity: MipB (iceplant; Yamada *et al. Plant Cell* 7:1129-42 (1995)) and SUCS (root nodules; broadbean; Kuster *et al. Mol Plant Microbe Interact* 6:507-14 (1993)) for roots, OsSUT1 (rice; Hirose *et al. Plant Cell Physiol* 38:1389-96 (1997)) for leaves, Msg (soybean; Stomvik *et al. Plant Mol Biol* 41:217-31 (1999)) for siliques, cell1 (Arabidopsis; Shani *et al. Plant Mol Biol* 34(6):837-42 (1997)) and ACT11 (Arabidopsis; Huang *et al. Plant Mol Biol* 33:125-39 (1997)) for inflorescence.

Still other promoters are affected by hormones or participate in specific physiological processes, which can be used in combination with those of present invention. Some examples are the ACC synthase gene that is induced differently by ethylene and brassinosteroids (mung bean; Yi *et al. Plant Mol Biol* 41:443-54 (1999)), the TAPG1 gene that is active during abscission (tomato; Kalaitzis *et al. Plant Mol Biol* 28:647-56 (1995)), and the 1-aminocyclopropane-1-carboxylate synthase gene (carnation; Jones *et al. Plant Mol Biol* 28:505-12 (1995)) and the CP-2/cathepsin L gene (rat; Kim and Wright. *Biol Reprod* 57:1467-77 (1997)), both active during senescence.

H. Vectors

Vectors are a useful component of the present invention. In particular, the present promoters and/or promoter control elements may be delivered to a system such as a cell by way of a vector. For the purposes of this invention, such delivery may range from simply introducing the promoter or promoter control element by itself randomly into a cell to integration of a cloning vector containing the present promoter or promoter control element. Thus, a vector need not be limited to a DNA molecule such as a plasmid, cosmid or bacterial

phage that has the capability of replicating autonomously in a host cell. All other manner of delivery of the promoters and promoter control elements of the invention are envisioned. The various T-DNA vector types are a preferred vector for use with the present invention. Many useful vectors are commercially available.

It may also be useful to attach a marker sequence to the present promoter and promoter control element in order to determine activity of such sequences. Marker sequences typically include genes that provide antibiotic resistance, such as tetracycline resistance, hygromycin resistance or ampicillin resistance, or provide herbicide resistance. Specific selectable marker genes may be used to confer resistance to herbicides such as glyphosate, glufosinate or broxynil (Comai *et al.*, *Nature* 317: 741-744 (1985); Gordon-Kamm *et al.*, *Plant Cell* 2: 603-618 (1990); and Stalker *et al.*, *Science* 242: 419-423 (1988)). Other marker genes exist which provide hormone responsiveness.

(1) Modification of Transcription by Promoters and Promoter Control Elements

The promoter or promoter control element of the present invention may be operably linked to a polynucleotide to be transcribed. In this manner, the promoter or promoter control element may modify transcription by modulate transcript levels of that polynucleotide when inserted into a genome.

However, prior to insertion into a genome, the promoter or promoter control element need not be linked, operably or otherwise, to a polynucleotide to be transcribed. For example, the promoter or promoter control element may be inserted alone into the genome in front of a polynucleotide already present in the genome. In this manner, the promoter or promoter control element may modulate the transcription of a polynucleotide that was already present in the genome. This polynucleotide may be native to the genome or inserted at an earlier time.

Alternatively, the promoter or promoter control element may be inserted into a genome alone to modulate transcription. *See*, for example, Vaucheret, H *et al.* (1998) *Plant J* 16: 651-659. Rather, the promoter or promoter control element may be simply inserted into a genome or maintained extrachromosomally as a way to divert transcription resources of the system to itself. This approach may be used to down-regulate the transcript levels of a group of polynucleotide(s).

(2) Polynucleotides to be Transcribed

The nature of the polynucleotide to be transcribed is not limited. Specifically, the polynucleotide may include sequences which will have activity as RNA as well as sequences which result in a polypeptide product. These sequences may include, but are not limited to antisense sequences, ribozyme sequences, spliceosomes, amino acid coding sequences, and fragments thereof.

Specific coding sequences may include, but are not limited to endogenous proteins or fragments thereof, or heterologous proteins including marker genes or fragments thereof.

Promoters and control elements of the present invention are useful for modulating metabolic or catabolic processes. Such processes include, but are not limited to, secondary product metabolism, amino acid synthesis, seed protein storage, oil development, pest defense and nitrogen usage. Some examples of genes, transcripts and peptides or polypeptides participating in these processes, which can be modulated by the present invention: are tryptophan decarboxylase (tdc) and strictosidine synthase (str1), dihydrodipicolinate synthase (DHDPS) and aspartate kinase (AK), 2S albumin and alpha-, beta-, and gamma-zeins, ricinoleate and 3-ketoacyl-ACP synthase (KAS), *Bacillus thuringiensis* (Bt) insecticidal protein, cowpea trypsin inhibitor (CpTI), asparagine synthetase and nitrite reductase. Alternatively, expression constructs can be used to inhibit expression of these peptides and polypeptides by incorporating the promoters in constructs for antisense use, co-suppression use or for the production of dominant negative mutations.

(3) Other Regulatory Elements

As explained above, several types of regulatory elements exist concerning transcription regulation. Each of these regulatory elements may be combined with the present vector if desired.

(4) Other Components of Vectors

Translation of eukaryotic mRNA is often initiated at the codon which encodes the first methionine. Thus, when constructing a recombinant polynucleotide according to the present invention for expressing a protein product, it is preferable to ensure that the linkage between the 3' portion, preferably including the TATA box, of the promoter and the polynucleotide to be transcribed, or a functional derivative thereof, does not contain any intervening codons which are capable of encoding a methionine.

The vector of the present invention may contain additional components. For example, an origin of replication allows for replication of the vector in a host cell. Additionally, homologous sequences flanking a specific sequence allows for specific recombination of the specific sequence at a desired location in the target genome. T-DNA sequences also allow for insertion of a specific sequence randomly into a target genome.

The vector may also be provided with a plurality of restriction sites for insertion of a polynucleotide to be transcribed as well as the promoter and/or promoter control elements of the present invention. The vector may additionally contain selectable marker genes. The vector may also contain a transcriptional and translational initiation region, and a transcriptional and translational termination region functional in the host cell. The termination region may be native with the transcriptional initiation region, may be native with the polynucleotide to be transcribed, or may be derived from another source. Convenient termination regions are available from the Ti-plasmid of *A. tumefaciens*, such as the octopine synthase and nopaline synthase termination regions. See also, Guerineau *et al.*, *Mol. Gen. Genet.* 262:141-144 (1991); Proudfoot, *Cell* 64:671-674 (1991); Sanfacon *et al.*, *Genes Dev.* 5:141-149 (1991); Mogen *et al.*, *Plant Cell* 2:1261-1272 (1990); Munroe *et al.*, *Gene* 91:151-158 (1990); Ballas *et al.*, *Nucleic Acids Res.* 17:7891-7903 (1989); Joshi *et al.*, *Nucleic Acid Res.* 15:9627-9639 (1987).

Where appropriate, the polynucleotide to be transcribed may be optimized for increased expression in a certain host cell. For example, the polynucleotide can be synthesized using preferred codons for improved transcription and translation. See U.S. Patent Nos. 5,380,831, 5,436,391; see also Murray *et al.*, *Nucleic Acids Res.* 17:477-498 (1989).

Additional sequence modifications include elimination of sequences encoding spurious polyadenylation signals, exon intron splice site signals, transposon-like repeats, and other such sequences well characterized as deleterious to expression. The G-C content of the polynucleotide may be adjusted to levels average for a given cellular host, as calculated by reference to known genes expressed in the host cell. The polynucleotide sequence may be modified to avoid hairpin secondary mRNA structures.

A general description of expression vectors and reporter genes can be found in Gruber, *et al.*, "Vectors for Plant Transformation, in *Methods in Plant Molecular Biology & Biotechnology*" in *METHODS IN PLANT MOLECULAR BIOLOGY & BIOTECHNOLOGY*, (Glich *et al.*, eds. 1993) pp. 89-119. Moreover GUS expression vectors and GUS gene cassettes are available from Clontech Laboratories, Inc., Palo Alto, California while luciferase expression

vectors and luciferase gene cassettes are available from Promega Corp. (Madison, Wisconsin). GFP vectors are available from Aurora Biosciences.

I. Polynucleotide Insertion Into A Host Cell

The polynucleotides according to the present invention can be inserted into a host cell. A host cell includes but is not limited to a plant, mammalian, insect, yeast, and prokaryotic cell, preferably a plant cell.

The method of insertion into the host cell genome is chosen based on convenience. For example, the insertion into the host cell genome may either be accomplished by vectors which integrate into the host cell genome or by vectors which exist independent of the host cell genome.

The nucleic acids of the invention can be used to confer desired traits on essentially any plant. Thus, the invention has use over a broad range of plants, including species from the genera Asparagus, Atropa, Avena, Brassica, Citrus, Citrullus, Capsicum, Cucumis, Cucurbita, Daucus, Fragaria, Glycine, Gossypium, Helianthus, Heterocallis, Hordeum, Hyoscyamus, Lactuca, Linum, Lolium, Lycopersicon, Malus, Manihot, Majorana, Medicago, Nicotiana, Oryza, Panieum, Pannesetum, Persea, Pisum, Pyrus, Prunus, Raphanus, Secale, Senecio, Sinapis, Solanum, Sorghum, Trigonella, Triticum, Vitis, Vigna, and, Zea.

(1) Polynucleotides Autonomous of the Host Genome

The polynucleotides the present invention can exist autonomous or independent of the host cell genome. Vectors of these types are known in the art and include, for example, certain type of non-integrating viral vectors, autonomously replicating plasmids, artificial chromosomes, and the like.

Additionally, in some cases transient expression of a polynucleotide may be desired.

(2) Polynucleotides Integrated into the Host Genome

The promoter sequences, promoter control elements or vectors of the present invention may be transformed into host cells. These transformations may be into protoplasts or intact tissues or isolated cells. Preferably expression vectors are introduced into intact tissue. General methods of culturing plant tissues are provided for example by Maki *et al.* "Procedures for Introducing Foreign DNA into Plants" in METHODS IN PLANT MOLECULAR BIOLOGY & BIOTECHNOLOGY, (Glich *et al.*, eds. 1993) pp. 67-88; and by Phillips *et al.* "Cell-

Tissue Culture and In-Vitro Manipulation" in CORN & CORN IMPROVEMENT, 3rd Edition (Sprague *et al.*, eds. 1998) pp. 345-387.

Methods of introducing polynucleotides into plant tissue include the direct infection or co-cultivation of plant cell with *Agrobacterium tumefaciens*, Horsch *et al.*, Science, 227:1229 (1985). Descriptions of *Agrobacterium* vector systems and methods for *Agrobacterium*-mediated gene transfer provided by Gruber *et al.* supra.

Alternatively, polynucleotides are introduced into plant cells or other plant tissues using a direct gene transfer method such as microprojectile-mediated delivery, DNA injection, electroporation and the like. More preferably polynucleotides are introduced into plant tissues using the microprojectile media delivery with the biolistic device. See, for example, Tomes *et al.*, "Direct DNA transfer into intact plant cells via microprojectile bombardment" in PLANT CELL, TISSUE AND ORGAN CULTURE: FUNDAMENTAL METHODS (: Gamborg and Phillips, eds. 1995).

In another embodiment of the current invention, expression constructs can be used for gene expression in callus culture for the purpose of expressing marker genes encoding peptides or polypeptides which allow identification of transformed plants. Here, a promoter that is operatively linked to a polynucleotide to be transcribed is transformed into plant cells and the transformed tissue is then placed on callus-inducing media. If the transformation is conducted with leaf discs, for example, callus will initiate along the cut edges. Once callus growth has initiated, callus cells can be transferred to callus shoot-inducing or callus root-inducing media. Gene expression will occur in the callus cells developing on the appropriate media: callus root-inducing promoters will be activated on callus root-inducing media, etc. Examples of such peptides or polypeptides useful as transformation markers include, but are not limited to barstar, glyphosate, chloramphenicol acetyltransferase (CAT), kanamycin, spectinomycin, streptomycin or other antibiotic resistance enzymes, green fluorescent protein (GFP), and β -glucuronidase (GUS), etc. Some of the promoters of the invention will also be capable of sustaining expression in some tissues or organs after the initiation or completion of regeneration. Examples of these tissues or organs are somatic embryos, cotyledon, hypocotyl, epicotyl, leaf, stems, roots, flowers and seed.

Integration into the host cell genome also can be accomplished by methods known in the art, for example, by the homologous sequences or T-DNA discussed above or using the cre-lox system (A.C. Vergunst *et al.*, *Plant Mol. Biol.* 38:393 (1998)).

J. Utility

Common Uses

The polynucleotides of the invention have a variety of uses. For example, modulation of expression of the gene products of the invention can be used to modulate suspensor cell and/or embryo size, shape or rates of development.

The suspensor-specific promoters of the invention are also useful for expression of any number of polynucleotides in a suspensor-specific fashion. Exemplary gene products that can be expressed under the control of the promoters of the invention include toxic gene products. In some embodiments, toxic gene products are also expressed in the embryo under the control of the same or a second promoter. By preventing development of the suspensor cell and/or the embryo, plants with modulated fertility and/or that produce seedless fruit can be developed.

Examples of toxic genes include, e.g., those which produce toxic substances, disrupt cell function, suppress genes required by the cell (such as by using anti-sense, sense suppression, or ribozymes), and disruption of mitochondrial function. Particular examples include, barnase (Sancho & Fersht, *J. Mol. Biol.* 224:741-47 (1992)), diphtheria toxin (DT) A chain, which adenoribosylates elongation factor EF-2, thus blocking protein synthesis (Herrera *et al.*, *Proc. Natl. Acad. Sci., USA* 91:12999-13003 (1994)), and the thymidine kinase (tk) gene, which provides a conditional cell-lethal function, requiring the presence of a nucleoside analog such as ganciclovir for lethality (Brady *et al.*, *Proc. Natl. Acad. Sci., USA* 91:365-69 (1994)).

Alternatively, growth regulators such as gene products that modulate gibberellin expression, can be specifically expressed within the suspensor, thereby modulating (e.g., increasing or decreasing) the attached embryo's size, shape or rate of development.

An additional utility includes the expression of gene products that induce embryonic features to the suspensor cell, thereby leading to the development of a second embryo. Examples of the gene products that induce embryonic features include the *LEC1* (see, e.g., Lotan, *et al. Cell* 93(7):1195-205 (1998)).

In yet another use, nucleic acids of the invention can be used in the development of apomictic plant lines (*i.e.*, plants in which asexual reproductive processes occur in the ovule, see, Koltunow, *A. Plant Cell* 5: 1425-1437 (1993) for a discussion of apomixis). Apomixis provides a novel means to select and fix complex heterozygous

genotypes that cannot be easily maintained by traditional breeding. Thus, for instance, new hybrid lines with desired traits (e.g., hybrid vigor) can be obtained and readily maintained.

In yet another use, expression cassettes comprising the promoter polynucleotides of the invention can be used to express genes that result in apomictic plants.

5 Examples of genes useful in creating apomictic plant include LEC1 nucleic acids as described by Lotan, *et al. Cell* 93: 1195-1205 (1998) and in USSN 09/026,221 as well as FIE and MEDEA nucleic acids as described in Ohad *et al., Plant Cell* 11:407-415 (1999); Grossniklaus *et al., Science* 280:446-450 (1998) and USSN 09/177,249. In these
embodiments, constructs providing expression of a LEC1, FIE, MEDEA or other nucleic
10 acids capable of inducing apomictic fruit are used alone or in combination.

The following examples are provided for a further understanding of the invention, however, the invention is not to be construed as limited thereto.

EXAMPLES

MATERIALS AND METHODS

Plant materials and maintenance

Seeds of the day neutral Scarlet Runner Bean cultivar 'Hammond's Dwarf Red Flower' (Vermont Bean Seed Company, Fair Haven, Vermont; Nagl, 1990) were
20 germinated in a soil mixture of vermiculite, perlite, sandy-loam soil, sphagnum peat moss, and plaster sand respectively at a ratio of 3:3:2:2:2. Plants were maintained in a 16:8 hour light/dark cycle in the greenhouse. Flowers were hand-pollinated by lightly brushing the stigma with a watercolor brush containing pollen. Hand-pollinated flowers were tagged and seeds were harvested at specific days after pollination.

Suspensor isolation

25 The micropylar half of a 6 days after pollination (DAP) seed was cut and placed upright on its cut side under a dissecting microscope. Approximately 1 mm was sliced from the left and right sides of the seed coat "flat face." The seed was turned on its "flat face" and the remaining seed coat and endosperm were removed from the exposed embryo
30 proper. The entire embryo was isolated and then the suspensor was separated from the embryo proper by microdissection. Generally, ten suspensors were isolated per hour.

RNA isolation and gel blot analysis

Polysomal RNAs were isolated according to the procedure of Cox and Goldberg (1988). Poly(A) mRNA was isolated from total polysomal RNA using the PolyA Tract® mRNA isolation system (Promega: Madison, WI) and the protocol supplied by the manufacturer. Total RNAs, used for the Differential Display Reverse Transcription Polymerase Chain Reaction (DD-RT-PCR) and RNA gel blot experiments, were isolated using the RNeasy® plant total RNA kit (Qiagen: Chatsworth, CA). RNAs were treated with RNase-free DNase (Boehringer Mannheim: Indianapolis, IN) following the protocol of Ausubel *et al.* (1992). RNA gel blots were carried out as described by Sambrook *et al.* (1989). ³²P-labeled DNA probes for the RNA gel blots were prepared by the random-priming procedure of Feinberg and Vogelstein (1984).

cDNA library construction and screening

A cDNA library of 5-9 DAP Scarlet Runner Bean seeds containing globular-stage embryos was constructed using the ZAP Express® cDNA synthesis kit (Stratagene: La Jolla, CA). Poly(A) mRNA was used as a template to generate first-strand cDNA using MMLV reverse transcriptase and a 50-base oligonucleotide linker-primer [5'-(GA)₁₀ACTAGTCTCGAG(T)₁₈-3']. Double-strand cDNAs were blunt-ended and ligated to an EcoRI adapter. After phosphorylation of EcoRI 5' ends, the cDNAs were digested with XhoI and size-fractionated on a Sephacryl S-400 column to exclude cDNAs that were smaller than 250 bp. The fractionated cDNAs were ligated to the λZAP vector. About 3,000 recombinants from the unamplified library were differentially screened with ³²P-labeled first-strand cDNAs generated from: (1) 5-9 DAP seed micropylar region poly(A) mRNA and (2) leaf poly(A) mRNA. cDNA clones representing mRNAs preferentially present in the micropylar region were screened two more times following the strategy used in the primary screen.

Differential display reverse transcription polymerase chain reaction

Differential display procedures of Liang and Pardee (Liang, P., *et al.*, *Science*, 257:967-971 (1992)) were followed using the RNImage™ kit (GenHunter Corp.: Nashville, TN). Differential display reactions were carried out using total RNA templates from: (1) 6-8 DAP dissected suspensors of globular-stage embryos, (2) 6 DAY embryo-containing micropylar seed regions, (3) 6 DAP non-embryo-containing chalazal seed regions, (4) 6-8 DAP isolated globular-stage embryo propers, (5) leaves, (6) ovules, (7) 2 DAY whole seeds, and (8) 3 DAP whole seeds. Briefly, first-strand cDNAs were generated by reverse

transcription (RT) of 200 ng of total RNA using MMLV reverse transcriptase and an anchor/reverse primer (G primer: 5'-AAGCT₁₁G-3' or C primer: 5'-AAGCT₁₁C-3'). Aliquots of the first-strand cDNAs were used as templates for the polymerase chain reaction (PCR) using combinations of forward and anchor/reverse primers in the presence of ³³P-dCTP and AmpliTaq® polymerase (Perkin Elmer; Branchburg, NJ). The forward primers used were: H-AP49, 5'-AAGCTTTAGTCCA-3'; H-AP50, 5'-AAGCTTTGAGACT-3'; H-AP51, 5'-AAGCTTCGAAATG-3'; H-AP52, 5'-AAGCTTGACCTTT-3'; H-AP53, 5'-AAGCTTCCTCTAT-3'; H-AP54, 5'-AAGCTTTTGAGGT-3'; H-AP55, 5'-AAGCTTACGTAG-3', and H-AP56, 5'-AAGCTTATGAAGG-3', where H-AP refers to the primers supplied by the RNAimage™ kit. The RT-PCR products were size-fractionated in a 6% acrylamide gel and visualized by autoradiography.

Candidate suspensor-specific cDNAs as bands were identified that were (1) over 200 bp in size, (2) present at the same position in lanes containing cDNAs amplified from 6-8 DAP suspensor and micropylar-region mRNAs, and (3) absent in lanes containing cDNAs amplified from chalazal region, embryo proper, and leaf mRNAs. Isolated cDNA fragments were PCR-amplified, cloned into the pCR2.1® vector (Invitrogen: San Diego, CA), and sequenced. cDNAs were designated with (1) a C or G, indicating the anchor/reverse primer used, (2) a two-digit number between 49 and 56, indicating the forward primer used, and (3) a one-digit number indicating, the band position on the DD-RT-PCR gel. For example, C541 represents a cDNA band that was amplified by a C anchor/reverse primer, an H-AP54 forward primer, and that was in position number 1 on the DD-RT-PCR gel.

Gel blot analysis of PCR-amplified population cDNAs

For pre-screening of differential display cDNA clones, PCR-amplified cDNAs from different mRNA populations were generated following the procedures of Kelly *et al.* (1990), with minor modifications. Suspensor (6 DAP), ovule, 2 DAP seed, 3 DAP seed, 6 DAP micropylar region, 6 DAP chalazal region, and leaf total RNAs were isolated. First-strand cDNA was generated from 5 µg of each RNA using MMLV reverse transcriptase and 50 ng/µl of oligo(dT₂₀) as primer. The first-strand cDNAs were 3' tailed with poly(dA) using terminal transferase. PCR amplifications were carried out using tailed first-strand cDNAs as templates and 2 µM of dT₂₀dN (where dN = dG, dC, dA, or dT) as primer in 100 µl containing 20 mM Tris (pH 8.4), 50 mM KCl, 1 mM MgCl₂, and 0.2 µM dNTPs at 94°C/1 minute, 42°C/2 minutes, and 72°C/5 minutes for 30 cycles, followed by a 10 minute extension at 72°C. A 1 µl aliquot from each reaction was used to perform another round of

amplification using the same conditions. The reactions were extracted with phenol/chloroform and precipitated in ethanol. An aliquot equivalent to 1 µg from each reaction was size-fractionated in a 1% agarose gel, which was then used for DNA gel blot analysis according to the procedures of Sambrook *et al.*, *supra*.

5 DNA sequencing and analysis

DNA sequencing was performed following the dideoxy sequencing procedures recommended by USBiochemicals (Cleveland, OH). For genomic clone pG564g7.2.79, unidirectional, nested deletion set was prepared using the Erase-a-Base® system (Promega: Madison, WI). Compilation and analysis of sequences were carried out using the Wisconsin Genetics Computer Group (GCG) software. ORFs and exon-intron junctions were identified by using GENSCAN (<http://ccr-081.mit.edu/GENSCAN.html>; Burge, C., *et al.*, *Journal of Molecular Biology*, 268:78-94 (1997)). The G564 intron-exon junctions were confirmed by comparing the cDNA and gene sequences. Protein sorting sequences were identified using PSORT (<http://psort.nibb.ac.jp>; Nakai, K., *et al.*, *Genomics*, 14:897-911 (1992)). DNA and protein sequence comparisons were performed using the NCBI Genbank BLAST programs (<http://www.ncbi.nlm.nih.gov>; Altschul, S. F., *et al.*, *Nucl. Acids Res.*, 25:3389-3402 (1997)). The complete C541 and 0564 cDNA sequences were based on sequences from (1) DD-RT-PCR cDNA clones, (2) cDNA clones isolated from a 5-9 DAP seed cDNA library, and (3) from cDNAs generated from 5' random amplification of cDNA ends (RACE-RT-PCR; Chenchik, A., *et al.*, *Clontechniques*, 10:5-8 (1995)).

In situ hybridization

In situ hybridization studies were carried out as described by Cox and Goldberg (Cox, K. H., *et al.*, *PLANT MOLECULAR BIOLOGY: A PRACTICAL APPROACH* (C. H. Shaw, ed. 1988) pp. 1-34) and Yadegari *et al.* (Yadegari, R., *et al.*, *Plant Cell*, 6:1713-1729 (1994)) with minor modifications. Briefly, for Scarlet Runner Bean, unfertilized ovules and individual seeds (4-7 DAP) were harvested from pods, and seeds were cut at their chalazal ends before fixing to enhance penetration of the fixative. For tobacco, seeds up to 7 DAP were collected while still attached to the placenta. Older tobacco seeds were separated from the placenta prior to collection. Tissues were fixed overnight at 4°C in 1% glutaraldehyde solution prepared in 0.1 M phosphate buffer (pH 7.0) (Meyerowitz, E. M., *Plant Mol. Biol. Rep.*, 5:242-250 (1987)), dehydrated, cleared, and embedded in paraffin. Eight to 10 µm sections were hybridized to ³³P-labeled sense or anti-sense RNA probes at a specific activity of 4-5 x 10⁸ dpm/µg. After hybridization and emulsion development, sections were stained

with 0.05% toluidine blue in 0.05% borate solution. Photographs were taken using either bright-field or dark-field illumination with a compound microscope (Olympus BH2: Olympus Corporation, Lake Success, NY). The photographs were digitized, adjusted for optimum silver grain resolution using the KPT-Equilizer program (Metacreations Corp., Carpinteria, CA), and assembled in Adobe Photoshop 5.0 (Adobe Systems Inc., San Jose, CA).

Light microscopy

Bright-field microscopy

Seeds and unfertilized ovules from Scarlet Runner Bean were collected as described for in situ hybridization and fixed overnight in 5% glutaraldehyde, 0.1 M phosphate buffer (pH 7.0), and 0.01% Triton X-100 at 4°C. After dehydration, samples were embedded in Spurr's (Spurr, 1969) plastic resin (Polysciences: Warrington, PA). 1 µm thick sections were stained for 18 to 20 minutes at 42°C with 0.05% toluidine blue in 0.05% borate solution. Bright-field photographs were taken with Kodak Gold 100 film (ISO 100/21°) using a compound microscope (Olympus BH-2: Olympus Corporation, Lake Success, NY).

Whole mount microscopy

Dark-field photographs of seeds were taken using a dissecting microscope (Olympus SZH). Dark-field and bright-field photographs of dissected embryos were taken using a compound microscope (Olympus BH-2).

G564/GUS construction and tobacco plant transformation

A 21 kb *G564* genomic clone was isolated from a Scarlet Runner Bean λDASHII (Stratagene: La Jolla, CA) genomic library by screening with a ³²P-labeled *G564* cDNA clone. A 7 kb genomic fragment was recloned in pBluescript (Stratagene: La Jolla, CA) generating plasmid p*G564g7.2.79*. 4.8 kb of this plasmid was sequenced to confirm that the sequence of the coding region corresponded to that of the *G564* cDNA clone. The entire *G564g7.2.79* genomic clone was transferred into pGV1501AN, a pGV1500-derived plant transformation vector (DeBlare, R., *et al.*, *Methods in Enzymology*, 153:277-292 (1987)).

The region surrounding the ATG start codon in *G564g7.2.79* was converted into an SphI endonuclease restriction site by PCR using a T3 primer and a mutagenic oligo (5'-ATTGGACTGCATGCTTACGCTAGTCTGTGCAGAG-3'). A 4.2 kb *G564* promoter region was cloned in the SphI site upstream of the *E coli β-Glucoronidase (GUS)* gene coding region (Jefferson, R. A., *et al.*, *EMBO. J.*, 6(13):3901-3907 (1987)) in pGEM5*GUS*. After cloning, the *G564* promoter region was re-sequenced. pGEMSG*GUS* was constructed by

inserting the *GUS* coding region and the Ti-plasmid *gene 7 3'* end from *TPI2/GUS* gene (Drews, G. N., *et al.*, *Plant Cell*, 4:1383-1404 (1992)) into the *NcoI/NotI* sites of pGEM5 (Promega: Madison, WI). The *G564/GUS* gene was transferred to the pHYGA (Hygromycin^R) plant transformation vector (Klucher, K. M., *et al.*, *Plant Cell*, 8:137-153 (1996)). Tobacco plants were transformed and regenerated using the leaf disk procedure of Horsch *et al.* (Horsch, *et al.*, *Science*, 227:1229-1231 (1985)).

GUS histochemical assay

Transgenic tobacco seeds were harvested at different stages of development (Barker, S. J., *et al.*, *Proc. Natl Acad. Sci. USA*, 85:458-462 (1988)). Embryos were dissected from seeds in 50 mM sodium phosphate (pH 7.0). Dissected embryos were incubated in GUS assay buffer [50 mM sodium phosphate (pH 7.0), 0.1 % Triton X-100, 0.5 mM ferricyanide, 0.5 mM ferrocyanide, 2mM 5-bromo-4chloro-3indolyl- β D-glucuronide] for 30 minutes to 16 hours at room temperature (Jefferson, R. A., *et al.*, *EMBO. J.*, 6(13):3901-3907 (1987)). Embryos were photographed under bright-field or dark-field illumination using a compound BH2 Olympus microscope.

RESULTS

The Scarlet Runner Bean embryo forms a "giant" suspensor early in development

The early developmental stages of Scarlet Runner Bean embryogenesis were characterized to link these stages to morphological markers of the developing seed and to specific times after pollination. Table 1 summarizes the morphological characteristics of the unfertilized ovule and developing seeds from 0 DAP until maturity at 35 DAP. From the ovule until 7 DAP, the seed length increased from 0.75 mm to 2-4 mm and the seed gradually adopted a green color (Table 1). At 11 DAP, the seed began to acquire red pigmentation in the area contiguous to the hilum region (Table 1) and the red color gradually spread and covered the entire seed by 20-25 DAP (Table 1). At 25 DAP, the seed length had increased and was 15 mm (Table 1). At 35 DAP, the mature dry seed had a purple seed coat with magenta streaks near the hilum and was 20 mm in length (Table 1).

The embryonic stages corresponding to seeds at different DAP were characterized from micrographs of longitudinal sections of the micropylar region containing the embryo. In the unfertilized ovule, the egg cell was identified from the orientation of its nucleus and cytoplasmic-dense region towards the chalaza and its vacuolated region towards the micropyle. These cytological features were inverted in the adjacent synergids. The egg

cell and synergids were bordered by the central cell at their chalazal ends. At 2 DAP, the embryonic cells were irregularly organized, the apical and basal regions were morphologically indistinguishable, and endosperm had started to form. Just prior to globular stage (4 DAP), the suspensor of the filamentous embryo was distinguished from the embryo proper by its large and irregularly-shaped cells and was approximately 200-250 μm in length. By contrast, the embryo-proper cells were smaller and more uniform in size and shape.

The suspensor developed two distinct regions – a file of neck cells that connected suspensor to embryo proper and a set of large basal cells that protruded into the seed tissue. In the suspensor-basal region, the number of cells remained constant and the increase in length of the suspensor-basal region was mainly due to cell enlargement. The total suspensor length increased from 500 μm to 1000 μm , which was its maximum size (Table 1). The embryo proper increased in cell size and number, and developed from globular stage to heart stage, to cotyledon stage. At the cotyledon stage, the embryo proper was bigger than the suspensor and contained chlorophyll, whereas the suspensor remained white.

Globular embryos were dissected at the rate of approximately 10 per hour and collect separately the embryo-proper and suspensor regions (see Materials and Methods). Twenty micrograms of total RNA was isolated from 250 suspensors and 300 ng total RNA from 200 embryo-proper regions. Together, these data show that the suspensor of Scarlet Runner Bean embryo developed early in seed development (2-11 DAP) and that it was feasible to surgically dissect globular stage embryos into embryo-proper and suspensor regions in order to isolate region-specific embryo RNAs.

DD-RT-PCR of RNA from micro-dissected suspensor regions yields two suspensor-specific cDNA clones

Two strategies were used to identify suspensor-specific mRNAs (Materials and Methods): (1) differential screening of a 5-9 DAP seed cDNA library representing mRNAs present in seeds containing globular-stage embryos and (2) DD-RT-PCR (Liang, P., *et al.*, *Science*, 257:967-971 (1992)) of total RNA from micro-dissected suspensors of globular-stage embryos. Candidates for suspensor-specific cDNA clones were rescreened using: (1) DNA gel blots containing PCR-amplified population cDNAs (Materials and Methods) and (2) RNA gel blots (Materials and Methods).

Differential screening

In the first approach, two 'seed-specific' candidates for suspensor cDNA clones were identified, designated as SRB8 and SRB13, which hybridized with a 5-9 DAP micropylar-region seed cDNA probe, but not with a leaf cDNA probe (Materials and Methods). SRB8 and SRB13 were sequenced and used BLAST searches (Altschul, S. F., *et al.*, *Nucl. Acids Res.*, 25:3389-3402 (1997)) to show that the encoded proteins are homologous to ribosomal proteins and Bowman-Birk trypsin inhibitor, respectively (Materials and Methods).

DD-RT-PCR analysis

In the second approach, 25 candidate suspensor-specific cDNAs were identified that were displayed in the lane containing cDNAs amplified from 6 DAP suspensor RNA and in the lane containing cDNAs amplified from RNA of the micropylar half of 6 DAP seed, and that were not present in lanes containing cDNAs amplified from 6 DAP seed chalazal region RNA, globular-stage-embryo-proper RNA, and leaf RNA. All candidate cDNAs longer than 200 bp were cut from the gel, re-amplified, cloned, and sequenced (Materials and Methods).

Total cDNA gel blot analysis

Because the amount of RNA from the suspensor was too limited to screen a large number of clones by standard RNA blot analysis, a DNA gel blot procedure was devised using PCR-amplified population cDNAs (Kelly, A. J., *et al.*, *Plant Cell*, 2:963-972 (1990)) to pre-screen the candidate cDNA clones (Materials and Methods). Total cDNA blot analysis of SRB8 and SRB13 showed that they hybridized with 6 DAP suspensor cDNA, unfertilized ovule, 2 DAP seed, 3 DAP seed, 6 DAP seed micropylar region cDNAs, and 6 DAP seed chalazal region cDNA but not with leaf cDNA. In addition, three DD-RT-PCR cDNAs were identified that hybridized with suspensor and seed micropylar-region cDNAs, but did not hybridize with ovule, seed chalazal-region, and leaf cDNAs. These three clones were designated as G541, G564, and G563, and represented putative suspensor-specific cDNAs. Sequence analysis and homology searches with these cDNAs indicated that they were not related to any protein of known function. However, G564 and C541 proteins were predicted to be secreted or to be targeted to the vacuole, respectively (Materials and Methods).

RNA gel blot analysis

SRB8, SRB13, G564, C541, and G563 probes were hybridized to gel blots, containing 6 DAP suspensor RNA, unfertilized ovule RNA, 2 DAP seed RNA, 3 DAP seed RNA, 6 DAP seed micropylar region RNA, 6 DAP seed chalazal region RNA, and leaf RNA to verify the results of the total cDNA blots. SRB8 and SRB13 probes hybridized with unfertilized ovule and all seed tissue RNAs, but not with leaf RNA. The SRB8 probe yielded a stronger hybridization signal with micropylar-region RNA than with chalazal-region RNA. By contrast, the SRB13 probe produced a stronger signal with chalazal-region RNA as compared to micropylar-region RNA.

G564 and C541 probes did not hybridize with unfertilized ovule, 2 DAP seed, 3 DAP seed, 6 DAP chalazal region, and leaf RNAs. By contrast, G564 and C541 probes yielded a low signal with 6 DAP seed micropylar-region RNA. This signal was strongly amplified with suspensor RNA isolated from 6 DAP micropylar-region seed, suggesting that the lower signal with 6 DAP seed micropylar-region RNA was caused by dilution of the suspensor RNA by non-embryonic seed tissue RNA. G563 produced a similar hybridization pattern, but yielded equal hybridization signals with suspensor and 6 DAP micropylar RNAs. Together, these data showed that during seed development different patterns and levels of RNA accumulation occur. In addition, the higher hybridization signals from G564 and C541 probes with suspensor RNA versus micropylar RNA suggested that G564 and C541 cDNAs represent suspensor-specific mRNAs.

G564 and C541 are suspensor-specific markers

In situ hybridization was used to visualize directly regions that the G564, C541, G563, SRB8, and SRB13 mRNAs were localized in unfertilized ovules and 7 DAP seeds.

Localization of G564 and C541 mRNA

Dark field images of 7 DAP embryo sections hybridized with G564 and C541 anti-mRNA probes showed that G564 and C541 mRNAs were localized specifically in the suspensor. The G564 hybridization signal was spread evenly over the suspensor neck and basal cells. The C541 signal, on the other hand, was higher in the suspensor basal cells than in the suspensor neck cells. In addition, compared to the G564 probe, the C541 probe produced fewer hybridization grains, suggesting that the C541 mRNA is present at a lower prevalence than the G564 mRNA. No hybridization signal was detected above background level in the embryo proper, nor in any other cell or tissue of the developing seed. No G564 or

C541 hybridization signals above background were observed in any unfertilized ovule cell or tissue type, similar to that observed with the sense control probe.

Localization of G563 mRNA

The G563 anti-mRNA probe hybridized specifically with transcripts in the endothelial layer surrounding the embryo but not in the embryo or any other seed tissue. The G563 hybridization signal was first detected at 3 DAP. By contrast, no hybridization signal above background level was obtained in the chalazal endotheium, nor in the endothelium or any other tissue of the unfertilized ovule.

Localization of SRB8 and SRB13 mRNAs

The SRB8 and SRB13 mRNAs were highly prevalent within unfertilized ovule and seed, and were not localized exclusively within the suspensor. However, both mRNAs displayed different and changing accumulation patterns within pre- and post-fertilization ovule/seed. In the ovule, the SRB8 anti-mRNA probe detected transcripts in the endotheium and the epidermal layer. In addition, in the developing seed, SRBS hybridization grains accumulated to a high level in the endosperm and in the embryo. A stronger SRB8 hybridization signal was observed in the embryo proper than in the suspensor. The SRB13 anti-mRNA probe yielded hybridization signal in the outer integument of the unfertilized ovule and seed. Although SRB13 mRNA was present in the suspensor, its prevalence was not as high as in the integument.

Taken together, these data show that in the unfertilized ovule and developing seed various and partially overlapping transcript-accumulation patterns occur that change after fertilization has occurred. In addition, these results show that G563 mRNA is a marker for seed micropylar endothelium and that G564 and C541 mRNAs are suspensor-specific markers.

G564 and C541 are markers for the basal-region of the four-cell embryo

In situ hybridization was used to investigate the accumulation pattern of G564 and C541 mRNAs during embryo development. Before fertilization, no hybridization signal was obtained with either G564 or C541 anti-mRNA probes in the egg or the synergids, even after a 6-9 month emulsion exposure. After fertilization, and before the suspensor and embryo-proper region were morphologically distinguishable (2 DAP), the G564 and C541 anti-mRNA probes detected transcripts exclusively in the two basal cells of the four-cell embryo, but did not detect any transcripts in the two apical cells. From early globular stage, after 3 DAP, G564 and C541 transcripts were detectable in the suspensor and not in the

embryo proper. In addition, the higher concentration of C541 mRNA in the suspensor-basal region, compared with the suspensor-neck region.

The G564 mRNA accumulation pattern at later stages of embryo development was investigated in 23 DAP early-maturation-stage embryos. The dark field image of an axis and cotyledon section that was hybridized with a G564 anti-mRNA probe showed that G564 transcripts accumulated in the axis, but not in the cotyledons or in any other seed tissue.

Together, these data show that late G564 transcripts mark the embryo axis, and that G564 and C541 mRNAs are suspensor-specific markers. In addition, these results show that within two cell divisions after fertilization, G564 and C541 mRNAs mark the two basal cells of the four-cell embryo.

Basal-region specific G564 mRNA accumulation is transcriptionally regulated

The *G564* gene was isolated from a Scarlet Runner Bean genomic library to determine whether the basal-region-specific and suspensor-specific G564 mRNA accumulation pattern was regulated at the transcriptional or post-transcriptional levels. A 6.99 kb genomic fragment from the Scarlet Runner Bean was isolated. The *G564* coding region was 659 bp long, consisted of 2 exons of 107 and 388 bp, and contained one 164 bp intron. The 5' and 3' regions, included in the genomic fragment, were 4242 bp and 2085 bp in length respectively. In the 5' region, another gene, at position -4214 to -2588, similar to the *Arabidopsis Pol3* gene (accession no. AC005561) was identified.

G564 mRNA localization in transgenic tobacco embryos carrying the Scarlet Runner Bean G564 gene

The Scarlet Runner Bean *G564* genomic clone was introduced into tobacco and localized G564 mRNA accumulation in transgenic embryos to investigate whether the basal-region-specific and suspensor-specific G564 mRNA accumulation patterns were conserved in a heterologous plant. At the pie-globular embryo stage, similar to the Scarlet Runner Bean embryo, the G564 mRNA accumulated specifically in the embryo basal region, but not in the apical region. At this stage of tobacco embryo development the suspensor is distinguishable from the embryo proper. At the globular stage, the G564 mRNA was detected in the suspensor and in the hypophyseal region of the embryo proper. In heart- and torpedo-stage embryos, G564 transcripts accumulated in the axis similar to the G564 mRNA accumulation pattern in the Scarlet Runner Bean early maturation-stage embryo. In addition, G564 transcripts accumulated in the endosperm. No hybridization signal above background level was detected in non-transformed tobacco embryos. Together, these results suggested

that the basal-region-specific and suspensor-specific *G564* mRNA accumulation pattern is conserved across the plant kingdom and that all regulatory elements for correct suspensor-specific *G564* mRNA accumulation are contained within the 6.99 kb *G564* genomic clone. Analysis of the gene sequence indicated that the coding sequence was interrupted by an intron. As measured from the first identified nucleotide of the *G564* cDNA sequence (i.e., position 4242 of SEQ ID NO:2), the first exon is located from positions 1 to 107 and the second exon from positions 271-659.

G564/GUS expression in transgenic tobacco embryos

A chimeric *G564*-promoter/*GUS* gene was introduced (see Materials and Methods) into tobacco and accumulation of *GUS* mRNA and *GUS* enzyme activity in transgenic tobacco embryos was monitored to study *G564* transcription regulation. The *G564/GUS* gene was active in the two suspensor cells of the five-cell pre-globular embryo. In the embryo proper, by contrast, no *GUS* activity was detected. No *GUS* hybridization grains were detected above background level, indicating that – in the suspensor – *GUS* mRNA had accumulated below the detection level of the in situ hybridization. At globular stage, both *GUS* activity and *GUS* mRNA accumulation were detectable in the suspensor and in the hypophyseal region of the embryo proper. At heart and torpedo stages, *GUS* activity and mRNA accumulation were detectable in the axis. *GUS* transcripts were also detected in the endosperm. Together, these data show that in transgenic tobacco embryos, *G564/GUS* expression and *GUS* mRNA accumulation follow the same developmental pattern as was observed for *G564* transcripts in transgenic tobacco embryos carrying the entire *G564* gene and as observed in Scarlet Runner Bean embryos. In addition, these results indicate that the *G564* mRNA basal-region-specific and suspensor-specific accumulation is controlled at the transcriptional level by the 4.2 kb 5' upstream region of the *G564* gene, and that the transcription-regulatory function of this region was conserved between plant species.

To further analyze the *G564* promoter, a series of 5' deletions were constructed and tested for suspensor-specific activity (Figure 6). Promoters with deletions of nucleotides –4242 to –921 retained suspensor-specific *GUS* activity, while promoters with deletions up to nucleotide –662 did not have *GUS* activity in suspensor cells. These results indicate that a suspensor-specific control element is present between positions –921 and –662.

Sequence analysis of the Scarlet Runner Bean *G564* promoter region revealed four sequences of approximately 100 base pairs long within the promoter region. Each repeat

is highly homologous to the other repeats. These repeats can be found between positions -1327 to -1225, -1206 to -1103, -1030 to -928, and -908 to -800. Each homologous repeat contains either the sequence GAAAAGCGAA (SEQ ID NO:10) or the related sequence GAAAAGTGAA (SEQ ID NO:11).

Additional promoter fragments from the Scarlet Runner Bean G564 promoter were isolated and linked to a minimal 35S promoter operably linked to the GUS gene. As indicated in Figure 7, two fragments encompassing the region between -921 and -662 resulted in GUS activity in the suspensor cell. These fragments were from positions -1524 through -99 and -2064 through -99. In addition, a 187 base pair fragment (positions -913 through -713 of Figure 1) linked to the minimal 35S promoter lead to GUS expression in the suspensor cell. This result suggests that at least one suspensor-specific control element is located within the 187 base pair fragment.

A comparison of the Scarlet Runner Bean G564 promoter (SEQ ID NO:1) and the Scarlet Runner Bean C541 promoter identified a conserved 10 base pair sequence which may confer suspensor-specific activity. Supporting this assertion, the sequence, GAAAAGCGAA (SEQ ID NO:10), is found at positions -846 to -837, i.e., within the area which the deletion results indicate controls suspensor-specific activity. Identical motifs can also be found at positions -1144 through -1135 and between -713 through -704 of Figure 1. The motif is also found at positions -684 through -675 of the Scarlet Runner Bean C541 promoter region (Figure 4). Interestingly, the Arabidopsis G564 ortholog promoter region comprises a motif (GAAAAGCCAA - SEQ ID NO:11) that is highly homologous to SEQ ID NO:10.

As a further analysis, a series of embryo-specific promoters that do not initiate transcription in the suspensor cell were screened for SEQ ID NO:10. None of the promoters screened (*Kti1* (Accession No. 45035), *Kti2* (Accession No. S45035), *Kti3* (Accession No. K00821) or the lectin promoter (Accession No. S45092)) contained SEQ ID NO:10.

A listing of other motifs identified in the region defined by -921 to -662 of the Scarlet Runner Bean G564 promoter region is provided as Figure 8.

DISCUSSION

The Scarlet Runner Bean embryo was used as a model system to investigate gene expression programs during early embryogenesis. Two suspensor-specific mRNAs designated as G564 and C541 were identified. In four-cell embryos, G564 and C541 mRNAs accumulate exclusively in the two basal cells, but are not detectable in the two apical cells. A

chimeric *G564/GUS* reporter gene is transcribed specifically in two basal cells of transgenic tobacco embryos at a similar stage (five-cell). From these results it is concluded that as early as the four-cell embryo stage the apical and basal cells transcribe different gene sets and are specified at the molecular level.

5 The Scarlet Runner Bean suspensor is a novel system to study the mechanisms regulating specification of the basal region of the early plant embryo

Scarlet Runner Bean has been used historically to study the role of the suspensor in embryo development. The suspensor size facilitated its micro-dissection (Fig. 10-Q) and rendered it accessible for physiological and cytological studies (Nagl, W., Z. *Pflanzenphysiol.*, 73:1-44 (1974); Sussex, I., *et al.*, *Caryologia*, 25:261-272 (1973); Yeung, E. C., *et al.*, *Protoplasma*, 94:19-40 (1978); Yeung, E. C., *et al.*, *Plant Cell*, 5:1371-1381 (1993); Yeung, E. C., *et al.*, *Zeitschrift für Pflanzenphysiologie*, 91:423-433 (1979)). Because the suspensor is simple, terminally differentiated, and only few cell generations removed from the basal cell, we have adopted this model to study the mechanisms specifying basal-cell fate. Scarlet Runner Bean suspensors were collected separately from embryo propers and used the suspensors to identify two genes, *G564* and *C541*, that are transcribed specifically in the suspensor and in the basal region of the embryo shortly after division of the zygote. The *G564* promoter maintains transcriptional activity in suspensors of tobacco embryos. Therefore, this promoter can be used to identify regulatory genes and thus as an entry point to penetrate the regulatory circuits that control basal cell specification. In addition, *Arabidopsis* genes corresponding to *G564* and *C541* were identified (SEQ ID NO:4 and SEQ ID NO:8, respectively). We can use these genes to find mutants important for suspensor function in embryo development. Thus, the *Arabidopsis* model system is complemented by the Scarlet Runner Bean suspensor as a model to investigate the earliest events in plant embryogenesis.

25 A mosaic of gene expression programs is active during seed development

In flowering plants, fusion of the sperm cells with both the egg cell and central cell initiates embryo and endosperm development, respectively (Table 1). In addition, fertilization causes the integument and the endothelium to differentiate and to contribute to the development of the seed (Table 1 and EMBRYOLOGY OF ANGIOSPERMS (Johri, B. M., ed. 1984); Miller, S. S., *et al.*, *Annals of Botany London*, 84:297-304 (1999); EMBRYOGENESIS IN ANGIOSPERMS: A DEVELOPMENTAL AND EXPERIMENTAL STUDY (Raghavan, V., ed. 1986)). Simultaneously, a cascade of different gene expression programs is initiated that are correlated with the various events occurring during embryo and seed development (Goldberg,

R. B., *et al.*, *Cell*, 56:149-60 (1989); Goldberg, R. B., *et al.*, *Science*, 266:605-614 (1994)). For example, SRB8 mRNA accumulates in the ovule chalazal endothelium and after fertilization, it accumulates in endosperm and embryo proper. SRB8 is homologous to a ribosomal protein L10A indicating a greater need for ribosome and protein synthesis in these tissues before and during early seed development SRB13 transcripts accumulate in the integuments and, after fertilization, in the seed coat and to a lesser extent in the developing embryo. SRB13 is homologous to a Bowman-Birk trypsin inhibitor illustrating the protective function of integuments and seed coat.

G563 mRNA starts to accumulate specifically at 3 DAP in the seed micropylar endothelium surrounding the developing embryo. The micropylar-endothelium cell layer is suggested to function as an embryo-nursing tissue by exchanging metabolites with the suspensor via extensive cell wall ingrowths that appear at 3 DAP (Natesh, S., *et al.*, *Embryology of angiosperms*, (ed. B. M. Johri) pp. 377-444, Berlin: Springer Verlag (1984); Yeung, E. C., *et al.*, *Protoplasma*, 94:19-40 (1978); Yeung, E. C., *et al.*, *Can. J. Bot.*, 57:120-136 (1979)). Probably because of this tight contact between endothelium and suspensor, some residual endothelial cells were present in our hand-dissected suspensor preparations, which explains why we were able to identify G563 as a micropylar-endothelium-specific transcript. The correlation of G563 transcript accumulation with the appearance of cell wall ingrowths contiguous to the suspensor of the developing embryo suggests that G563 marks the specification of the micropylar endothelium as an embryo-nursing tissue. Although the function of the predicted G563 protein is unknown, its high glycine and proline content (47.5 and 12.5 percent, respectively) suggests a structural function perhaps in the formation of the specialized cell wall ingrowths.

G564 and C541 transcripts accumulate specifically in the suspensor. G564 transcripts are distributed evenly over the whole suspensor, while C541 transcripts accumulate to a higher concentration in the suspensor-basal region than in the suspensor-neck region. Based on physiological and cytological studies, the main activities of the suspensor are importing, producing and transporting nutrients and growth regulators to the developing embryo proper (Alpi, A., *et al.*, *Planta*, 147:225-228 (1979); Brady, T., *Cell Differentiation*, 2:65-75 (1973); Ceccarelli, N., *et al.*, *Zeitschrift für Pflanzenphysiologie*, 102:37-44 (1981); Clutter, M., *et al.*, *Journal of Cell Biology*, 63:1097-1102 (1974); Schnepf, E., *et al.*, *Protoplasma*, 69:133-143 (1970); Sussex, I., *et al.*, *Caryologia*, 25:261-272 (1973); Yeung, E. C., *et al.*, *Can. J. Bot.*, 57:120-136 (1979); Yeung, E. C., *et al.*, *Plant Cell*, 5:1371-1381 (1993)). The exact functions of G564 and C541 in these activities are unknown, but the fact

that G564 protein is predicted to be secreted suggests that it might play a role in metabolite exchange in the intercellular space of the cell wall ingrowths. C541 is predicted to be targeted to the vacuole, which explains the higher concentration of C541 mRNA in the highly vacuolate suspensor-basal region.

Together, the different SRB8, SRB13, G563, G564, and C541 mRNA accumulation patterns illustrate that an array of different gene regulatory programs is active to make a seed. However, how these programs are regulated coordinately remains to be established.

Differentiation of early-embryo apical and basal regions is marked by the accumulation of different transcript sets

The suspensor is derived from the basal cell of the two-cell embryo, however it is not known what mechanisms direct the basal cell to become specified and develop into a suspensor, nor is it known when these mechanisms are active. To gain entry into the mechanisms regulating suspensor development and thus into the mechanisms regulating apical-basal cell specification events, two suspensor-specific transcripts were identified, designated as G564 and C541. The G564 and C541 transcripts first accumulate in the two basal cells of the four-cell embryo, before the suspensor is morphologically distinguishable and thus marking the embryo-basal region for suspensor specification. By contrast, in *Arabidopsis* pro-embryos a homeobox mRNA, designated as ATML1, has been found to accumulate selectively in the apical cell (Lu *et al.*, *Plant Cell* 8(12):2155-68 (1996).

Together, this shows that at the four-cell embryo stage the apical and basal regions have differentiated and that this specification process is marked by accumulation of different transcript sets. In addition, it indicates that the mechanisms activating the apical and basal-region-specification processes are active earlier either in the two-cell embryo or in the zygote or egg.

Apical and basal-region specific accumulation of mRNA is caused by specific transcriptional programs

G564 mRNA accumulation pattern in the basal-region and the suspensor is similar to that in Scarlet Runner Bean embryos. This shows that the 6.99 kb *G564* genomic clone is a marker for the specification mechanism of the basal region of the four-cell embryo and that within this 6.99 kb genomic fragment an element is present that is recognized by this mechanism. In addition, we conclude that although early-embryo cell division patterns are different between Scarlet Runner Bean and tobacco (Kaplan, D. R., *et al.*, *Plant Cell*,

9:1903-1919 (1997); Natesh, S., *et al.*, EMBRYOLOGY OF ANGIOSPERMS, (B. M. Johri, ed. 1984) 377-444), the mechanisms specifying cell fate are conserved (Goldberg, R. B., *et al.*, *Science*, 266:605-614 (1994)).

In transgenic tobacco embryos containing the chimeric *G564/GUS* gene, GUS enzyme activity in a basal-region-specific and suspensor-specific pattern are similar to the *G564* mRNA accumulation pattern in Scarlet Runner Bean embryos and *G564* transgenic tobacco embryos. This shows that the mechanism regulating basal-region specific *G564* mRNA accumulation works at the transcriptional level. Therefore, the differentiation of the basal and the apical regions of the early embryo, which is marked by differential accumulation of transcript sets, is caused by specific apical and basal-region transcription programs. Initial analysis was performed of the basal-region transcription program by dissecting the GYM promoter for cis-regulatory elements to identify its regulatory factors. Preliminary data indicate that the elements directing basal-region-specific transcription are present at -921 to -662.

A model for the mechanism of specification for the apical and basal cell of the two-cell embryo

How is the *G564* transcriptional program activated specifically in the embryo basal region and how does this provide clues to the general mechanism specifying basal-cell fate? A possible explanation might reside in the apical-basal polarized cyto-architecture of the egg cell and zygote (Fig. 1E and Willemse, M. T. M., *et al.*, EMBRYOGENY OF ANGIOSPERMS, (B. M. Johri, ed. 1984) 159-196). The asymmetric distribution of cytoplasm, and/or its contents within the egg and/or zygote may play a role in activating specific apical and basal-region transcription programs (Goldberg, R. B., *et al.*, *Science*, 266:605-614 (1994)). Based on this suggestion, a simple model is proposed for the specification of basal cells leading to suspensor differentiation. This model assumes that there is an asymmetric distribution of "morphogenetic factors" (e.g. transcription factors) within either the egg cell or the zygote or both. In addition, it assumes that the basal cell (and suspensor) is specified autonomously as a consequence of inheriting the 'morphogenetic factors' following zygotic division. These factors trigger a cascade of events leading to the transcription of basal-region-specific genes, like *G564*, and suspensor differentiation (Fig. 8).

The model outlined above is consistent with analogous autonomous specification processes that occur for specific cell types during embryo development in various animal systems (Davidson, E. H., *et al.*, *Development*, 125:3269-3290 (1998)). In

plants, this model predicts that the embryo-basal-region-specific transcription of *G564* (Fig. 5B, 7B, J) is programmed by one or more basal-cell-specific transcription factors, and that these transcription factors are derived initially from the basal region of the egg cell or zygote. It is possible that these regulatory factors are bound by the cytoskeleton to the basal pole of the egg and/or the zygote and that these factors automatically become part of the basal cell after zygote division. This would be similar to the mechanism responsible for targeting factors to unique intracellular cytoplasmic locations in animal embryos (Lall, S., *et al.*, *Cell*, 98:171-180 (1999); Yisreali, J. K., *et al.*, *Development*, 108:289-298 (1990)) and to the mechanism by which the polarized axis is fixed in *Fucus* eggs (Kropf, D. L., *Plant Cell*, 9:1011-1020 (1997); Quatrano, R., *Cold Spring Harbor Symposia on Quantitative Biology*, 57:65-70 (1997)).

Alternatively, it is also possible that a signalling mechanism is responsible for basal cell specification similar to that which establishes dorsal/ventral polarity in *Drosophila* embryos (Davidson, E. H., *et al.*, *Development*, 125:3269-3290 (1998); Sen, J., *et al.*, *Cell*, 95:471-481 (1998)). In this case, a signal derived from the maternal seed tissues contiguous with the basal cell (e.g. endothelium) would interact with a basal cell ligand which would then trigger a signal transduction cascade leading to transcription of basal-region-specific genes like *G564* and suspensor differentiation. One prediction of this model is that the transcription factors which activate *G564* transcription should be present in both the apical and basal cells of the embryo, but remain inactive within the apical cell (Davidson, E. H., *et al.*, *Development*, 125:3269-3290 (1998)).

Table 1. Description of Scarlet Runner Bean seed development stages.

Stage	DAPs after Pollination (DAP)	Suspensor length	Seed length	Seed color
Ovule	0	-	<0.75 mm	white
Proembryo	1 to 4	<50 μ m to 250 μ m	0.75 to 1.5 mm	pale green
Globular	5 to 9	320 μ m to 600 μ m	2 to 4 mm	green
Heart	10 to 12	700 μ m to 900 μ m	4.5 to 6 mm	green with red pigment contiguous to the hilum
Early cotyledon	13 to 17	~1000 μ m	7 to 9 mm	green with heavy red pigment in the area surrounding the hilum
Late cotyledon	~25	ND	~15 mm	scarlet red
Mature	~30 to 35	ND	~20 mm	purple

ND, not determined

It is understood that the example and embodiments described herein are for illustrative purposes only and that various modifications or changes in light thereof will be suggested to persons skilled in the art and are to be included within the spirit and purview of this application and scope of the appended claims. All publications, patents, and patent applications cited herein are hereby incorporated by reference for all purposes.

SEQUENCE LISTING

SEQ ID NO:1 Scarlet Runner Bean G654 promoter

	-4242	GCATGCACTG	CCACAAGTAG	TGAACCTCATG	GTTTACCTC	CTCAAGTAGA
	-4192	AAACCTTTTG	AGTGAATTG	AAGATTTATT	CTCCCAAGAA	GGACCCATTG
5	-4142	GGCTTCCTCC	TCTTAGGGGG	ATAGAACATC	AAATTGACTT	TATACCGGGG
	-4092	GCAAGCCTAC	CAAAATAGGCC	TCCTTATAGA	ACCAACCCCG	AGGAACAACAA
	-4042	GGAGATAGAA	TCACAAGTTC	AAGACTTGTT	GGAGAAGGGT	TGGGTTCAAA
	-3992	AGAGCCTAAG	CCCTTGCTCT	GTACCTGTCT	TGTTGGTGCC	AAAAAAGAT
	-3942	GGAAATGGC	GTATGTGTTG	TGATTTGTAGA	GCAATCAACA	ACATCACCAT
10	-3892	CAAGTATAGG	CATCCAATCC	CAAGGCTTGA	CGATATGCTT	GATGAATTGC
	-3842	ATGGGTCAAC	TCTATTCTCC	AAAATTGACC	TTAAAAGTGG	ATATCACCAG
	-3792	ATTCGAATCA	AGGAGGGTGA	TGAGTGGAAA	ACCGCTTTTA	AGACCAAAAT
	-3742	TGGATTATAT	GAGTGGTTGG	TGATGCCCTT	TGCTCTTACT	AACGCTCCAA
	-3692	GTACATTCAT	GAGGCTTATG	AATCACACCT	TGAGGGATTG	TATAGGTAAA
15	-3642	TATGTAGTAG	TTTATTTTGA	TGATATCTTA	GTATATAGTA	AAACCCCTAGA
	-3592	AGACCATCTA	AGTCACCTTA	GGGAAGTTCT	TCTAGTCTTT	AGGAAAAATA
	-3542	GTCTTTTTCG	CAATAGGGAT	AAGTGTACCT	TTTGTGTAGA	TAGCGTAGTC
	-3492	TTTTTAGGCT	TTATAGTAAA	CCAAAAGGGG	GTGCATGTAG	ATCCCGAGAA
	-3442	AATCAAAGCC	ATCCGCGAGT	GGCCAACTCC	ACAAAATGTA	AGTGTGTGTA
20	-3392	GAAGTTTTCA	TGGGTTAGCT	AGCTTCTATA	GAAGGTTTGT	TCCCAATTTT
	-3342	TCTAGCCTAG	TCTCTCCCTT	GAATGAACCT	GTAAAAAAG	ATGTTGCATT
	-3292	TTGTTGGAAT	GAAAAGCATG	AGCAAGCCTT	TCAAAGGCTA	AAAGCTCACT
	-3242	GACCAATGCA	CCCATCCTAT	CTCTTCCAAA	TTTTTCCAAA	CTTTTGGAGA
	-3192	TAGAGTGTGA	TGCATCGGGA	GTAGGCATAG	TGCGGTTTGT	TTGCAAGGTG
25	-3142	GACACCCCTT	GCTTATTTTA	GTGAAAAACT	CCATGGTGCC	ACCCCTACTA
	-3092	CCCCCCTAT	GACAAAGACT	CTATGCTCTT	GTGCGACCTT	AAAGACTTGG
	-3042	GGAAACACTAC	CTTGTGTCCT	AAAGAAATTTG	GnTATCCATA	GTGATCACGA
	-2992	GTCTTTAAAA	TATTTAAAGG	GCCAAACAAA	GCTCAATAAG	AGACATGCTA
	-2942	AATGGATGGA	ATTTCTTGAA	CAATTTCTCT	ATGTCATCAA	ATACAAGAAA
30	-2892	GGGAGCAGCA	ATATAGTGGC	CGATGCTCTT	TCTAGACGGC	ACACTCTCTT
	-2842	TTCAAAACTA	GGTGCCCAAA	TTCTTGGATT	TGACCCATA	AGAGAGCTTT
	-2792	ATCAAGAAGA	TCAAGAATCT	TCATCCATCT	ATGCCCAATG	TCTACATAGA
	-2742	GACCAAGGAG	GTTACTATGT	GTCCGAGGGA	TATCTTTTAA	AAGAAGGAAA
	-2692	ACTTTGCATT	CCCCAAGGAA	CACATAGAAA	ACTCCTTGTC	AAAGAACTAT
35	-2642	ATGAAGGGGG	ACTCATGGGC	CATTTTGGAG	TTGATAAAAC	TCTAGACTTT
	-2592	TAAAGGCAAA	ATTTTGTGTT	CCACACATGA	GGAAAGATGT	CCACGACATT
	-2542	GCTCTAGAGTA	TCTCATGTTT	AAAAGCAAA	TCTAGAACAA	TGCGGCTGGA
	-2492	CTCTACACCC	CTTTGCCGAT	TGCAAAAGCTC	CTTTGTGAAGA	CATTAGCATG
	-2442	GATTTCAATT	TAGGACTTCC	TAGGACTGCA	AGAGGCCATG	ACTCTATCTT
40	-2392	TGTGGTAGTG	GACCGTTTTA	GCAAAATGTC	TCACCTTTAT	CCATGCCACA
	-2342	AAGTAGATGA	TGCTCAAAAT	ATTTCTAAAC	TCTCTTTTAG	AGGAAGTGGT
	-2292	AGACTCCATG	GTCTCCCTAG	AAGTATAGTG	TCCGATAGAG	ATCACCTTAA
	-2242	ATATATAATT	ATACACTTGT	TTTTTTCTCT	TTTTTTATTT	TATCAAGTAA
	-2192	AAAGATTATTG	TTCTAGATTA	TTATGAGTAT	ATACCTTACT	TCTGTATTTT
45	-2142	ATTTCTTTCT	ATTTTTTATG	ACGATGAAAT	TTCTTATTAT	ATCCGAGCTT
	-2092	TTCAATATATA	TTTTTATTTT	TTTTCCATCT	AGATGCTCTG	TACTTTTCTT
	-2042	CAGTTGAAAT	TTCCACTCTC	CAACAAAAAC	TCTATTCAAGT	TTTTGTATAAC
	-1992	ACTGTGACGT	TAAACGAGTTA	AAATAAGAAA	ATCATGTAAT	ATAAATTATT
	-1942	TCTAGTAGATA	TTTTAGAATT	ACAAATACGA	TAAATATTTA	AAATTTAAAA
50	-1892	ATTATTAAAC	AATGAATTTT	TTTGGAAATT	AATATAAAAC	TTAGACTTGT
	-1842	GGTTTCTTCA	TTCACTCAAA	ACCTTTTTTCT	ATTGTGTGGC	GTGTGCGTGA
	-1792	ACATCGAATT	TGGGTGCTTT	ATGCGGCTTT	ATCTTCATCT	GCACTTCCAA
	-1742	ATTAATAAAT	TAATTCGGGA	AAATAATAAA	CCCACACACT	GTTTTATGCA
	-1692	TATATTAAAG	TAAATAAAGG	AGAACTATTT	TAAAGAATAT	AAAAATAATA
55	-1642	ATGTAACAAA	TGATGTCAC	AAAGAAGAAA	AAAAATTAACA	AGAAATTTGAA
	-1592	TATATTTCTT	TATGAAATGT	TTTGTGCACT	ACCGAGAGAG	GTGCAACATG
	-1542	ATACACGCAA	GCACTCTAAT	AGTTTGGTAA	TTCTTTTCA	ACATCGNTAA
	-1492	GACACATACA	CTAAAATTAC	TTTAAATAGA	TAAATTAGAT	TCAATTGGAT
	-1442	GACATTAAAT	TATAATACTC	TATCCAAAAT	TATAACTATA	AATAAAAAAT

5
10
15
20
25
30

```

-1392 TATTTTGTAGA AAATAAGTAA TGAAAAATTTA ATTCTAAAAAT TTATAACACT
-1342 TTTATGCTGT GTTTGTTTCG AAGCATAGAA AAAATAAAAG TTATTGTTGG
-1292 GAATGAAAAAG TGAAGAAAAAT CATGTAATAA AAACAAAATG ACACGACAAT
-1242 CAAAAAATAA GTTTTCATGC AAAAAGTTTT TCATAAATTA CACTTTTATG
-1192 ATGTGTTTGT TTCGAAGTGT AGAAAAACGA AAGTTTATTA TTGGTAATGA
-1142 AAAGCGAAGA AAATCACGTA ATAAAAACAA AGCAAGATGG CACGACAATC
-1092 AAAAAAAGT TTCTACACAA AACTTTATTC AAAATTTTACA ACATCTTTTAT
-1042 GTTGTGTTT GTTTCGAGG TATAGAAAAA CAAAGAATTA GTGTGGTAA
-992 TGAAAAAGTGA AGAAAACCAT GTAATGAAAA CAAAATGGCA CGACAATCAA
-942 AAAAAGTTTT CAGCAAAAAT TTTCTTCAAA ATTTATAACA TTTTCATGTT
-892 GTGTTTGTIT CAAAGCCTAG AAAAAAGGAG AGTTACTATT GGTAATGAAA
-842 AGCGAAGAAA ACCACATAAT AAAAAACAAA TGGCAGCACA ATCAAGAAAAA
-792 AGTTTTCACA CAAAAGTTTT TTTCAAAATTT ACTATGTTTA TTTGCAAAAT
-742 TAGAAAAACG AAGAGTTTAT ATTAGTAATG AAAAGCGAAG AAAACTACGT
-692 AATAAAAAAC AAAATGGCAC GACATAAAAA AAAGTTTCA CGCAAAATTT
-642 TCTTGGTGCG CAGAAAGTTA TATATATTA TTAATTAATT TCCATTACT
-592 TTTTCCCTTT TTTATTTTAA AGTTAAATTA TTATTATTTT CATTTAAAT
-542 ATAAATATTA TTTAAATATA AAAAATATA CCTTAATCAA AACAAAGCCT
-492 TAATCTAAAA TTTACACAC TTTTAACCTT AAAATTAAC TTAAAAGGAA
-442 AATGATAGTG TGACAACTAA AAAAGTTGTA TACAACCTGT TCAATAGTTT
-392 AGAAATAAAT ATATATAATA AAGAGTAAAT TTGTAATTA ATGATATAAA
-342 AAAGTATTA AAATAATAATA TTTAGAGTAG TAATATGTTT GTATAAAAAA
-292 ATGTGTTTGT CCATATATCA TTATTCACIT TAAATAATCA TGACAATAT
-242 TTTCCCGAA AGATGGAAAG AACGAAAGA GCGTTGGATA ATGGAAAAAT
-192 ACAAGCAATC TCCCTCCAGT ACTTTGCGATA ACATTTTGA TTAGTAGATGA
-142 GTTTTTTATC ATATATATTT AGAATATAGG AAAATTTTAG AATCAGGTGG
-92 ATAGCTATAT AATAGTAATA TTTTAATTTA TAATGTAGTT GATTTTATTT
-42 GTCACCTGGT ATACATAAAT ATGTGTTGAT AGTGGGTGAC TTGTGGCTTA
9 AAGAAATGTC CAGAGGCTGA CAACAACCTC GCACAGACTA GCGTAAAC

```

SEQ ID NO:2 Scarlet Runner Bean G654 genomic region

35
40
45
50
55

```

-4242 GCATGCACTG CCACAAGTAG TGAACATCAT GTTTTACCTC CTCAAGTAGA
-4192 AAACCTTTTT AGTGAATTTG AAGATTTATT CTCCCAAGAA GGACCCATTG
-4142 GGCCTCCCTC TCTTAGGGGG ATAGAACATC AAATTGACTT TATACCGGGG
-4092 GCAAGCCTAC CAAATAGGCC TCCCTATAGA ACCAACCCCG AGGAAACAAA
-4042 GGAGATAGAA TCACAAGTTC AAGACTTGTT GGAGAAGGGT TGGGTTCAAA
-3992 AGAGCCTAAG CCCTTGTGCT GTACCTGTCT TGTGTGTGCC AAAAAAGAT
-3942 GGAATATGGC GTATGTGTTG TGATTTGAGA CCAATCAACA ACATCAACAT
-3892 CAAGTATAGG CATCCAATCC CAAGGCTTGA CGATATGCTT GATGAATTGC
-3842 ATGGGTCAAC TCTATTCTCC AAAATTGACC TTAAGAGTGG ATATCACCAA
-3792 ATTCGAATCA AGGAGGGTGA TGAGTGGAAT ACCGCTTTTA AGACCAAAA
-3742 TGGATTATAT GAGTGGTTGG TGATGCCCTT TGGTCTTACT AACGCTCCAA
-3692 GTACATTCAT GAGGCTTATG AATCACACCT TGAGGGATAG TATAGGTAAA
-3642 TATGTAGTAG TTTATTTTGA TGATATCTTA GTATATAGTA AAACCTTAGA
-3592 AGACCATCTA AGTCACCTTA GGGAGTTTCT TCTAGTTCTT TGAAGAAATTA
-3542 GTCTTTTTCG CAATAGGGAT AAGTGTACCT TTTGTGTAGA TAGCGTAGTC
-3492 TTTTTAGGCT TTATAGTAAA CCAAAAGGGG GTGCATGTAG ATCCGAGAA
-3442 AATCAAGGCC ATCCGCGAGT GGCCCACTCC ACAAATGTGA AGTATGTGA
-3392 GAAGTTTICA TGGGTTAGCT AGCTTCTATA GAAGGTTTGT TCCCAATTTT
-3342 TCTAGCCTAG CTCTCCCTTT GAATGAACCT GTAAAAAAG ATGTTGCATT
-3292 TTGTTGGAAT GAAAAAGCAT AGCAAGCCTT TCAAGGGCTA AAAGCTCACT
-3242 CACCAATGCA CCCATCCTAT CTCTTCCAAA TTTTCCAAA TTTTGGAGA
-3192 TAGAGTGTGA TGCAATCGGA GTAGGCATAG TCGGGTTTTG TTCAAGGTG
-3142 GACACCCCTT GCTTATTTTA GTGAAAAACT CCATGGTGCC ACCCTCACTA
-3092 CCCCACCTAT GACAAGACT CTATGCTCTT GTGCAACCTT AAAGACTTGG
-3042 GGAACACTAC CTTGnGTCCC AAAGAATTTG GnTATCCTTA GTGATCAGA
-2992 GTCTTTAAAA TATTTAAAGG GCCAACACAA GCTCAATAAG AGACATGTGA
-2942 AATGGATGGA ATTTCTTGA CAATTTCCCT ATGTCACTAA ATACAAGAAA
-2892 GGGAGCACCA ATATAGTGGC CGATGCTCTT TCTAGACGGC ACACCTCTCT

```

-2842 TTCAAACCTA GGTGCCCAAA TTCTTGGATT TGACCACATA AGAGAGCTTT
 -2792 ATCAAGAAGA TCAAGAACTC TCATCCATCT ATGCCCAATG TCTACATAGA
 -2742 GCACAAGGAG GTTACTATGT GTCCGAGGGA TATCTTTTAA AAGAAGGAAA
 5 -2692 ACTTTGCATT CCCCAGGAA CACATAGAAA ACTCCTTGTC AAAGAATCAC
 -2642 ATGAAGGGGG ACTCATGGGC CATTTGGAG TTGATAAAAC TCTAGACTTT
 -2592 TAAAGCAAA ATTTTGTGG CCACACATGA GGAAGATGT CCACGACATT
 -2542 GTCTAGAGTA TCTCATGTTT AAAAGCAAG TCTAGAACAA TGCCGCTGGA
 -2492 CTCTACACCC CTTTGCCGAT TGCAAAGCTC CTTGTGAAGA CATTAGCATG
 -2442 GATTTCAATT TAGGACTTCC TAGGACTGCA AGAGGCCATG ACTCTATCTT
 10 -2392 TGTGGTAGTG GACCGTTTTA GCAAAATGTC TCACTTTATT CACTGCCACA
 -2342 AAGTAGATGA TGCTCAAAT ATTTCTAAAC TCTCTTTTAG AGAAGTGGTG
 -2292 AGACTCCCATG GTCTCCCTAG AAGTATAGTG TCCGATAGAG ATCACTCTAA
 -2242 ATATATAATT ATACACTTGT TTTTCTTCTC TTTTCTATT TATCAAGTAA
 -2192 AAAGTATTGT TTCTAGATTA TTATGAGTAT ATACTTACT TCTGTATTC
 15 -2142 ATTTCTTTCT ATTTTATGAT ACGATGAAAT TTCTTATTAT TCTCAGACTT
 -2092 TTCAATATA TTTTATTTTC TTTTCCATCT AGATGCTCTG TACTTTCTTT
 -2042 CAGTTGAAAT TTCCACTCTC CAACAAAACA TCATTCAAGT TTTGTATAAC
 -1992 ACTGTGACGT TAACCAGTTA AAATAAGAAA ATCATGTAAT ATAAATTTAT
 -1942 TCAGTAGATA TTTTAGAATT ACAAAATACGA TAAATAAATA ATATTAAGAA
 -1892 ATTATTAAC AATGAATTTT TTTGGAAATT AATATAAAC TTAGACTTGT
 -1842 GGTTCCTTCA TTCAGTCAAA ACCTTTTCTT ATTTGTGGGC GTGTGCGTGA
 -1792 ACATCGAATT TGGGTGCTTT ATGCCGCTTT ATCTTCATCT GCACCTTCAA
 -1742 ATTAATAATT TAATTCGGGA AAATAATAAA CCCACACACT GTTTTATGCA
 25 -1692 TATATTAAGA TAAATAAAG AGAACTATT TAAAGATAT AAATAATAAA
 -1642 ATGTAACAAA TGATGCTACT AAAGAAGAAA AAAATTAACA AGAATTGTAA
 -1592 TATATTCTTT TATGAATGT TTTGTGCATT ACCGAGAGAG GTGCAACATG
 -1542 ATACACGCAA GCATCTAAT AGTTTGGTAA TTTCTTTTCA ACATCnTAA
 -1492 GCACATCACA CTAAATTTAC TTTAAATAGA TAAATTAGAT TCAATGGAT
 -1442 GACATTAATT TATAATCTC TATCCAAAAT TATAACTATA AATAAAAGT
 -1392 TATTTTLAGA AAATAAGTAA TGAAAATTTA ATCTCAAAAT TTATAACAT
 -1342 TTTATGCTGT GTTTGTTTCG AAGCATAGAA AAATAAAGAG TTTATGTGG
 -1292 GAATGAAAG TGAAGAAAT CATGTAATAA AACACAAATG ACACGCAAT
 -1242 CAAAAAATA GTTTTCATGC AAAACTTTTT TCAAAATTTA CACTTTTATG
 -1192 ATGTGTTTGT TTGGAAGTGT AGAAAAACGA AAGTTATTTA TTGTAATGA
 35 -1142 AAAGCGAAGA AAATCACGTA ATAAAAACAA AGCAAGATGG CACGCAATC
 -1092 AAAAAAAGT TTCTACAAA AACITTTATC AAAATTTACA ACACITTTAT
 -1042 GTTGTGTGTT GTTCCGAGG TATAGAAAA CAAAGAATTA GTGTTGGTAA
 -992 TGAAAAGTGA AGAAAAACAT GTAATGAAAA CAAAATGGCA CGACAATCAA
 40 -942 AAAAAGTTTT CACGCAAAAT TTTCTTCAA ATTATAACA TTTTATGTT
 -892 GTGTTGTTT CAAAGCCTAG AAAAACGAG AGTTACTATT GGTATGAAA
 -842 AGCGAAGAAA ACCACATAAT AAAACAGAAA TGGCAGCAGA ATCAAGAAAA
 -792 AGTTTTCACA CAAAACTTTT TTTCAAAATT ACTATGTTA TTTTCAAATT
 -742 TAGAAAAACG AAGAGTTATT ATTAGTATAA AAAAGCGAAG AAAACTACGT
 45 -692 AATAAAAAAC AAAATGGCAC GACAATAAAA AAGTTTTCa CGCAAAATTT
 -642 TCTTGGTGGC CAGAAAGTTA TATATATTTA TTAATTAATT TTTACTTACT
 -592 TTTTCCCTTT TTTATTTTAA AGTTAAATTA TTATTTATTT CATTTAAAT
 -542 ATAAATATTA TTTAAATATA AAAAATATAA CCTTAATCAA AACAAAGCCT
 -492 TAATCTAAAA TTTACAACAC TTTTAACTTT AAAATTAAC TTTAAAGGAA
 -442 AATGATAGTG TGCAACTTAA AAAAGTTGTA TACAACCTGT TCATAGGTTT
 50 -392 AGAAATAAAT ATATATAATA AAGAGTAAAT TTGTAATTAA TGTATATAAA
 -342 AAAGTATTAA AATAATAATA TTTAGAGTAG TAATATGTTT GTATAAAAAA
 -292 ATGTGGTTGT CCAATATATCA TTAATTCATT TAAATATACA TGACAAATAT
 -242 TTTCAACGAA AGATGGAAG AACGAAAAGA CGGTTGGATA ATGGAATAAT
 -192 ACAAGCAATC TCCCTCCAGT ACTTTGCATA ACATTTTGTG TTAGTATGTA
 55 -142 GTTTTATATC ATATATATTT AGAATATAGG AAAATTTTGA AATCACTGGT
 -92 ATGCTATAT AATAGTAATA TTTTAAATTA TAATGTAGTT GATTTTATTT
 -42 GTCAACTGGT ATACATAAAT ATGTGTTGAT AGTGGGTGAC TTGTGGCTTA
 9 AAGAAATGTC CAGAGGCTGA CAACAACCTC GCACAGACTA GCGTAAAC
 57 **ATG** AAG TCC AAT TTT GCT ATT TTC GTA GTC TTT TCT CTT CTT CTT
 60 1 M K S C N F A I F V V F S L L L
 102 CTG **GTACCTCTTCACTCTCTCTCAAAAACCTCTGTTGCTCTTTCACTCTGTTGTA**

16 L
 160 ATTTTGTTTACACTTTTGGAAAAATGAAGCTGATATATATGTAACAACCTTTCAGTTT
 219 GTCTGCACCTGAAACTGATAGAAAAATATACGTTTGTGGATATATAG GTT GGC
 17 V G
 274 AGT TGC AGC TGC GCA AGA AAA GAC ATG AGA GGG TAT TGG AAG GAT
 19 S C S C A R K D M R G Y W K D
 319 ATG ATG AAG GAG CAA CCT ATG CCA GAA GCA ATC AAA GAC CTT ATT
 34 M M K E Q P M P E A I K D L I
 GAG GAT TCA GAA GAA GTG TCA GAA GCA GGG AAG GGT CGT TTT GTT
 49 E D S E E V S E A G K G R F V
 AGG GAC TTC GAT GTA AAG CCT AAT GTC ATA TTA TAT CAC ACA CAT
 64 R D F D V K P N V I L Y H T H
 GTT GTG CCC ATG AAG CAG AGG CAG AAG AAT AAA GAT TGA
 79 V V P M K Q R Q K N K D •
 493 AGACTATGTGATTGGCAGTTTCAGACTTATTTGGCACCAAAATTTATGATGCTCTTGTGCTG
 555 TTTCAAATTTGTACTCAAACCTTTGAACCCCTTTGCAGCATCTTGTCTCTTTTGGTCTGTGT
 617 GAATTTTGTACAGTTATACCTGTACGAAATAGTTTCTCTTCATAATAAGCAACTTTTCTCTCT
 679 C

SEQ ID NO:3 Scarlet Runner Bean G654 amino acid sequence

57 **ATG** AAG TCC AAT TTT GCT ATT TTC GTA GTC TTT TCT CTT CTT CTT
 1 M K S N F A I F V V F S L L L
 102 CTG GTACCTCTTCAATCTTCTCTACAAAACCTCTGTGCTCTTTCCACCTCTGTTTGA
 16 L
 160 ATTTTGTTTACACTTTTGGAAAAATGAAGCTGATATATATGTAACAACCTTTCAGTTT
 219 GTCTGCACCTGAAACTGATAGAAAAATATACGTTTGTGGATATATAG GTT GGC
 17 V G
 274 AGT TGC AGC TGC GCA AGA AAA GAC ATG AGA GGG TAT TGG AAG GAT
 19 S C S C A R K D M R G Y W K D
 319 ATG ATG AAG GAG CAA CCT ATG CCA GAA GCA ATC AAA GAC CTT ATT
 34 M M K E Q P M P E A I K D L I
 GAG GAT TCA GAA GAA GTG TCA GAA GCA GGG AAG GGT CGT TTT GTT
 49 E D S E E V S E A G K G R F V
 AGG GAC TTC GAT GTA AAG CCT AAT GTC ATA TTA TAT CAC ACA CAT
 64 R D F D V K P N V I L Y H T H
 GTT GTG CCC ATG AAG CAG AGG CAG AAG AAT AAA GAT TGA
 79 V V P M K Q R Q K N K D •

SEQ ID NO:4 Arabidopsis G654 genomic region

101001 CAAAACAAAAGCAAATGCCGGTTTTCTTATTATTATTTGCAACTTTAGAC
 100151 CTTTTTGTAACGTTTCTTAATTTTTTTCCTTGATAAAGAACCCCTATTAT
 45 100201 ATCTTAGCTAAATATTACCTCATTGTGTTATGAGCTAAACACCCCAA
 100251 AAATATTGTAGTTTGTCTTCGGATTAACTGCCAAGCAAGTGATTAGAT
 100301 ATATTAAAGGAAAAATGAATGAAAGGACAAAAAATATAAACGACCAATATT
 100351 TGAATACTGATATTATCTCCATTCTCAATATTTTTGATTATTGTGAC
 100401 AATATTGGTTGTTTCCCATTTGCTACATCTTTGAGGACATGAAATGATA
 50 100451 ACATATATATGAACGAGTATAATACATTTCTCGTTTTCATTTTACAAATAAT
 100501 GTCAATTTATGCTAACATTTTTATTAAAAAATATCCTTTATAGATTTTC
 100551 AGTGTATTATTTTACCATGGTACTGTAAAGTCGGATGCTATATATATATA
 100601 TATATATATATATATCAAAATGACACTGAAGAATTTATTGAACTAAAA
 55 100651 CTA AAAACGTA AAAATAAAAAGAATTTTCAAAAATCAAAAATTTATATA
 100701 AAAATATAGATAAAATGTTAATATAGTACAACCTTCATTTCAACAGAGAG
 100751 AATAAATCTTCTATAGACAGTGAATATCCATTATAATAACGAGCAATAGT
 100801 TGTAATGTTGCAGTACAAAAAGAGAATTTGTAATATTGTGCATGATTGAG
 100851 AAATCAAGTTGACTTTGAAATAAAAGGCTAATCCAACAAGTACATGTA

100901 GAAGTTGACTATAGCTATATATTTACTACAAATTTGATCATTTTCAAGAAAG
 100951 ACATTTAAATTAAGATATGTCATGCACTGATTGATGAACCCCACTCGCTT
 101001 GCTTCGTGCCATTTCGACAAGATGTTACTTTTAAATGCAAGGTAAATATG
 101051 GATATACTCTCTGTAATTTTGTAGTAGATATTTTACGAAAAATTTGTTT
 101101 TTTTCCAAAAACAAATGATATTTATTAATTTTCAATAGCAATTAATTA
 101151 AATTTTAAATTAATTTTGAAGATTTATATGCTGCAGATTTAGATTACCATTTG
 101201 GTGAAATCATGTTTAGGTAAATAATAAATGATGTTGTAGTTTAGGAAAAA
 101251 AAAAAATCTTTAATCTTTATGTAAGAAATGTTAACTTCAATTTAAAAA
 101301 TATGAAGCAGTATATATAAGATGTTTAACTAATCGAATTAATTTTGTG
 101351 GGATGAAATTTTCTTGCAATGTTTCTAAAAAAATAATATGTGAAAAAT
 101401 AACATTCAATTTGATGTTTATAAGAAATATATGTGAGTTTGTGTTAGATAATG
 101451 ATAATACCTAAAAATGAAGATTTGTAAAGTTATACCTGCATCTCAAAATG
 101501 TTATTTTTCTCTTTATTTAAAAATACAGCAACATCTCAATGATTTTAT
 101551 TTTCTTTAAAAAATGAAAAAATGAAATTAGCAAAATATGTAAAAATTTAAA
 101601 ACGAATTTAAGAAAAAATTTGTAAAGATATGATATGCTTTATAAAAAAT
 101651 ACTGGTGGCGTACCTACTAAATATGATCACATTAGAGATTTGTATCCCTT
 101701 TAGCATATAGTATGTAGTATAGATATCTATATTTTATTTATTAAGAGC
 101751 ATATTATAATATAGTATATATATGTTAATTACAATAAACGTTCAATTCG
 101801 TTATGTTAGTTTGTGAAAACTTATTGGCTGTGCATATCAATGTGAGAAA
 101851 GCGACTCCACATGTGAGATGTTGGTCTGAGAAAGCTTTCTGCACCTGGTC
 101901 GGAACACTTTCATGGACTAGAATGCAATCCATCTATTCAAAGAAAAAGCAG
 101951 TTGTCATGCACTGCTCGGTTTTTCACATTTTGAAGCAGCGCAACAATGT
 102001 CTTACATAATATGCGATCGATCACTCTGCAACCAATATTCAAGTACATAG
 102051 ACCATGACATCAAAAACATTTATCACACCGAGAAGAAAGAAACGTCAAATTT
 102101 GGTAACCTTAATGGCGTTATGCTCGCGGTGAATTTCTCTAAGAGTTCTCCC
 102151 AAATTTTATGATTCTCTGTTTTTAACTTTTTCGCCAAAGAAATCATACAT
 102201 ATAGATTTGACACCACTTTCAACTTTTCAAAATACAAGTAAATTAATTT
 102251 CAAGCTTGAAAGGAATTTAAATCATGATCTAAACCTAAACGACAAATCTT
 102301 CACAAGTGAGAATCACTAATTGACTACCCCTTGGTCGATATACATCATTT
 102351 GTGTAAATCTGAAAAATTTGGTTTGGATTTTGATCTGATATGTCATCATAT
 102401 AAAACTTGTATTTATTTTATTTAGAAATTTTGGCGCAACAGATAAATCATC
 102451 ATCTATTTAGAAAAATTTTCATTGCAACCAATTAATCAGGGGAAAGAGT
 102501 GAAATCACAATCTTATCTACACTCTTTATTAATTAACCGCCATAAATATA
 102551 ACAAATTTTCAAAATACCACTTATGAGAAGCACTAAGATCACTTTTCTCT
 102601 TATGACTTTCTTTCTAAAGCTAAGCTGGTAGTCTAGCTCATGATTATCC
 102651 TTTTCTTAATGGGAATATTGTGGAAGCGGTTTCAAAATCTTTAGACAAAAA
 102701 TCCATGGCCACTAAAAGTTAGCAAAAGTTAAAAATAGTTTAAAAAAATATG
 102751 AGTGTACTTGGCCATATGCCATATTGTTGAGATCATAAACAAGAGAAATA
 102801 TAGTTTATTGAAGTTTATGATCATAAATCAACAATACATCATGCTTTCATCA
 102851 ACATTTTCCATGGATTGTAGAGGATCAACTTCAATACTAATGGTGGAGGTC
 102901 TTATTCATCCATTGCTCTCATGCCAATTAAGCAGTTAGGTTATTTGTGTGTA
 102951 CTCTAGTAGTTGCCAAATCAATCTTAATATTCACAAATGTTGTAATTTCTTA
 103001 ATTACGTATAGATAAATGACTAGATAAACAACGTTGGCTTTGTTTATCAGG
 103051 AAAGTTTCCAAATCATATATATGAATGTAGAATAGTGTCTCTCAATTAAT
 103101 TATTAATTAGCATCTCACCATTCTGAGACTGGGAGCATGTGACCAAGTTGAC
 103151 ATGTGTATTAAAGAACTTTGAGAAAAACCACTTTATGATACTCCCATCT
 103201 GAGACTGGGATGAGTACCATTTTATAAAAAATGAGTAGTGAAGAAAAATG
 103251 TCAAAAAAATTTCAACATGTCTTTTAAACATTTTAACTTTATAATTTT
 103301 AACAAACACTCTTCCAATATGCGTTATGAAGAACTTTATAAACTTTTATAT
 103351 AACATGCTTTTGAATAATTTATAAAATCTGATTTTGAAGAAAGAGTAT
 103401 ACTTTTGAATAAGACAAATGAAGTGCTATTTTAAAAATGATATCATATA
 103451 AGTCTTAACTGTGTTTGTGTTGAATTTTATTTATATACCTGTGCAAAATATC
 103501 AACTAAATAAATAAATTAATTTATTAATCATGAAGATAAATTAATATC
 103551 ATAAAAAGATAAATAAATAACCAACAAATTTATATTTGTTATAAAAAATAC
 103601 TTTGAGCTCTTCTCATAGACTTTTCCAGCTTCCATCTAGAAAAATCACA
 103651 TAAATTAAGATAAATAAACAAGAAATAAACATAGTTTCACTCTTAAGTCTT
 103701 AGTCTTAGATTTGTTTAAATTTTCAAGGTTTAGGTATGTTATATGTTTT
 103751 TTTTATGGGTTGCTAGATTTTGATCTCAAGAAGAAATGACGGGTGTAGT
 103801 ATAGATGGTTTGTGTTGAGTTTTCCTTGGTTTACTTCGTTTGGTTTT
 103851 TGTCCCAAGAAATGTTCTGTACTCGCTGGTTTATGCTCTCAAGAAAGTC
 103901 ACGACCAATGCGCGCTCTTGTATTTCAACTTGAATTTCAAAATTCGATTC

103951 ATGAAAAAAAAATGTATCTCTTAAAGTCCATTAGTACCAAAAAATACTAT
 104001 ATCATTACTACATAAAATAGTCTTGGGTTTCCAAAGTATTTCGTGTGATA
 104051 TATGTTAAGAGTTCGAAATAGACACATAGATATAATGTTGAAATGGGACC
 5 104101 TCTCACATAAATTATCTCCTTTCTCTTCATTCTCTACACTCTCAAGTTTC
 104151 CAATCCACCCCTAAGGTAATTTATTTCTTAACTAAGTAAATTTGTTAAC
 104201 AAATCTTAACTAGCTACAAATGTGTATTACAAGTCTTAAATAAAAACCTA
 104251 CTTTAATTCAAAGGTATTAAACCTTCTCTAAATTGATACCTTACTTAGTATC
 104301 GATCGGTCTAGTTTGGGTTTGGACACACACCATCATGGGACGAAATT
 104351 AGTCATTCTACGGTGTCCAAGACACAAATCTCGGACTCGATGTGGATATG
 10 104401 ACACCTTCATTATACTTTTAACTTCAATAAACTAATTTAGGAGGGAAG
 104451 AATCGGAATCTGCATATCAATCACAATAGACTATAGTATACTTAGATTAT
 104501 GATCTAATCAATGGGCTCCTTCAACTAATAAGTAGGCCCATCAACAAATTT
 104551 GAAATCATAAGACATTATAAATTAATCAATGTTCTAAAAAATCTTTGGT
 15 104601 TATGTGTCCCGTAGAGCTAATGTGCACACACAAATGAAAGTTGACCCGTTT
 104651 CACTTGTCCTCACTTTTATGATCTTTCTTTTAGTTTAAATCCAACCTTTTA
 104701 TAATCTCATCTTGTATATCAACAAAACCTTTTGGCGTCTCTTTTTCATAAT
 104751 TTAAGTAACCTCTCAGCGAGAAAAGCCAACTTTTCTCTGTTTATATTC
 104801 TTTTAAAGAAAATGAATTCAAGGGGACCCAAATTTAAAGGAAAACCA
 104851 AAACCTCCTTTCTATGTATTTTATCTTGAAGTTTCTATGTAATCAACAA
 20 104901 TCCTAACAGTAGAGAATAAAAAACATCGTTTGGGAGGTTTATATATAGC
 104951 ATATGAGAATAGTCTTAAAAATGTTTACACAAAAATTAGATTTCCTTTT
 105001 CCTCTGTCAATGGAGCTATATCACTTGTCATTTTGCTTAAACCTTTGCGG
 105051 GAAGATTGTTATGAAACAGTTTAAATGGAATCTAGTTGCCAATGTCAAG
 105101 TTTAATATGTTTGTCCCTATACCTTTATGTAATCTTATAATCTTTGTTAT
 25 105151 AGAATTATCTACTTTTAGTATTTTACATTAACATAATCTATAGAATTCT
 105201 CTTTGTCTATACAAATAAACAAAGTAATATATTCTTAATACATATTAAAA
 105251 ATGGTGGTGTGCTATCTGAGCTGTAAATGTTGATTGCTCCAGAGAAGAA
 105301 TAGACAAAAATCCTTACTTTAAGAGGCCCACTCTGAAAAATTAGACAA
 105351 GAAAAATTAACAAAATTAGGTTACACATATTATCATTATATATATATGCA
 30 105401 CAACACAAAGTTGACCTTGCAATGTACTATTGAATAAAATTAATTAATGTC
 105451 AAGAAGAGGAGGAATTATCACTGTTACCAAGAAAACAACCTCCTCTAAAC
 105501 AGGCTCTATATATATAAACTTTAAACCTAAAGAAATTAACACAGACTCAA
 105551 GAAAAATCCTCAAAACAAAAGTTAAAGCAGAC **ATG** A G CAA CAG CAA
 1 M K Q Q Q
 35 105599 CGT TAC TTG GTC GTC TTC ATC GTC CTT TTA AGC TTT CTT
 6 R Y L V V F I V L L S F L
 105638 CTG GTAAAGCTTCTCTCTTAAATATATTAAACCCCTAATTAAAGATCTCATATA
 19 L
 105691 TCGTAATGTTGTATATATTGTTGGTATAG TTT GTG AAT CTG AGT
 40 20 F V N L S
 105736 GAA GGA AGA ACA GGA GGA GTT GCA GAA GAA TAT TGG AAG
 25 E G R T G G V A E E Y W K
 105775 AAG ATG ATG AAG AAT GAA CCG TTG CCT GAA CCA ATC AAA
 38 K M M K N E P L P E P I K
 45 105814 GAG CTT CTC AAC AAT CCT TTT AGG ACC GCA CAA GAG AGA
 51 E L L N N P F R T A Q E R
 105853 TTC ATC CAG AAT TTC GAC ACC AAA TCT GTT GTC ATC ATC
 64 F I Q N F D T K S V V I I
 105892 TAC CAC AAT CCT AAT GAA TAA TCAATGAAGTCTCTCATATAG
 77 Y H N P N E •
 105934 ATATCTATGACTTTAATTTGTGTTTATGTATGGATCGACTTATACGTGCA
 105984 CGTATATGTTATTAATTAAGAAAAGAAAAGCTGCTTGAGTTGTTGTGTT
 106034 ATACACGTATACTAAATATGTTCTGTTTAGTGCAGAAATGTTAACCCCTAG
 106094 CTATAAGGGATTTTGTGTTCTTTTGTGTTTACCATTAAATGTGAGTGA
 55 106144 GTGAGTTTGTGATGAAATTAGATTGCTTCCACATTGTTGTTTGTGATA
 106194 TATATAAATCAATATACTGTGCTTCTGTGCTGTTGTTCTTATATTTT
 106244 TGTGACATTAAATTAATCTTATCAAAAAATTTATTTTAACTGTGTC
 106294 CTATGAAAAAGATGAACATATGAGTTAACCTCATCTCAAGAGAGATTCT
 106344 TTTTGTGTTTGTGTTTCT

2341 AAATTAATAATCCGAGGAAAAGACCAAGCCCTTAATAGACATGAAAAGGCTACTCGTTG
 2401 AAAAAGATGAACCCAGTTTCATCTTCACAAAAACCTGAATTTATAGGATATGATTTTAAAG
 2461 AAATATTGAGAAAAGCGAAAACATCACATAAAGAAATAACCATTTAGCGATCTTAAATAGTG
 5 2521 AAATAAAATAATTAAGGCGGAAATCGAATCTATAAAAGTCGAGCTACAAGAATTAAGG
 2581 ATAAAAATTATACATGAGGAATCCATCTCCTCTGCCGACGAAATTCACAGAAGAGGGAAG
 2641 CTAGTAGACCTTCCATCAAGAAATTAACATACAAAAGACAAAGTGGCATGTAAAAATAG
 2701 CCCTAGAAATTTGTTTGTGTTTGTGACCGTTTCATGTTGGTCAAAGATGAGTCTTACCTAA
 2761 CACAATAAAAAAGTTACTCTTAAATATCAAGGAGAGCTACAATATCAATGAATGAAT
 2821 GACATTAAATTTTTCTTTAGTTTAAAACTTGAATGAGTTGTTTTCATAAATATCTGAC
 10 2881 TGACTGACATTTTATTTTCTGAAAAATGAGGAAGTTTATACGTTAAACACCATATAT
 2941 ATATTTTATCTCAAAGTCAACGAAATATTATAAAGAATCAATTTAAAAAAATTTATCT
 3001 TTTGCAAAAAAATAATTAATAATATGAAACTCCTCCACACCATATTAACCATATTATAAA
 3061 TATAAAAAACCTCTCACAAATGTGCATCTTGAATTTCTTTATGTTGAGAGATTAATCTC
 3121 TAAAGAAAAAGGTTGAGAAAGGTGCAGCAACA ATG TCT CCA TTC TGT AGA
 15 1 M S P F C R
 3172 AAC TTT TCA ATG GCA TGG GTG CTT ATG GCA TTT GTG TTG TTT
 7 N F S M A W V L M A F V L M F
 3214 GCA AAC AGT GCT ATG CCC ACA AAT GGA TCC ACT GTT GGG GTA
 21 A N S A M P T N G S T V G V
 20 3256 AAA AAC ATG TTG GGT GGT AAA TTG ATG CTA AAC GTT TTA TGT
 35 K N M L G G K L M L N V L C
 3298 CCC CAT ATT GAT AAG CAA CAC ATT ATC CCG AAT GGT GGT TCA
 49 P H I D K Q H I P N G G G S
 3340 TTT GAG TGG AAG TAC AAT GGT GGT GCT CCA CCA ATA GGA CAA
 63 F E W K Y N G G A P F I G Q
 3382 TCA CCA TTC ATG TGT TTC TTT CGG TGG AAT AAT GTT CAT CAC
 77 S P F M C F F R W N N V H H
 3424 TCC CTT GAT CTG TGT TCA CCA AGC AAG TAT ACT GGT TGT GAA
 91 S L D L C S P S K Y T G C E
 3466 AAT GCC ATT TGG GAA ATC AAA GAA AAG CAA TTT TGT AGG TAC
 105 N A I W E I K E K Q F C R Y
 3508 AGA GGT GGA CCT ATT AAT TAT TTT TGC TAT GAC TGG GAT GAT
 119 R G G P I N Y F C Y D W D D
 3550 TAG TTATATAGATTATTCATGTTTCATCTCAATAAAAAATGACTTTAGAGTGATTCCT
 35 •
 3609 AGTTTGCCTTAACATTCTTACATATTCTTAACATTTCCGTCCTACCACCCGTAACATATAT
 3669 TTAATTTAAATAGTATCTGTGCAGTTTATTTTAAAAAAGGTTATGTGGATTAGAAG
 3729 AGAGATAAATATGTAGACGGTCACCAACCTTAATTTTGAAGCTATGTAAGACTATATTGA
 3789 CCAAGAATATATGTTTAAACTCATTCTTTAAAGACTATATCTCCATTTATGATTATGCA
 40 3849 AATGCAATTAGTTTCTTTTTCATTGAAGAATTCAAAAGAAAGTTATCATTTAAAAAGTAT
 3909 CATTAATCACTTATATGTTGTTTCTTAATATCCTTATGTTTAAATAGAAATTTTCTT
 3969 ATCCCTTTAAATGAAGTTATTACTACTTTTCTCATATCTTCATTATTTTGAATATTT
 4029 TAAATATTCAATTTTGTAAACACCCAGAAAATACATGTAACTATCACTTTTCTTTT
 4089 TATTACAAATTTATGACTTATAGAAATCAAAATATTAAGAGTTTCAAAACTACA
 4149 TCCTAAGTCTTTCAGACCCCTGACACATGTATCATCTGCTCGTATATGTGATCAGCT
 4209 ATCCGAGTTTCAAGATAACAAGAAAACCAAGGTTAAGCTAATGAAAAAATTTCCATAA
 4269 CATATTTTAATTCATGCAAAAAGAACCGTCAAGTAATCATTTATTAACATTTCTTTAA
 4329 TATTGTTATATAAAATTTCAATATCAATTTTCATCATTATATAGACCCACATGGAATCTA
 4389 TTTTCAATCACAATCATTTGGATTTCATTTTAATCCTACTTCGhCTTCCAGAAGACTCAT
 4449 AATGATGCCCTTACCAGAGACTAACACCTAATCAAGAGAAATGATCAAGGTAAGTTCAA
 4509 ACATCCAAATAACGAGTGCCCTACAGTGGGACCCAAATGTGATGAACCTTATCAGCTTT
 4569 CACCACCTGATATCTTATTCTATATGACGTAGATCATCAGTAACTAGAGGATCTCCGT
 4629 TAAACATATGTTTTTATACTTAAATGTCATCAACCAACACTCACACATTATCCAAATAT
 4689 TATGACATCAATTTTATACAAATTTTCATCATTCATATATATATCATATGATCAATCACA
 55 4749 TAACTATTAAAAATTCATACCATTCAGAAGCTTTTCCAACATCAAAAGCAATATTTACTT
 4809 TCAAACTATCAAAATATAATTATTATTTAATAAAGCTT

SEQ ID NO:7 Scarlet Runner Bean C541 amino acid sequence

ATG TCT CCA TTC TGT AGA

M S P F C R

1
3172 AAC TTT TCA ATG GCA TGG GTG CTT ATG GCA TTT GTG TTT
7 N F S M A W V L M A F V L F
5 3214 GCA AAC AGT GCT ATG CCC ACA AAT ACF GTT GGG GTA
21 A N S A M P T N G S T V G V
3256 AAA AAC ATG TTG GGT AAA TTG ATG CTA AAC GTT TTA TGT
35 K N M L G G K L M L N V L C
10 3298 CCC CAT ATT GAT AAG CAA CAC ATT ATC CCG AAT GGT GGT TCA
49 P H I D K Q H I I P N G G S
3340 TTT GAG TGG AAG TAC AAT GGT GGT GCT CCA CCA ATA GGA CAA
63 F E W K Y N G G A P P I G Q
15 3382 TCA CCA TTC ATG TGT TTC TTT CGG TGG AAT AAT GTT CAT CAC
77 S P F M C F F R W N N V H H
3424 TCC CTT GAT CTG TGT TCA CCA AGC AAG TAT ACT GGT TGT GAA
91 S L D L C S P S K Y T G C E
3466 AAT GCC ATT TGG GAA ATC AAA GAA AAG CAA TTT GT AGG TAC
105 N A I W E I K E K Q F C R Y
20 3508 AGA GGT GGA CCT ATT AAT TAT TTT TGC TAT GAC TGG GAT GAT
119 R G G P I N Y F C Y D W D D
3550 TAG

SEQ ID NO:8 Arabidopsis C541 genomic region

142000 TTATCTTATTTCATATAATTTGTTGTTTACTTTCAAAATTTTAAATTTT
141950 TTATATTTTATCTTTTACAGTTTAAAAATTAATAAAATGAAACTTTTTC
141900 TTAATATGTGTAAAAATATAAAATCAAAAAAGTTGTTATATGGTACATGGC
30 141850 ACAATCTTATAAAATTAATAATTTGAAAACGATACTTTATATAATAAAAT
141800 ATCTTAGTTGACATTTTATTAGTGTGTTTCAATCATATTTTGTGTTGCTT
141750 GATAAGCGTAAAAACAAATCAAACTTAACGATATCTTATATAATAAAATTA
141700 TCTTAGTTGACATTTTATTAGTGTCTTCAATCATATCTTTGTTGCTTG
141650 ATAAGCGTAAAAACAAATCAAGTAAAGTTGGGCACCTCAATTTGTTTAAAA
35 141600 AAGTTTGGGTACCTCAAAAATTAATAGTCTTGTGAGATTCTTCAAAAA
141550 AAATCTGGAAGAATTTATGAAGAAGGGGGGGAGGGGGGGAGGGGGGG
141500 AAGTGAAGATGAATATTCAACAAAAGAGGGGTAGGCATGATGTTAAGTGA
141450 TTAATAAACTATGTTAATGGAGACAATTTCTGTTAAACAAACCCGTTAAT
141400 TGAACACGATAGCATCTTCTCTAACAATGTAAACGATATTGTTTATTC
141350 ATAACTACTCATTAATTTCTGAGTTTCAAATCATATAAGATTATGGGG
40 141300 GGTGTATTCAATTAAGGATTTGAAATGATTTGTATTAATAATGACAAATCC
141250 CATGTTATTTCACCAATGAATTTGAAAAACTTTTAAAAATCAAGTTTA
141200 TTGATTTAGTGATTTTAAAAATGTACAACCAACCCACTGTTATTGGAAAC
141150 ATTTTAAAGTAGTGGATTAAAAATGACTTGTGATGATTTGGGTGGGATTG
45 141100 AGAAAAATTTCTTAGTTAAGAATTCACCAATCCCAATCTCATGTTTCAAG
141050 TAGAATTTGGGAGAATTTAATAACAAATCTCTAATTTACCAAAAGTCA
141000 CCAAAATCATTTAAAACTCATTAATTAATGATTTCAATCTCCAG
140950 TTGAATACATCCCTTGGAAATAGAGATTTTGCTCGATTTGGGACCTAAG
140900 ATTGAATTTTGGGGATTAGTTTAAATCGTTACAAACAAATGACATCGTAT
140850 TATTGTTATAGGAACAATGTGCTTTTCAGTTGACATGTATGTTAATAGA
50 140800 AAATTAACCTATTAAACGGGATTGCTAACCCATTAAACATCGTAACATA
140750 ATGGTCAAGTCAATAAAAGTTTGGTATTATTGAAAAGTCAACGTAAGT
140700 TTGATATTTATTGAAAAGTCAACATAAAATTTGATATCTTATTTCGTTTC
140650 GACAGACATAAGGATTACATCAATGTTTTTAATAAATTAAGAGATTATTA
140600 TGACATTTTCCATTTAAAAATGGCAATGTTTCGAAACCAAGATATC
55 140550 AAAATTGACATACCTAATTCATCTACATTTGTTGACAGCAATTCACGT
140500 GCCTTGACCAATGGCACATCTGGCAATCATCAATTTTAAGGAAAAGG
140450 TAGATTCCGATACAATATAATGGAAATAGGTGGAAGGATCATTGACTAC
140400 TTGACTTGTAACAAACACACACAGTATATAACTCATTGACATTTACAA

140350 ACAACATTGTGCTAGCTTAAACTCCCTCTCCTATTCAAAAAA **ATG**
 1 M
 140305 GAT ATT CCA AAG CAA TAT CTA TCA CTA TTC ATA TTG
 2 D I P K Q Y L S L F I L
 5 140269 ATT ATC TTC ATA ACT ACA AAA TTA TCA CAA GCC GAC
 14 I I F I T T K L S Q A D
 140233 CAT AAA AAC GAC ATT CCA GTT CCC AAC GAT CCA TCA
 26 H K N D I P V P N D P S
 10 140197 TCA ACA AAT TCT GTG TTT CCT ACC TCG AAA AGA ACC
 38 S T N S V F P T S K R T
 140161 GTG GAA ATC AAT AAT GAT CTC GGT AAT CAG CTA ACG
 50 V E I N N D L G N Q L T
 140125 TTA CTG TAT CAT TGT AAA TCA AAA GAC GAT GAT TTA
 62 L L Y H C K S K D D D L
 15 140089 GGT AAC CGG ACT CTG CAA CCA GGT GAG TCG TGG TCT
 74 G N R T L Q P G E S W S
 140053 TTT AGT TTC GGG CGT CAA TTC TTT GGA AGG ACG TTG
 86 F S F G R Q F F G R T L
 20 140017 TAT TTT TGT AGT TTT AGT TGG CCA AAT GAA TCG CAT
 98 Y F C S F S W P N E S H
 139981 TCG TTC GAT ATA TAT AAA GAC CAT GAG AAT AGC GGC
 110 S F D I Y K D H R D S G
 139945 GGT GAT AAC AAG TGC GAG AGC GAC AGG TGT GTG TGG
 122 G D N K C E S D R C V W
 25 139909 AAG ATA AGA AGA AAC GGA CCT TGT AGG TTT AAC GAT
 134 K I R R N G P C R F N D
 139873 GAA ACG AAG CAG TTT GAT CTT TGT TAT CPT TGG AAT
 146 E T K Q F D L C Y P W N
 30 139837 AAA TCT TTG TAT TGA CAACAATATGCTGATGTTCTGCTCTTTAC
 158 K S L Y •
 139793 GACTCATGGAGTTTTCATTGTTTGAACAATAATATAAAACATATAAAAT
 139743 TCTATTATCCAAAGTTCCAACTTATAATAATTGATAATCATATCATATT
 139693 ATCATCTTAAAGCATTCATGCTACAAAGATAATACCCCCAAGCTATTTTA
 139643 CATTTAAAGCTGAACAGAGACACAATACTAACGATAAAAGTTCGTAGTA
 35 139593 TCTTTATGCAACCATACATACATATACACAAAGATAGACAGGTAGTGCC
 139543 TAATAATTCTACTTTGGGTGAGGTATGAACAGCAGCAACAGTAGATACCAT
 139493 TGTATCCATACCACATATTATGAGGCCCTCTGCAGATTTTGTAGTAAC
 139443 CATGCTCTCCACATCGCTCCCCAGAGTTCTTGATAATCCAA

SEQ ID NO:9 Arabidopsis C541 amino acid sequence

1 M
 140305 GAT ATT CCA AAG CAA TAT CTA TCA CTA TTC ATA TTG
 2 D I P K Q Y L S L F I L
 45 140269 ATT ATC TTC ATA ACT ACA AAA TTA TCA CAA GCC GAC
 14 I I F I T T K L S Q A D
 140233 CAT AAA AAC GAC ATT CCA GTT CCC AAC GAT CCA TCA
 26 H K N D I P V P N D P S
 50 140197 TCA ACA AAT TCT GTG TTT CCT ACC TCG AAA AGA ACC
 38 S T N S V F P T S K R T
 140161 GTG GAA ATC AAT AAT GAT CTC GGT AAT CAG CTA ACG
 50 V E I N N D L G N Q L T
 140125 TTA CTG TAT CAT TGT AAA TCA AAA GAC GAT GAT TTA
 62 L L Y H C K S K D D D L
 55 140089 GGT AAC CGG ACT CTG CAA CCA GGT GAG TCG TGG TCT
 74 G N R T L Q P G E S W S
 140053 TTT AGT TTC GGG CGT CAA TTC TTT GGA AGG ACG TTG
 86 F S F G R Q F F G R T L

140017 TAT TTT TGT AGT TTT AGT TGG CCA AAT GAA TCG CAT
 98 Y F C S F S W P N E S H
 139981 TCG TTC GAT ATA TAT AAA GAC CAT CGA GAT AGC GGC
 110 S F D I Y K D H R D S G
 139945 GGT GAT AAC AAG TGC GAG AGC GAC AGG TGT GTG TGG
 122 G D N K C E S D R C V W
 139909 AAG ATA AGA AGA AAC GGA CCT TGT AGG TTT AAC GAT
 134 K I R R N G P C R F N D
 139873 GAA ACG AAG CAG TTT GAT CTT TGT TAT CCT TGG AAT
 146 E T K Q F D L C Y P W N
 139837 AAA TCT TTG TAT TGA CAACAATATGCTGATGTTCTGCTTTTAC
 158 K S L Y •

SEQ ID NO:10 promoter control element

GAAAAGCGAA

SEQ ID NO:11 promoter control element

GAAAAGCCAA

WHAT IS CLAIMED IS:

1. An isolated polynucleotide which specifically modulates transcription in a plant suspensor cell and/or basal region of a plant embryo, the polynucleotide comprising a promoter control element comprising,

(a) a nucleotide sequence having at least 50% sequence identity to nucleotides 3329 to 3475 of SEQ ID NO:1; or

(b) a nucleotide sequence which hybridizes to nucleotides 3329 to 3475 of SEQ ID NO:1 under a condition establishing a T_m minus 20°C.

2. The isolated polynucleotide of claim 1, comprising

(a) a nucleotide sequence having at least 50% sequence identity to SEQ ID NO:1; or

(b) a nucleotide sequence which hybridizes to SEQ ID NO:1 under a condition establishing a T_m minus 20°C.

3. The isolated polynucleotide of claim 1, wherein the polynucleotide comprises nucleotides 3324 to 3580 of SEQ ID NO:1.

4. An expression cassette comprising a promoter sequence, the promoter sequence comprising,

i. a nucleotide sequence having at least 50% sequence identity to nucleotides 3329 to 3475 of SEQ ID NO:1; and

ii. a promoter polynucleotide with at least basal promoter activity, which promoter polynucleotide is operably linked to a heterologous polynucleotide,

wherein when the expression cassette is inserted into a plant, the heterologous polynucleotide is specifically expressed in a suspensor cell and/or basal region of a plant embryo.

5. The expression cassette of claim 4, wherein the nucleotide sequence comprises nucleotides 3329 to 3475 of SEQ ID NO:1

6. An isolated polynucleotide which specifically modulates transcription in a plant suspensor cell and/or basal region of a plant embryo, the polynucleotide comprising a promoter comprising,

4 (a) a nucleotide sequence having at least 50% sequence identity to SEQ ID
5 NO:1 or nucleotides 1 to 3154 of SEQ ID NO:6;

6 (b) a nucleotide sequence which hybridizes to SEQ ID NO:1 or
7 nucleotides 1 to 3154 of SEQ ID NO:6 under a condition establishing a T_m minus 20°C.

1 7. The isolated polynucleotide of claim 6, wherein the promoter
2 comprises SEQ ID NO:1.

1 8. The isolated polynucleotide of claim 6, wherein the promoter
2 comprises nucleotides 1 to 3154 of SEQ ID NO:6.

1 9. The isolated polynucleotide of claim 6, further comprising a G564
2 polynucleotide operably linked to the promoter.

1 10. The isolated polynucleotide of claim 9, wherein the isolated
2 polynucleotide comprises SEQ ID NO:2.

1 11. The isolated polynucleotide of claim 6, further comprising a G541
2 polynucleotide operably linked to the promoter.

1 12. The isolated polynucleotide of claim 9, wherein the isolated
2 polynucleotide comprises SEQ ID NO:6.

1 13. The isolated polynucleotide of claim 6, further comprising a
2 heterologous polynucleotide operably linked to the promoter.

1 14. A vector comprising a promoter of claim 6 operably linked to a
2 heterologous nucleic acid sequence.

1 15. The vector of claim 14, wherein the promoter is SEQ ID NO:1.

1 16. The vector of claim 14, wherein the promoter comprises nucleotides 1
2 to 3154 of SEQ ID NO:6.

1 17. A host cell comprising a promoter of claim 6.

1 18. The host cell of claim 17, wherein the promoter comprises SEQ ID
2 NO:1.

1 19. The host cell of claim 17, wherein the promoter comprises nucleotides
2 1 to 3154 of SEQ ID NO:6.

1 20. The host cell of claim 17, wherein the host cell is a plant cell.

1 21. A host cell comprising the vector of claim 14.

1 22. A plant comprising the polynucleotide of claim 13.

1 23. A plant of claim 22, wherein the promoter comprises SEQ ID NO:1.

1 24. A plant of claim 22, wherein the promoter comprises nucleotides 1 to
2 3154 of SEQ ID NO:6.

1 25. A plant comprising a vector of claim 14.

1 26. A method of modulating transcription in a plant suspensor cell and/or
2 basal region of a plant embryo, the method comprising introducing into a plant an expression
3 cassette comprising the promoter of claim 1.

1 27. The method of claim 26, wherein the promoter comprises SEQ ID
2 NO:1.

1 28. The method of claim 26, wherein the promoter comprises nucleotides 1
2 to 3154 of SEQ ID NO:6.

1 29. The method of claim 26, wherein a G564 polynucleotide is operably
2 linked to the promoter.

1 30. The method of claim 26, wherein the promoter is operably linked to a
2 heterologous polynucleotide.

1 31. The method of claim 30, wherein the promoter is operably linked to
2 the heterologous polynucleotide in an antisense orientation.

1 32. An isolated nucleic acid comprising a polynucleotide, or complement
2 thereof, encoding a G564 polypeptide exhibiting at least 50% sequence identity to SEQ ID
3 NO:3.

1 33. The isolated nucleic acid of claim 32, wherein the G564 polypeptide
2 comprises SEQ ID NO:3.

1 34. The isolated nucleic acid of claim 32, wherein the nucleic acid further
2 comprises a promoter operably linked to the polynucleotide.

1 35. The isolated nucleic acid of claim 34, wherein the promoter is a
2 constitutive promoter.

1 36. The isolated nucleic acid of claim 34, wherein the polynucleotide is
2 linked to the promoter in an antisense orientation.

1 37. An isolated nucleic acid comprising a polynucleotide, or complement
2 thereof, encoding a C541 polypeptide exhibiting at least 50% sequence identity to SEQ ID
3 NO:7.

1 38. The isolated nucleic acid of claim 37, wherein the C541 polypeptide
2 comprises SEQ ID NO:7.

1 39. The isolated nucleic acid of claim 37, wherein the nucleic acid further
2 comprises a promoter operably linked to the polynucleotide.

1 40. The isolated nucleic acid of claim 39, wherein the promoter is a
2 constitutive promoter.

1 41. The isolated nucleic acid of claim 39, wherein the polynucleotide is
2 linked to the promoter in an antisense orientation.

1 42. An expression cassette comprising a promoter operably linked to a
2 heterologous polynucleotide sequence, or a complement thereof, encoding a G564
3 polypeptide exhibiting at least 50% sequence identity to SEQ ID NO:3.

1 43. The expression cassette of claim 42, wherein the G564 polypeptide
2 comprises SEQ ID NO: 3.

1 44. The expression cassette of claim 42, wherein the G564 polynucleotide
2 comprises nucleotides 4242 to 4901 of SEQ ID NO: 2.

1 45. The expression cassette of claim 42, wherein the promoter is a
2 constitutive promoter.

1 46. The expression cassette of claim 42, wherein the polynucleotide is
2 linked to the promoter in an antisense orientation.

1 47. An expression cassette comprising a promoter operably linked to a
2 heterologous polynucleotide, or a complement thereof, encoding a C541 polypeptide
3 exhibiting at least 50% sequence identity to SEQ ID NO:7.

1 48. The expression cassette of claim 47, wherein the C541 polypeptide
2 comprises SEQ ID NO:7.

1 49. The expression cassette of claim 47, wherein the C541 polynucleotide
2 comprises nucleotides 3155 to 3552 of SEQ ID NO: 6.

1 50. The expression cassette of claim 47, wherein the promoter is a
2 constitutive promoter.

1 51. The expression cassette of claim 47, wherein the polynucleotide is
2 linked to the promoter in an antisense orientation.

1 52. A host cell comprising an exogenous nucleic acid comprising a
2 polynucleotide, or complement thereof, encoding a G564 polypeptide exhibiting at least 80%
3 sequence identity to SEQ ID NO:3.

1 53. The host cell of claim 52, wherein the nucleic acid further comprises a
2 promoter operably linked to the polynucleotide.

1 54. The host cell of claim 53, wherein the promoter is constitutive.

1 55. The host cell of claim 53, wherein the promoter is operably linked to
2 the polynucleotide in an antisense orientation.

1 56. A host cell comprising an exogenous nucleic acid comprising a
2 polynucleotide, or complement thereof, encoding a C541 polypeptide exhibiting at least 50%
3 sequence identity to SEQ ID NO:7.

1 57. The host cell of claim 56, wherein the nucleic acid further comprises a
2 promoter operably linked to the polynucleotide.

1 58. The host cell of claim 57, wherein the promoter is constitutive.

1 59. The host cell of claim 57, wherein the promoter is operably linked to
2 the polynucleotide in an antisense orientation.

1 60. A transgenic plant comprising a recombinant expression cassette, the
2 recombinant expression cassette comprising a polynucleotide, or complement thereof,
3 encoding a G564 polypeptide exhibiting at least 50% sequence identity to SEQ ID NO:3.

1 61. The transgenic plant of claim 60, wherein the G564 polypeptide
2 comprises SEQ ID NO:3.

1 62. The transgenic plant of claim 60, wherein the polynucleotide
2 comprises nucleotides 4242 to 4901 of SEQ ID NO:2.

1 63. The transgenic plant of claim 60, wherein the nucleic acid further
2 comprises a promoter operably linked to the polynucleotide.

1 64. The transgenic plant of claim 63, wherein the promoter is a constitutive
2 promoter.

1 65. The transgenic plant of claim 60, wherein the polynucleotide is linked
2 to the promoter in an antisense orientation.

1 66. A transgenic plant comprising a recombinant expression cassette, the
2 recombinant expression cassette comprising a polynucleotide, or complement thereof,
3 encoding a C541 polypeptide exhibiting at least 50% sequence identity to SEQ ID NO:7.

1 67. The transgenic plant of claim 66, wherein the G541 polypeptide
2 comprises SEQ ID NO:7.

1 68. The transgenic plant of claim 66, wherein the polynucleotide
2 comprises nucleotides 3155 to 3552 of SEQ ID NO: 6.

- 1 69. The transgenic plant of claim 66, wherein the nucleic acid further
2 comprises a promoter operably linked to the polynucleotide.
- 1 70. The transgenic plant of claim 69, wherein the promoter is a constitutive
2 promoter.
- 1 71. The transgenic plant of claim 66, wherein the polynucleotide is linked
2 to the promoter in an antisense orientation.
- 1 72. An isolated polypeptide comprising an amino acid sequence at least
2 80% identical to SEQ ID NO:3.
- 1 73. The isolated polypeptide of claim 72, wherein the polypeptide is SEQ
2 ID NO:3.
- 1 74. An isolated polypeptide comprising an amino acid sequence at least
2 80% identical to SEQ ID NO:7.
- 1 75. The isolated polypeptide of claim 74, wherein the polypeptide is SEQ
2 ID NO:7.
- 1 76. An antibody capable of binding the isolated polypeptide of claim 72.
- 1 77. An antibody capable of binding the isolated polypeptide of claim 74.
- 1 78. A method of introducing an isolated polynucleotide into a host cell
2 comprising:
3 (a) providing an isolated polynucleotide according to claim 1; and
4 (b) contacting the polynucleotide with the host cell under
5 conditions that permit insertion of the polynucleotide into the host cell.
- 1 79. A method of detecting a polynucleotide in a sample, comprising
2 (a) providing an isolated polynucleotide according to claim 1;
3 (b) contacting the isolated polynucleotide with a sample under conditions
4 which permit a comparison of the sequence of the isolated polynucleotide with the sequence
5 of DNA in the sample; and
6 (c) analyzing the result of the comparison.

1 80. The method of claim 79, wherein the isolated polynucleotide and the
2 sample are contacted under conditions which permit the formation of a duplex between
3 complementary nucleic acid sequences.

POLYNUCLEOTIDES USEFUL FOR MODULATING TRANSCRIPTION

ABSTRACT OF THE DISCLOSURE

The invention provides polynucleotides for expression of genes in suspensor cells in plants and methods for using such polynucleotides.

SF 1160138 v1

1 23
2 24
3 25
4 26
5 27
6 28
7 29
8 30
9 31
10 32
11 33
12 34
13 35
14 36
15 37
16 38
17 39
18 40
19 41
20 42
21 43
22 44
23 45
24 46
25 47
26 48
27 49
28 50
29 51
30 52
31 53
32 54
33 55
34 56
35 57
36 58
37 59
38 60
39 61
40 62
41 63
42 64
43 65
44 66
45 67
46 68
47 69
48 70
49 71
50 72
51 73
52 74
53 75
54 76
55 77
56 78
57 79
58 80
59 81
60 82
61 83
62 84
63 85
64 86
65 87
66 88
67 89
68 90
69 91
70 92
71 93
72 94
73 95
74 96
75 97
76 98
77 99
78 100
79 101
80 102
81 103
82 104
83 105
84 106
85 107
86 108
87 109
88 110
89 111
90 112
91 113
92 114
93 115
94 116
95 117
96 118
97 119
98 120
99 121
100 122
101 123
102 124
103 125
104 126
105 127
106 128
107 129
108 130
109 131
110 132
111 133
112 134
113 135
114 136
115 137
116 138
117 139
118 140
119 141
120 142
121 143
122 144
123 145
124 146
125 147
126 148
127 149
128 150
129 151
130 152
131 153
132 154
133 155
134 156
135 157
136 158
137 159
138 160
139 161
140 162
141 163
142 164
143 165
144 166
145 167
146 168
147 169
148 170
149 171
150 172
151 173
152 174
153 175
154 176
155 177
156 178
157 179
158 180
159 181
160 182
161 183
162 184
163 185
164 186
165 187
166 188
167 189
168 190
169 191
170 192
171 193
172 194
173 195
174 196
175 197
176 198
177 199
178 200
179 201
180 202
181 203
182 204
183 205
184 206
185 207
186 208
187 209
188 210
189 211
190 212
191 213
192 214
193 215
194 216
195 217
196 218
197 219
198 220
199 221
200 222
201 223
202 224
203 225
204 226
205 227
206 228
207 229
208 230
209 231
210 232
211 233
212 234
213 235
214 236
215 237
216 238
217 239
218 240
219 241
220 242
221 243
222 244
223 245
224 246
225 247
226 248
227 249
228 250
229 251
230 252
231 253
232 254
233 255
234 256
235 257
236 258
237 259
238 260
239 261
240 262
241 263
242 264
243 265
244 266
245 267
246 268
247 269
248 270
249 271
250 272
251 273
252 274
253 275
254 276
255 277
256 278
257 279
258 280
259 281
260 282
261 283
262 284
263 285
264 286
265 287
266 288
267 289
268 290
269 291
270 292
271 293
272 294
273 295
274 296
275 297
276 298
277 299
278 300
279 301
280 302
281 303
282 304
283 305
284 306
285 307
286 308
287 309
288 310
289 311
290 312
291 313
292 314
293 315
294 316
295 317
296 318
297 319
298 320
299 321
300 322
301 323
302 324
303 325
304 326
305 327
306 328
307 329
308 330
309 331
310 332
311 333
312 334
313 335
314 336
315 337
316 338
317 339
318 340
319 341
320 342
321 343
322 344
323 345
324 346
325 347
326 348
327 349
328 350
329 351
330 352
331 353
332 354
333 355
334 356
335 357
336 358
337 359
338 360
339 361
340 362
341 363
342 364
343 365
344 366
345 367
346 368
347 369
348 370
349 371
350 372
351 373
352 374
353 375
354 376
355 377
356 378
357 379
358 380
359 381
360 382
361 383
362 384
363 385
364 386
365 387
366 388
367 389
368 390
369 391
370 392
371 393
372 394
373 395
374 396
375 397
376 398
377 399
378 400
379 401
380 402
381 403
382 404
383 405
384 406
385 407
386 408
387 409
388 410
389 411
390 412
391 413
392 414
393 415
394 416
395 417
396 418
397 419
398 420
399 421
400 422
401 423
402 424
403 425
404 426
405 427
406 428
407 429
408 430
409 431
410 432
411 433
412 434
413 435
414 436
415 437
416 438
417 439
418 440
419 441
420 442
421 443
422 444
423 445
424 446
425 447
426 448
427 449
428 450
429 451
430 452
431 453
432 454
433 455
434 456
435 457
436 458
437 459
438 460
439 461
440 462
441 463
442 464
443 465
444 466
445 467
446 468
447 469
448 470
449 471
450 472
451 473
452 474
453 475
454 476
455 477
456 478
457 479
458 480
459 481
460 482
461 483
462 484
463 485
464 486
465 487
466 488
467 489
468 490
469 491
470 492
471 493
472 494
473 495
474 496
475 497
476 498
477 499
478 500
479 501
480 502
481 503
482 504
483 505
484 506
485 507
486 508
487 509
488 510
489 511
490 512
491 513
492 514
493 515
494 516
495 517
496 518
497 519
498 520
499 521
500 522
501 523
502 524
503 525
504 526
505 527
506 528
507 529
508 530
509 531
510 532
511 533
512 534
513 535
514 536
515 537
516 538
517 539
518 540
519 541
520 542
521 543
522 544
523 545
524 546
525 547
526 548
527 549
528 550
529 551
530 552
531 553
532 554
533 555
534 556
535 557
536 558
537 559
538 560
539 561
540 562
541 563
542 564
543 565
544 566
545 567
546 568
547 569
548 570
549 571
550 572
551 573
552 574
553 575
554 576
555 577
556 578
557 579
558 580
559 581
560 582
561 583
562 584
563 585
564 586
565 587
566 588
567 589
568 590
569 591
570 592
571 593
572 594
573 595
574 596
575 597
576 598
577 599
578 600
579 601
580 602
581 603
582 604
583 605
584 606
585 607
586 608
587 609
588 610
589 611
590 612
591 613
592 614
593 615
594 616
595 617
596 618
597 619
598 620
599 621
600 622
601 623
602 624
603 625
604 626
605 627
606 628
607 629
608 630
609 631
610 632
611 633
612 634
613 635
614 636
615 637
616 638
617 639
618 640
619 641
620 642
621 643
622 644
623 645
624 646
625 647
626 648
627 649
628 650
629 651
630 652
631 653
632 654
633 655
634 656
635 657
636 658
637 659
638 660
639 661
640 662
641 663
642 664
643 665
644 666
645 667
646 668
647 669
648 670
649 671
650 672
651 673
652 674
653 675
654 676
655 677
656 678
657 679
658 680
659 681
660 682
661 683
662 684
663 685
664 686
665 687
666 688
667 689
668 690
669 691
670 692
671 693
672 694
673 695
674 696
675 697
676 698
677 699
678 700
679 701
680 702
681 703
682 704
683 705
684 706
685 707
686 708
687 709
688 710
689 711
690 712
691 713
692 714
693 715
694 716
695 717
696 718
697 719
698 720
699 721
700 722
701 723
702 724
703 725
704 726
705 727
706 728
707 729
708 730
709 731
710 732
711 733
712 734
713 735
714 736
715 737
716 738
717 739
718 740
719 741
720 742
721 743
722 744
723 745
724 746
725 747
726 748
727 749
728 750
729 751
730 752
731 753
732 754
733 755
734 756
735 757
736 758
737 759
738 760
739 761
740 762
741 763
742 764
743 765
744 766
745 767
746 768
747 769
748 770
749 771
750 772
751 773
752 774
753 775
754 776
755 777
756 778
757 779
758 780
759 781
760 782
761 783
762 784
763 785
764 786
765 787
766 788
767 789
768 790
769 791
770 792
771 793
772 794
773 795
774 796
775 797
776 798
777 799
778 800
779 801
780 802
781 803
782 804
783 805
784 806
785 807
786 808
787 809
788 810
789 811
790 812
791 813
792 814
793 815
794 816
795 817
796 818
797 819
798 820
799 821
800 822
801 823
802 824
803 825
804 826
805 827
806 828
807 829
808 830
809 831
810 832
811 833
812 834
813 835
814 836
815 837
816 838
817 839
818 840
819 841
820 842
821 843
822 844
823 845
824 846
825 847
826 848
827 849
828 850
829 851
830 852
831 853
832 854
833 855
834 856
835 857
836 858
837 859
838 860
839 861
840 862
841 863
842 864
843 865
844 866
845 867
846 868
847 869
848 870
849 871
850 872
851 873
852 874
853 875
854 876
855 877
856 878
857 879
858 880
859 881
860 882
861 883
862 884
863 885
864 886
865 887
866 888
867 889
868 890
869 891
870 892
871 893
872 894
873 895
874 896
875 897
876 898
877 899
878 900
879 901
880 902
881 903
882 904
883 905
884 906
885 907
886 908
887 909
888 910
889 911
890 912
891 913
892 914
893 915
894 916
895 917
896 918
897 919
898 920
899 921
900 922
901 923
902 924
903 925
904 926
905 927
906 928
907 929
908 930
909 931
910 932
911 933
912 934
913 935
914 936
915 937
916 938
917 939
918 940
919 941
920 942
921 943
922 944
923 945
924 946
925 947
926 948
927 949
928 950
929 951
930 952
931 953
932 954
933 955
934 956
935 957
936 958
937 959
938 960
939 961
940 962
941 963
942 964
943 965
944 966
945 967
946 968
947 969
948 970
949 971
950 972
951 973
952 974
953 975
954 976
955 977
956 978
957 979
958 980
959 981
960 982
961 983
962 984
963 985
964 986
965 987
966 988
967 989
968 990
969 991
970 992
971 993
972 994
973 995
974 996
975 997
976 998
977 999
978 1000
979 1001
980 1002
981 1003
982 1004
983 1005
984 1006
985 1007
986 1008
987 1009
988 1010
989 1011
990 1012
991 1013
992 1014
993 1015
994 1016
995 1017
996 1018
997 1019
998 1020
999 1021
1000 1022
1001 1023
1002 1024
1003 1025
1004 1026
1005 1027
1006 1028
1007 1029
1008 1030
1009 1031
1010 1032
1011 1033
1012 1034
1013 1035
1014 1036
1015 1037
1016 1038
1017 1039
1018 1040
1019 1041
1020 1042
1021 1043
1022 1044
1023 1045
1024 1046
1025 1047
1026 1048
1027 1049
1028 1050
1029 1051
1030 1052
1031 1053
1032 1054
1033 1055
1034 1056
1035 1057
1036 1058
1037 1059
1038 1060
1039 1061
1040 1062
1041 1063
1042 1064
1043 1065
1044 1066
1045 1067
1046 1068
1047 1069
1048 1070
1049 1071
1050 1072
1051 1073
1052 1074
1053 1075
1054 1076
1055 1077
1056 1078
1057 1079
1058 1080
1059 1081
1060 1082
1061 1083
1062 1084
1063 1085
1064 1086
1065 1087
1066 1088
1067 1089
1068 1090
1069 1091
1070 1092
1071 1093
1072 1094
1073 1095
1074 1096
1075 1097
1076 1098
1077 1099
1078 1100
1079 1101
1080 1102
1081 1103
1082 1104
1083 1105
1084 1106
1085 1107
1086 1108
1087 1109
1088 1110
1089 1111
1090 1112
1091 1113
1092 1114
1093 1115
1094 1116
1095 1117
1096 1118
1097 1119
1098 1120
1099 1121
1100 1122
1101 1123
1102 1124
1103 1125
1104 1126
1105 1127
1106 1128
1107 1129
1108 1130
1109 1131
1110 1132
1111 1133
1112 1134
1113 1135
1114 1136
1115 1137
1116 1138
1117 1139
1118 1140
1119 1141
1120 1142
1121 1143
1122 1144
1123 1145
1124 1146
1125 1147
1126 1148
1127 1149
1128 1150
1129 1151
1130 1152
1131 1153
1132 1154
1133 1155
1134 1156
1135 1157
1136 1158
1137 1159
1138 1160
1139 1161
1140 1162
1141 1163
1142 1164
1143 1165
1144 1166
1145 1167
1146 1168
1147 1169
1148 1170
1149 1171
1150 1172
1151 1173
1152 1174
1153 1175
1154 1176
1155 1177
1156 1178
1157 1179
1158 1180
1159 1181
1160 1182
1161 1183
1162 1184
1163 1185
1164 1186
1165 1187
1166 1188
1167 1189
1168 1190
1169 1191
1170 1192
1171 1193
1172 1194
1173 1195
1174 1196
1175 1197
1176 1198
1177 1199
1178 1200
1179 1201
1180 1202
1181 1203
1182 1204
1183 1205
1184 1206
1185 1207
1186 1208
1187 1209
1188 1210
1189 1211
1190 1212
1191 1213
1192 1214
1193 1215
1194 1216
1195 1217
1196 1218
1197 1219
1198 1220
1199 1221
1200 1222
1201 1223
1202 1224
1203 1225
1204 1226
1205 1227
1206 1228
1207 1229
1208 1230
1209 1231
1210 1232
1211 1233
1212 1234
1213 1235
1214 1236
1215 1237
1216 1238
1217 1239
1218 1240
1219 1241
1220 1242
1221 1243
1222 1244
1223 1245
1224 1246
1225 1247
1226 1248
1227 1249
1228 1250
1229 1251
1230 1252
1231 1253
1232 1254
1233 1255
1234 1256
1235 1257
1236 1258
1237 1259
1238 1260
1239 1261
1240 1262
1241 1263
1242 1264
1243 1265
1244 1266
1245 1267
1246 1268
1247 1269
1248 1270
1249 1271
1250 1272
1251 1273
1252 1274
1253 1275
1254 1276
1255 1277
1256 1278
1257 1279
1258 1280
1259 1281
1260 1282
1261 1283
1262 1284
1263 1285
1264 1286
1265 1287
1266 1288
1267 1289
1268 1290
1269 1291
1270 1292
1271 1293
1272 1294
1273 1295
1274 1296
1275 1297
1276 1298
1277 1299
1278 1300
1279 1301
1280 1302
1281 1303
1282 1304
1283 1305
1284 1306
1285 1307
1286 1308
1287 1309
1288 1310
1289 1311
1290 1312
1291 1313
1292 1314
1293 1315
1294 1316
12

FIGURE 1

SCHEMATIC OF A GENE




FIGURE 2

-4242 GCATGCACTG CCACAAGTAG TGAACATCATG GTTTTACCTC CTCAGTAGA
 -4192 AAACCTTTTG AGTGAATTTG AAGATTATT CTCCCAAGAA GGACCCATTG
 -4142 GGCTTCTCC TCCTAGGGGG ATAGAACATC AAATTGACTT TATACCGGGG
 -4092 GCAAGCCTAC CAAATAGGCC TCCTTATAGA ACCAAGCCCG AGGAAACAAA
 -4042 GGAGATAGAA TCACAAGTTC AAGACTTGT GGAGAAGGGT TGGGTTCAAA
 -3992 AGAGCCTAAG CCCTTGTGCT GTACCTGTCT TGTGTGTGCC AAAAAAAGAT
 -3942 GGAAAAATGGC GTATGTGTTG TGATTGTAGA GCAATCAACA ACATCCCAT
 -3892 CAAGTATAGG CATCCAATCC CAAGGCTTGA CGATATGCTT GATGAATTGC
 -3842 ATGGGTCAAC TCTATTCTCC AAAATTGACC TTAAAGTGG ATATCACCAA
 -3792 ATTCGAATCA AGGAGGGTGA TGAGTGGAAA ACCGCTTTTA AGACCAAATT
 -3742 TGGATTATAT GAGTGGTTGG TGATGCCCTT TGGTCTTACT AACGCTCCAA
 -3692 GTACATTCAT GAGGCTTATG AATCACACCT TGAGGGATTG TATAGGTAAA
 -3642 TATGTAGTAG TTTATTTTGA TGATATCTTA GTATATAGTA AAACCCCTAGA
 -3592 AGACCATCTA AGTCACCTTA GGGAGTTCT TCTAGTTCTT AGGAAAAATA
 -3542 GTCTTTTTCG CAATAGGGAT AAGTGTACCT TTTGTGTAGA TAGCGTAGTC
 -3492 TTTTTAGGCT TTATAGTAAA CCAAAAGGGG GTGCATGTAG ATCCCGAGAA
 -3442 AATCAAAGCC ATCCGCGAGT GGCACCAATCC ACAAATGTA AGTGTGTAGA
 -3392 GAAGTTTTC TGGGTTAGCT AGCTTCTATA GAAGGTTTGT TCCCAATTTT
 -3342 TTAGCCTAG CTCTCCCTT GAATGAACCT GTAAAAAAAG ATGTTGCATT
 -3292 TTGTTGGAAT GAAAAGCATG AGCAAGCCCTT TCAAAGGCTA AAAGCTCACT
 -3242 CACCAATGCA CCCATCCTAT CTCTTCCAAA CTTTGGGAGA
 -3192 TAGAGTGTGA TGCATCGGGA GTAGGCATAG TGCGGTTTGT TTGCAAGGTTG
 -3142 GACACCCCTT GCTTATTTTA GTGAAAAACT CCATGGTGCC ACCCTGACTA
 -3092 CCCCACTAT GACAAAGACT CTATGCTCTT GTGCGACCTT AAAGACTTTGG
 -3042 GGAACACTAC CTTGnGTCCC AAAGAATTG GntATCCATA GTGCATCAGA
 -2992 GTCTTTAAAA TATTTAAAGG GCCAACACAA GCTCAATAAG AGACATGCTA
 -2942 AATGGATGGA ATTTCTTGAA CAATTGCTCTT ATGTCACTAA ATACAAGAAA
 -2892 GGGAGCACCA ATATAGTGGC CGATGCTCTT TCTAGACGGC ACACCTCTCT
 -2842 TTCAAAACTA GGTGCCCAAA TTCTTGGATT TGACACATA AGAGAGCTTT
 -2792 ATCAAGAAGA TCAAGAACTC TCATCCATCT ATGCCCAATG TCTAGATAGA
 -2742 GCACAAGGAG GTTACTATGT GTCCGAGGGA TATCTTTTAA AAGAAGGAAA
 -2692 ACCTTTCGATT CCCCAAGGAA CACATAGAAA ACTCCTGTGC AAAAATCAAC
 -2642 ATGAAGGGGG ACTCATGGGC CATTTTGAGG TTGATAAAAC TCTAGACTTT
 -2592 TAAAAGCAAA ATTTTGTGG CCACACATGA GGAAGAGGT CCACGACATT
 -2542 CTCTAGAGTA TCTCATGTTT AAAAGCAAAG TCTAGAACAA TGCCGCTGGA
 -2492 CTCTACACCC CTTTGCCGAT TGCAAGCTC CTGTGTAAGA CATTTAGCAT
 -2442 GATTTCATT TAGGACTTCC TAGGACTGCA AGAGGCCATG ACTCTATCTT
 -2392 TGTGGTAGTG GACCGTTTTA GCAAAATGTC TCACTTTAT CCATGCGACA
 -2342 AAGTAGATGA TGCTCAAAAT ATTTCTAAAC TCTCTTTAT AGAGGCTGAG
 -2292 AGACTCCATG GTCTCCCTAG AAGTATAGTG TCCGATAGAG ATCACCTTAA
 -2242 ATATATAATT ATACACTGT TTTTTTCTC TTTTTTATT TATCAAGTAA
 -2192 AAAGTATTG TTCTAGATTA TTATGAGTAT ATACTTACTT TCTGTATTTT
 -2142 ATTTCTTTCT ATTTTTFATG ACGATGAAAT TTCTTATTAT ATCCAGACTT
 -2092 TTCAATATA TTTTATTTC TTTTCCATCT AGATGCTCTG TACTTTTCTT
 -2042 CAGTGTAAAT TTCCACTCTC CAACAACAAA TCATTCAGT TTTGTATAAC
 -1992 ACTGTGACGT TAACCAGTTA AAATAAGAAA ATCATGTAAAT ATAAATTAAT
 -1942 TCAATAGATA TTTTGAART ACAATACGA TAAATAATAA AATTTAAAAA
 -1892 ATTATTAAC AATGAATTTT TTTGGAART AATATAAAC TTAGACTGTG
 -1842 GGTTCCTTCA TTCAGTCAAA ACCTTTTCTT ATGTGTGGTG GTGTGCGTGA
 -1792 ACATCGAATT TGGGTGCTT ATGCGCGTTT ATCTTCATCT GCACCTTCAA
 -1742 ATTAATAATT TAATTCCGGA AAATAATAAA CCCACACACT GTTTTATGCA
 -1692 TATATTAAGA TAAATAAAG AGAECTATTT TAAAGAAAT AAATAAATAT
 -1642 ATGTACAAA TGATGTCACT AAAGAAGAAA AAAATTAAAC AGAATTGTAA
 -1592 TATATTCTT TATGAAATGT TTTGTGCAAT ACCGAGAGAG GTCGAAGTGA
 -1542 ATACACGCAA GCATCTAACT AGTTTGGTAA TTCCTTTTCA ACATCGnTAA
 -1492 GCACATCACA CTAATAATT TTTAAATAGA TAAATTAGAT TCAATTGTGA
 -1442 GACATTAAAT TATAATACTC TATCCAAAT TATACTATA AATAAAAGT
 -1392 TATTTTAGA AAATAAGTAA TGAAAATTTA ATTTCAAAAT TTATAACCT
 -1342 TTTATGCTGT GTTTGTTTCG AAGCATAGAA AAATAAAAG TTATTGTTGG

-1292 GAATGAAAAG TGAAGAAAAT CATGTAATAA AAACAAAATG ACACGACAAT
 -1242 CAAAAAATAA GTTTTCATGC AAAAATTTT TCAAAATTTA CACTTTTATG
 -1192 ATGTGTTTGT TTCGAAGTGT AGAAAAACGA AAAGTTATTA TTGGTAATGA
 -1142 AAAGCGAAGA AAATCACGTA ATAAAAACAA AGCAAGATGG CACGACAATC
 -1092 AAAAAAAGT TTCTACACAA AACTTTTATC AAAATTTACA ACATTTTAT
 -1042 GTTGTGTTT GTTTCGGAGG TATAGAAAAA CAAAGAATTA GTGTGGTAA
 -992 TGAAAAGTGA AGAAAACCAT GTAATGAAAA CAAAATGGCA CGACAATCAA
 -942 AAAAAAGTTT CACGCAAAAT TTTCTTCAAA ATTTATAACA TTTTCATGTT
 -892 GTGTTTGTGTT CAAAGCCTAG AAAACGCAAG AGTTACTATT GGTAAATGAA
 -842 AGCGAAGAAA ACCACATAAT AAAAAACAAA TGGCACGACA ATCAAGAAAA
 -792 AGTTTTTACA CAAAACCTTTT TTTCAAAATTT ACTATGTTTA TTTCGAAAT
 -742 TAGAAAAACG AAGAGTTATT ATTAGTAATG AAAAGCGAAG AAAACTACGT
 -692 AATAAAAAAC AAAATGGCAC GACAATAAAA AAAGTTTCA CGCAAAATTT
 -642 TCTTGGTGCG CAGAAAGTTA TATATATTAA TTAATTAATT TTCATTACT
 -592 TTTTCCCTTT TTTATTTTAA AGTTAAATTA TTATTATTTT CATTTAAAT
 -542 ATAAATATTA TTTAAATATA AAAAAATATA CCTTAATCAA AACAAAGCCT
 -492 TAACTAAAAA TTTACAACAC TTTTAACTTT AAAATTAAC TTTAAAGGAA
 -442 AATGATAGTG TGACAACATA AAAAGTTGTA TACAACCTTG TCATAGTGT
 -392 AGAAATAAAT ATATATAATA AAGAGTAAT TGTATATTA ATGATATAAA
 -342 AAAGTATTAA AATAATAATA TTTAGAGTAG TAATATGGTT GTATAAAAAA
 -292 ATGTGGTGTG CCAATATATCA TTATTCACTT TAAATATCAA TGCAAAATAT
 -242 TTTCAACGAA AGATGGAAGG AACGAAAAGA GCGTTGGATA ATGGAAAAAT
 -192 ACAAGCAATC TCCCTCCAGT ACTTTGCATA ACATTTTGTA TTAGTATGA
 -142 GTTTTTTATC ATATATATTT AGAATATAGG AAAATTTTAG AATCACGTGG
 -92 ATAGCTATAT AATAGTAATA TTTTAATTAA TAATGTAGTT GATTTTATTT
 -42 GTCAACTGGT ATACATAAAT ATGTGTGTAT AGTGGGTGAC TTGTGGCTTA
 9 AAGAAATGTC CAGAGGCTGA CAACAACCTC GCACAGACTA GCGTAAC
 57 **ATG** AAG TCC AAT TTT GCT ATT TTC GTA GCT TTT TCT CTT CTT CTT
 1 M K S N F A I F V V F S L L L
 102 CTG GTACCTCTTCAATCTCTCTACAAAACCTCTGTGTCTCTTTCACCTCTGTTTGTA
 16 L
 160 ATTTTGTTTTACACTTTTGGAAAAATTGAAGCTGATATATATGTAACAACCTTTTCACTTTT
 219 GTCTGCACTGAAACTGATAGAAAAATATACGTTTTGTGGATATATATAG GTT GGC
 17 V G
 35 274 AGT TGC AGC TGC GCA AGA AAA GAC ATG AGA GGG TAT TGG AAG GAT
 19 S C S C A R K D M R G Y W K D
 319 ATG ATG AAG GAG CAA CCT ATG CCA GAA GCA ATC AAA GAC CTT ATT
 34 M M K E Q P M P E A I K D L I
 40 GAG GAT TCA GAA GAA GTG TCA GAA GCA GGG AAG GGT CGT TTT GTT
 49 E D S E E V S E A G K G R F V
 AGG GAC TTC GAT GTA AAG CCT AAT GTC ATA TTA TAT CAC ACA CAT
 64 R D F D V K P N V I L Y H T H
 GTT GTG CCC ATG AAG CAG AGG CAG AAG AAT AAA GAT TGA
 79 V V P M K Q R Q K N K D •
 45 493 AGACTATGTGATTGGCAGTTTCAGACTTATTTGGCACCAAAATTTATGATGCTCTTGTGTGCTG
 555 TTTCAAATTTGTACTCAAACCTTTGACCCCTTGCAGCATCTTGTCTCTTTTGGTCTTGTCT
 617 GAATTTTGTACAGTTATATCTGTACGAATAGTTTCTCTCATATAAGCAACATTTTCTCT
 679 C

100%
 90%
 80%
 70%
 60%
 50%
 40%
 30%
 20%
 10%
 0%
 25
 30
 35

FIGURE 3

101001 CAAAACAAAAGCAAATGCCGGTTTTCTTATTATTATTTCGAACCTTTAGAC
5 100151 CTTTTGTAACTGTTCTTTAATTTTTTCTTGTATGATAAAGAACCCCTATTAT
100201 ATCTTAGCTAAATATTACCTCATTTTGTATTAGCTAAACCAACCCCAA
100251 AAATATTGTAGTTTGTCTTCGGATTAACTGCCAAGCAAGTGATTAGAT
100301 ATATTAAAGGAAAATGAATGAAGGACAAAAAATATAACGCACAATAT
100351 TGAATACTGATATTATCTCCATTCTCAAAATTTTTGTGTTATTGTGAC
10 100401 AATATTGGTTGTTTCCCATTTGCTACATCTTTGAGGACATGAATGATA
100451 ACATATATATGAACGAGTATAATACATTTCTCGTTTCATTTTCAAAATAAT
100501 GTCAATTTATGCTAAACATTTTTATTAAAAAATTATCCTTATAAGATTTC
100551 AGTGATTATTATTTTACCATTGGTACTGTAAAGTCGGATGCTATATATATATA
100601 TATATATATATATATCAAAAATGACACTGAAGAATTATTTTGAACTAAAA
15 100651 CTAAAAACGTAAAAATAAAGAAATTTTTCAAAAATCAAAAATTTTATATA
100701 AAAATATAGATAAAATGTTAATATAGTACAACCTTCTATTCAAACAGAGAG
100751 AATAAATCTTCTATAGACAGTGAATATCCATTATAATAACGAGCAATAGT
100801 TGTAAATGTTGCAGTACAAAAAGAGAATTGTAATATTGTGCATGATTGAG
100851 AAATCTAAGTTGACTTTGAATTTAAAGGCTAATTCACAAAGTACATGTA
20 100901 GAAGTTGACTATAGCTATATATTACTACAAATTGATCATTTTCAAGAAAG
100951 ACATTTAAATTAAGATATGCTATGCTAGCTGATTGTAAGAACCCCATCTCGCTT
101001 GCTTCGTGCCATTGACACAGATGTTACTTTTAAATGCAAGGTAAATATG
101051 GATATACTCTTCGTATTTTTTGTAGTAGATATTTTACGAAAAATTTGTT
101101 TTTTCCAAAATCAAAATGATATTATTAAATTTCAATATAGAAATTAATTA
25 101151 AATTTTAATTAATTTGAAGATTATATGCTGCAGATTAGATTACCATG
101201 GTGAAATCATGTTTAGGTAAATAATAATGATGTTGTGATTTAGGAAAAA
101251 AAAAAATCTTTAATCTTTATGTAGAAGATTAAACCTCAATTTATAAAAA
101301 TATGAAGCAGTATTATATAAGATGTTTAACTAATCGAAATAATTTTTTG
101351 GGATGAAATTTTCTGTCATATGTTTCTAAAAAATAATATGTGAAAAATT
30 101401 AACATTCATTGTATGTTTATAAGAAATATATGTGAGTTTGTGTTAGATAA
101451 ATAATACTTAAAAATTAAGAAATTTGAAAGTTATACTGCACTTCAAAATATG
101501 TTATTTTTCTCTTTTATTAAAAATATCAGCAACATTTCAAAATGATTTTAT
101551 TTTCTTTAAAAAATGAAAAAATGAAGTTAGCAAAATATGTAAATTTTAAA
101601 ACGAATTTAAGAAAAAACTTTGTAAGATATGATATGCTTTATAAAAAAA
35 101651 ACTGGTGGCGTACCTACTAAATATGATCACATTAGAGATTGTATCCCT
101701 TAGCATATAGTATGTAGTATAGATATCTATATTTTTATTATTAAGAGAGC
101751 ATATTCAATAATAGGTATTATATGTAAATACAAATAACGTTCAATTCG
101801 TTATGTTAGTTTTTAGAAAACTTATTGCGGTGTCATATCAATGTAGTAA
101851 GCGACTCCACATGTGAGATGTTGGTCTGAGAAAGCTTCTGCACTTGCTC
40 101901 GGAACACTCTCATGGACTAGAATGCCATCCATCTATTCAAAGAAAAAGCAG
101951 TTGTCATGTCATGCTCGGTTTTTTCACATTTGGAAGCAGCGCAACAATGT
102001 CTTACATAATATGCGATCGATCACTCTGCAACCAATATTCAAGTACATAG
102051 ACCATGACATCAAAAACATTATCACACCGGAGAGAAAGAAACGTCAAATT
102101 GGTAACCTTAATGGCGTTATGCTGCGGTGAATCTCCTAAGAGTTCTCC
45 102151 AAATTTTATTGATTCTTGTGTTTTAACTTTTTGCGCAAAGAAATACATAT
102201 ATAGATTGACACCAATTTCAACTTATCAAAATACAAGTGAATAAATAATT
102251 CAAGCTGAAAGGAATTTAATCATGATCTAAACCTAAACGCACAAATCTT
102301 CACAAGTGAGAATCACTAATGACTACCCTTGGTGCATATACATCATT
102351 GTTGTAATCTGAAAAATGGTTGGGATTGATCTGATATGTCATTATAT
50 102401 AAACTGTGATTATTATTATTAGAATTTTGCCGCAACAGATAAATCATC
102451 ATCTATTAGAAAAATTTTCATTTGCAACCAAAATTAATCAGGGGAAAAAGGT
102501 GAAATCACATATCTTATCTACACTCTTATTATTAAATTAACGCCATATATA
102551 ACAAAATTTTCAATACCACTTATGAGAAGCACTAAGATCACCTTTTCTTCT
102601 TATGACTTTCTTTCAAAGCTAAGCTGGTAGTCATGACTCATGATTATCC
55 102651 TTTTCTAATGGGAATATTGTGAAGCGGTTTCAAAATCTTTAGACAAAAAT
102701 TCCATGGCCACTAAAGTTAGCAAAAGTTAAATAAGTTTAAAAAATAATG
102751 AGTGTACTTGGCCATATGCCATATTGTTGAGATCATAAAGAGAAATAA
102801 TAGTTTATTGAAGTTAGATCATAAATCAAAATACATCATTTGCTTCATCA
102851 ACATTTTCCATGGATTGAGAGGATCAACTCAATACTAATGGTGGGGTC

102901 TTATTCATCCATTGCTCTCTAGCCAATTAAAGCAGTTAGGTTATTTGTGTA
 102951 CTCTAGTAGTTGCCAAATCAATCTTAAATATCCAAATGTTGTAATTTCTA
 103001 ATTACGTATAGATAAAATGACTAGATAACACGTGGCTTTGGTTTATCAGG
 5 103051 AAAGTTTCCCAATCATATATATGAATGTAGAATAGTGTCTTCATTAAAT
 103101 TATTAAATAGCATCTCACCATCTGAGACTGGGAGCATGTGACAAGTTGAC
 103151 ATGTGTATTAAAGAACTTTTGAGAAAACCACTTTTATGATACATCCCATCT
 103201 GAGACTGGGATGAGTACCATTTTATAAAAAATATGAGTAGTGAAAAAATAT
 103251 TCAAAAAAAATTTCTACATGTCTTTTAAACATTTTAACTTTATAATTTT
 10 103301 AACAAACATCTTCCAATATGCGTTATGAAAACTTTATAAACTTTTTTAT
 103351 AACATGCTTTTGAAAAATTTATAAACTCTGTATTTTATAGAAAACAAAGTGAT
 103401 ACTTTTGAAAATAGACAAATGAAGTGCTATTTTAAAAATTGATATCATATA
 103451 AGTCTTAACTGTGGTTTGTTTGAAATTTATTTATATATCTTGTCAAAATAA
 103501 AACTAAATAAATAAATAAATAATTTATTTATAATCATGAAGATAAATATTATC
 103551 ATAAAGATAAAATATAAAATCAACAAATTTATATTTGTTAAATAAAATAC
 15 103601 TTTGAGCTCTCTCTCATAGACTTTTCCAGCTCCCATCTAGAAAAACACA
 103651 TAAATTAAGATAAAATAACCGAATAACATAGTTCACATCTTAACTCTT
 103701 AGTCTTAGATTGTGTTTAAATTTTCAAAGGTTTAGGTATTTGATATATGTTT
 103751 TTTTATTGGGTTGCTAGATTTTGATCCAAGAAGAAATGACGGGTTGTAGT
 20 103801 ATAGATGTTTGTGTTGAGTTTTCGCCCTTGGTTTACTCTGTTTGGTTT
 103851 TGTCCCCAGAATTGTTCTTGATCTGCTGGTTTATGTTCTCTACAAAGTGCC
 103901 ACGACCATTTGCCGGCTCTTTGTATTTCAACTTGAATTTCTAAATTCGATG
 103951 ATGAAAAAAATGTATCTCTTAAAGTCCATTAGTACCAAAAAATAACTAT
 104001 ATCATTACTACATAAAATAGCTTTGGGTTTCCAAAGTATTTCGTGTGATA
 25 104051 TATGTTAAGAGTTTCGAAATAGACACATAGATATAATGTTGAAATGGGACC
 104101 TCTCACATAAATATCTCCTTTCTCTTTCATTTCTTACCTCTCAAGTTTC
 104151 CAATCCCACCTTAAGGTAATTTATTTCTTAACTTAAGTAATTTGTTAAC
 104201 AAATCTTAACTAGCTACAAATGTGATTACAAAGCTTAAATAAACCTTA
 104251 CTTTAAATCAAAGGTATTAACCTTCTTAAATTTGATACTTACTTAGTATC
 104301 GATCGGTCTAGTTTAGGGTTTGGACACACACCATCATGGGAGCGAAAT
 30 104351 AGTCATTCTACGGTGTTCCAAGACACAAATCTCGGACTCGATGTGGATATG
 104401 ACACCTTCATTATAAATTTTAACTTCATAAAAACTAACTATTAGAGGAAG
 104451 AATCGGAATCTGCATATCAATCACAATAGACTATAGTATACTTAGATTTT
 104501 GATCTAAATCAATGGGCTCCTTCAACTAAATAGTAGCCCACTACCAATTAAT
 104551 GAAATCATAAGACATTATTAATAATCAATGTTCTAAAAATACGTTTGGT
 35 104601 TATGTGTCCCGTAGAGCTAAATGTGCACACAAATGAAAGTTGACCCGTT
 104651 CACTGTGCCCACTTTTATGATCTTTCTTTTAGGTAAATCCAACCTTTA
 104701 TAATCTCATCTGTTATCAAAACAAACCTTTTGCGCTGTCTTTTCTAATAT
 104751 TTAAGTAACCTCTCACGGAGAAAAGCCAACTTTTCTCTGTTTATCT
 40 104801 TTTTAAAGAAAAATGAATTCAGGGGACCCAAATTTAAAGGAAAAACCA
 104851 AAACCTCTTCTATGTATTTTAACTTGAAGTTTCTATGTAACTAACAA
 104901 TCTTAACAGTAGAGAAATAAAAAACATCGTTTGGGAGGTTTATATGATG
 104951 ATATGAGAAATGTTCTAAAAATGTTTACACAAAAATTAGATTTTCTTTT
 105001 CCTCTGTCAATGGAGCTATATCACTGTCTATTGTCTTAACCCCTTGGCG
 105051 GAAGATTGTTATGAACAGTTTAAATGGAATTTCTAGTGCCAAATGTCAAG
 45 105101 TTTAATATGTTTGTGCCCTATACTTTATGAACTCTATAATCTTTGTTAT
 105151 AGAATTATCTACTTTTAGTATTTTACATTAACATAATCTATAGAATTTCT
 105201 CTTTGTCTATACAATTAACCAAGTAATATATCTTAAATACATATAAAA
 105251 ATGGTGGTGTGCTATCTGAGCTGTAATAGTTGATTGCTCCAGAGAAGAA
 105301 TAGACAAAAATCCTTACTTAAGAGGCCCACTCTGAAAAATTTAGACAA
 50 105351 GAAAAATTAACACAAATTAGGTTACACATATTATCATTTTATATATATGCA
 105401 CAACACAAAGTTGACCTTGCATGTACTATTGAATAAAATAAATAAATG
 105451 AAGAAGAGAGGGAATTAATCACTGTTTACCAAGAAAACAACTCTCTCTAAAC
 105501 AGGCTCTATATATATAAACTTTAACACCTTAAGAAATTAACACAGATCAA
 55 105551 GTAAAAATCTCAAAACAAAAGTTAAAGCAGAC ATG AAG CAA CAG CAA
 1 CGT TAC TTG GTC GTC TTC ATC GTC CTT TTA AGC TTT CTT
 6 R Y L V V F I V L L S F L L
 105638 CTG GTAAAGCTTCTCTCTTAATTTATTTAAAGCCCTAATTAAGATCTCATATA
 19 L
 60 105691 TCTGAATGTTGTATATATTGTTGGTATAG TTT GTG AAT CTG AGT
 20 F V N L S

105736 GAA GGA AGA ACA GGA GGA GTT GCA GAA GAA TAT TGG AAG
 25 E G R T G G V A E E Y W K
 105775 AAG ATG ATG AAG AAT GAA CCG TTG CCT GAA CCA ATC AAA
 38 K M M K N E P L P E P I K
 5 105814 GAG CTT CTC AAC AAT CCT TTT AGG ACC GCA CAA GAG AGA
 51 E L L N N P F R T A Q E R
 105853 TTC ATC CAG AAT TTC GAC ACC AAA TCT GTT GTC ATC ATC
 64 F I Q N F D T K S V V I I
 105892 TAC CAC AAT CCT AAT GAA TAA TCAATGAAGTCTCTCATATAG
 77 Y H N P N E •
 10 105934 ATATCTATGACTTTAATTGTTGTTTATGTATGGATCGACTTATACGTGCA
 105984 CGTATATGTTATTAAATTAAGAAAAAGAAAAGCTGCTTGAGTTGTTGTGTT
 106034 ATACACGTATACTAAATATGTTCTGTTTAGTGCGAGAAATGTTAACCCCTAG
 106094 CTATAAGGGATTTTTTGTTCCTTTTTTTTGTGTACCAATTAATGTGAGTGA
 15 106144 GTGAGTTTTGTGTGATGAAAATAGATTTTGCTTACATTGTTGTTTGTGATA
 106194 TATATAAATCAATATACTGTGCCTTTTCGTGCTCTGTTTCTTATATTATTT
 106244 TGTGACATTAAATTAATTATCTTATCAAAAATTTATTTTATTAACTGTGTCT
 106294 CTATGAAAAAGATGAACAATATGAGTTAACCTCATCTCAAGGAGATTCT
 106344 TTTTGTGTTTGTGTTTTC

20

105736
 25
 105775
 38
 5
 51
 105853
 64
 105892
 77
 10
 105934
 105984
 106034
 106094
 15
 106144
 106194
 106244
 106294
 106344

FIGURE 4

1 AGCCTTTACAAATGTCTCCCAAGAAGATGAACCACGTTATTATTAGTAATCCTGAAAAGG
5 61 TTAACTGCTCTTGCTCTCGAATTTCTAAACCATCTGAAATATCTTAGTGTTTAAAATGGAG
121 ACTCTGAGGATATAGTCTTGCTGAACACGCTGACCGCTGAGTATAGATAACATCTAGTAAT
181 TTCTACCGGAGCGGTGTAAGAACTAGTCTGCCCTCTTAAAGAAAGCGCTACACCGGGCACC
241 TTCTTTTACGAATTCGAAATTCAGTCTTGCGGACGAGCTGTTGGATCGAAGGCTCTTTTCT
301 GATACCTGAGGAAATCTGACTTTCGAAGGAATAATCTCTAATCACCCACCACCCCTCGA
361 ATACACTTCAGGACCTCTGAAACCAACTCGTTTCGCGTAAATCACAAGAACTCTCCAC
10 421 TCATTCCGATTTTAGCCAAATTAATATGATATCGGTCTGGGAAGCCGATTAAGGAAATCTCC
481 ACAAAAGAGGATTTATGAATGAGGAAATAAGGAAAGAGAGAAGTAATTTTTAGGTAGTACCC
541 TGAAGAGACAGCGAAAAATTTAGAAAAAATACTACTCTCTGACATGCTTCAAA
601 GAATATCTCnnnnnAATGGTTAGTAATTAAGAAAGAGGATAGTATGATTTAAATCGAAAC
661 CACGTCCGGCAGAAACAAAGTGAATATCCCCCTCTTGTAGGAAGTTCGTTTCTTAAATATAG
15 721 AAAACAAAGAGTGTGCGCTCCCTTTTAAAATGATCTCAGAAAACGAGAAAGTATAGTAT
781 TAAAGATATCTCAAATCTACACAGTCACTAAATTTTACTAATCAAATGTTTCTTCAAT
841 TAGCAAATAAAAACAAAGAAAAAGmGAAATTTGAAGAAAATCGTTATATAAACACAT
901 TTAAATTTCTGAAAGAGAGATAAAACAGTTAAAAATTTGGTCAAATCTTGGATCTTTAT
961 ACGATGAAGTAAAAACAAAGTTATCTATCTCGGTAAATAAGAAAGAAACCGAAATCTAATA
20 1021 ATGATATGCTGCCAAAGGACAAATCCAAATCAGAAGATTTTAGACACAAATCGAAAAGAGAT
1081 TAAACCAAACTCTGAACGACCAATTAATGTGTAGAGAAGAACTAAACAACTCAGACTCAT
1141 GTTCAGAGTCTCCCGACGTGATTTAAAAAATAAAACGTAAATAACACAGAGATTTCCAGTA
1201 AGCCGAGATTTTACACTGCGCCCACTTCGATATCTCAGATGAGGCGATGGACCAACT
1261 GCATTTGTAGGAATGGATATCTGAGTATGATAGTATGATGACAGATGATGAATATAGAAAAAC
25 1321 TACGAAATTAACAATGGCCGCTCAGCAGTATAGAGTAAACATATCCAGGAAACAAACAT
1381 AAAATTAATATATCTGGATTACCGGAGTATTAAGAGGCTGGTGGGATAATTAACCTCAT
1441 CGCAGAAAGAAAGAAATATTTCTTAACTGTGTAAGAAATAGAAAACGAGAAAGGAATACC
1501 ACTAATGCTGGAAACATTGGTGGTAGCATAATTTCAATCACTTTAGAGAGATCCAAAGAT
1561 TTTTGAAGAAAGACCACTTTTATTCTTCATATCTAAGATGTCCCAACTGTTAGGTGACTT
30 1621 TAGATGGTATTCAGAAAAATTTTTAGCTATGGTTTAAACAGGCGAAGATCTTAGAGAAC
1681 TTTCTGGAAGAAACGGTTTATATAGCTGGAATACCGGATATCTTGTGGAAGAGTAAAGAA
1741 AATTTTCAAAAAGAGGTGCCAAACCCCAATTAAGAAGATGCACCATCGGGAATAAG
1801 TCGATTTGTAAAAAATCAGAGTCTTCAGTATATGCAATAATAGAAAATAGAAAATAGAT
35 1861 AAAAAGAGGTGAGAGTCAGGCGCTCAGGAATCAGGAAATTTGGGGAATTTTGTACTCAATCGGTA
1921 TGAACGAAATACCCCTCCATCAAAAAATAAAAAGAAATATAGCAAAAAGGAACAGGAGG
1981 AAACAGCGCGTAAAAACAGCGCTAAAAACGACGCTAAAAATTTAGAAAACCGGTATTT
2041 TTAGAAAACCATGAAGGTCTAATGATAGGCCACTTAGTCTGTTTATAAATGTGGACGCA
2101 TAGGACACATGAAGCGAGAGCTGAGACTATGAAGAAAAAATAGTAATTTAGACATATAGT
40 2161 ATGAAATTAAGAAAGAACTTGAAGAACTCTGATGAATTTCTCCAGAAGAGGAAGAAC
2221 GAGAATATAGTAGAGATCTGATTCAGAAAGTTTGACATAGGATCAACAATTTGAATTT
2281 GTGCTATAAAATATAACAGTATAGTAGTAATTAAGATTTGCGTTAGATGCAATGATA
2341 AAATTAATATATCCGGAGGAAACCAAGGCTTAATAGACATGAAAGGCTACTCGTGT
2401 AAAAAGTAGAACCCAGTTTCACTTCCACAAAACCTGAAATTTATAGGATATGATTTTAAAG
2461 AAATATTTGAGAAAGCGAAACATCCATCAAAAAGAAATACCAATTTAGCTATCTTAATAGT
45 2521 AATAATAATAATTAAGGCGGAATTCGAATCTATAAAGCTCGAGCTACAGAAGATTTAAAG
2581 ATAAATTTATACATGAGGAATCCATCTCTCTCGCAGCGAAATTCACAAGAGAGGAAG
2641 CTAGTAGACCTTCATCAAGAAATAATACATCAAAAGACAAAGTGGCATGTAAAAATAGT
2701 CCGTAGAATTTGTTTGTGTTTGTAGCCGTTTCATGTGTGTCACAAGATGAGTCTTACCTTA
50 2761 CACAAATAAAACGTTACTCTTAATATCAAGGAGAGCTACAAATCTCAATGAATGAAT
2821 GACATTAATATTTTCTTTAGTTTATAAATCTGAATGAGTTTCTTTCATAAATCTGAC
2881 TGACTGACATTTTATTTTCTGAAATGAGGAAGGTTTATTAGCTTTAACACCATATAT
2941 ATATTTTATCTCAAAGTCAAGAAATATTAATAAGGAATCAATTAATAAAATATTTCT
3001 TTTGCAGAAAAAAATTAATAAATATGAAGCTCTCCACCATATTACATATTATATAA
3061 TATAAAAAAACCTCTCACAAATGTGCATCTCGGAATCTTTATGTTAGAGATTTAATCTC
55 3121 TAAAGAAAAAGGTTGAGAAAGGTGCAGCAACA **ATG** TCT CCA TCT TGT AAT
1 M S P F C R
3172 AAC TTT TCA ATG GCA TGG GTG CTT ATG GCA TTT GTG TGT TT
7 N F S M A W L M A F V L F
3214 GCA AAC AGT GCT ATG CCC ACA AAT GGA TTC ACT GTT GGG GTA

5

10

15

20

25

30

35

21 A N S A M P T N G S T V G V
3256 AAA AAC ATG TTG GGT GGT AAA TTG ATG CTA AAC GTT TTA TGT
35 K N M L G G K L M L N V L C
3298 CCC CAT ATT GAT AAG CAA CAC ATT ATC CCG AAT GTT GGT TCA
49 P H I D K Q H I I P N G G S
3340 TTT GAG TGG AAG TAC AAT GGT GGT GCT CCA CCA ATA GGA CAA
63 F E W K Y N G G A P P I G Q
3382 TCA CCA TTC ATG TGT TTC TTT CGG TGG AAT AAT GTT CAT CAC
77 S P F M C F F R W N N V H H
3424 TCC CTT GAT CTG TGT TCA CCA AGC AAG TAT ACT GGT TGT GAA
91 S L D L C S P S K Y T G C E
3466 AAT GCC ATT TGG GAA ATC AAA GAA AAG CAA TTT TGT AGG TAC
105 N A I W E I K E K Q F C R Y
3508 AGA GGT GGA CCT ATT AAT TAT TTT TGC TAT GAC TGG GAT GAT
119 R G G P I N Y F C Y D W D D
3550 TAG TTATATAGATTATTCATGTTTCATCTCAATAAAAAAATGACTTTAGAGTGATTCTT
3609 AGTTTGCTTAAACATTCTTACATATTCTCAACTATTCGGTCACTACCACCCGTAACTATAT
3669 TTATTTAAAAATTAGTATCTGTACAGTTTTATTTTTAAAAAGGTTATGTGGATTAGAAG
3729 AGAGATAAAATGTTAGACGGTCACCAACCTTAATTTTTGAACATATGAAGACTATATTGA
3789 CCAAGAATATATGTTTAAACTCATTCAATTAAAGACTATATCTCCATTTATGATTATGCA
3849 AATGCAATTAGTTTTTTTTTTTCATTGAAGAATTCAAAAGAAAGTTATCATTAAAAAGTAT
3909 CATTAATCACTTATATGTTGTTTCTTAATATCCTTATTGTTAATAGAATAATTTTTTTT
3969 ATCCTTTAATTAAGGTTATTACTACTTTTTTTTCATATCTTCATTTATTTTGAAATATTTT
4029 TAAAATTATCAATTTTTGTACACCCAGAAAATACATGTAACATATCACTTTTTTTTTT
4089 TATTACAAATTTATGACTTATAGAAATACAAATATTAAAAATATAAGGTTCAAAACTACA
4149 TCCTAAGTCTTTCAGACCCCTGCACACATGTATCATCTGCTCGTATATGTGATACAGTC
4209 ATCCGAGTTACAAGATAACAAGAAAACCAAGGGTAAGCTAATGAAAAAAATTCATAA
4269 CATATTTAATTCATGCAAAAAGAACCAAGTCAAGTAATCATTATAAACATTTCTTTAAA
4329 TATTGTTATATAAAATTTCAATATCAATTTTCATCATTCATATAGACCACACATGGATCTA
4389 TTTTCAATCACAAATCATGGATTTCATTTTAATCCTACTTCGnCTTCCAGAAAGCTCAT
4449 AAGTATGCCCTACCAAGAGACTAACACCTAATCAAAGAGAAATGATCAAGGTAAAGTCAA
4509 ACATCCAATAACGAGTGCCCTACAGTGGGACCCAATGTGTATGAACTCCTTATCAGCTTCT
4569 CACCACCTGATATCTTATCTATATGACGTAGATCATCAGTGAACCTAGAGGATCTCCGT
4629 TAAACATATGTTTTTTTATACTTAATGTCTACAAACAACAACCTCACACATTATCCCAAATG
4689 TATGACATCAATTCATACAAATTTTCATCATTCATATATAATACATATCATTGAATCACA
4749 TAACATTTAAAAATTCATACCATTCAAGAACCTTTCCAACTCAAAAGCAATATTACTT
4809 TCAAACATCAAAATATAATTATTATTATTAATAAGCTT

FIGURE 5

142000 TTATCTTATTCCATATAATGTTGTTTTACTTTCAAAATTTTAAATTTT
 141950 TTATATTTATCTTTTTACAGTTTAAATTAATAAAATGAAATTTTTTTC
 141900 TTAATGTGTTAAATATATAAAATCAAAAAGTTGTTATATGGTACATGGC
 5 141850 ACAATCTTTATAAATTAATTAATTTGAAAACGATCTTTATATAATAAAAT
 141800 ATCTTAGTTGACATTTTATTAGTGTTTTCAATCATATTTTGTGCTT
 141750 GATAAGCGTAAAACAAACTAACTTAAAGTATCTTTATATAATAAAATTA
 141700 TCTTAGTTGACATTTTATTAGTGTCTTCAATCATATCTTTGTGCTTG
 141650 ATAAGCGTAAAACAAACTAACTTAAAGTTGGGCACCTCAATGTTTTAAAA
 141600 AAGTTTGGGTACCTCAAAAATTAATAGGCTTTGTCAGATCTTACAAAAA
 141550 AAATCTGGAAGAATTATGAAAGAAGGGGGGGAGGGGGGGGGGGGGGG
 141500 AAGTGAAGATGAATATTCAACAAAAGAGGGGTAGGATGATGTTAAGTGAG
 141450 TTAATAAACTATGTTAATGGAGACAATTTCTGTTAAACAAACCCGTTAAT
 141400 TGAAAAACGATAGCATTTCTTCTTAACAATGTAAAAACGATATTGTTTATC
 141350 ATAACACTCATTAATAATTTCTGAGTTTCAAAATCATATAAGATTAGGGG
 141300 GGTGTATTCAATTAAGGATTGAAATGATTTGTAATAAAATGACAAATCC
 141250 CATGTTATTTCAACATGAATTTGAAAACTTTTTTAAATCAAGTGTTA
 141200 TTAGATTAGTGATTTTAAATGTACAAACCAACCCACTGTTATTGGAAC
 141150 ATTTTAAGTAGTGGATTTAAATGACTTGAGTGATTTTGGGTGGGATTGC
 20 141100 AGAAAATTTCTAGTTAAGAATTCAAACATCCAAATCTCATGGTTTCAAG
 141050 TAGAATTTGGGAGAAATTTAATAACAAATCTCCTTAATTTACCAAAAGTCA
 141000 CCAAAATCATTTAAAACTCAATTAATAATTTAAATGATTTCAAACTCTCCAG
 140950 TTGAATACATCCCTTGGAAATTAGAGATTTTGCTCGATTGGGACCTAAG
 140900 ATTGAATTTGGGGATTAGTTTAATCGTTTACAAACAAATGACATCGTAT
 25 140850 TATTGTTATAGGAACAATTCGTTTTTCAGTTGACATGTATGTTAATAGA
 140800 AAATTAACCTCTATTAAACGGGATTGCTAACCCATTTAACATCGTAACATA
 140750 ATGGTCAAGTCAATAAAAGTTGGTATTATTGAAAGTCAACGTAAGT
 140700 TTGATATTATTGAAAAGTCAACATAAAATTGATATCTTATTGTTTTC
 140650 GACAGACATAAGGATTACATCAATGTTTTTAATAAATTAAGATTATTA
 30 140600 TGACATTTTTTCCATTTAAATTTGCCAATGTTTTCGAAACCAAGATACTC
 140550 AAAATTGACATACCTAATTCATCTACATTTGTTTGACAGCAATTCACGT
 140500 GCCTTGACCACATGGCACATCTGGCAATACATCAATTTTAAGGAAAAGG
 140450 TAGATTGCGATACATATAATGGAATTAAGTGGAAAAGGATCATTTGACTC
 140400 TTGACTTGTAAACAAACACACAGTATATAACTCAATTCGACATTTACAA
 140350 ACAACATTGTGCTAGCTTAAACTCCCTCTCCTATTCAAAAAA **ATG**
 1 M
 140305 GAT ATT CCA AAG CAA TAT CTA TCA CTA TTC ATA TTG
 2 D I P K Q Y L S L F I L
 140269 ATT ATC TTC ATA ACT ACA AAA TTA TCA CAA GCC GAC
 14 I I F I T T K L S Q A D
 40 140233 CAT AAA AAC GAC ATT CCA GTT CCC AAC GAT CCA TCA
 26 H K N D I P V P N D P S
 140197 TCA ACA AAT TCT GTG TTT CCT ACC TCG AAA AGA ACC
 38 S T N S V F P T S K R T
 45 140161 GTG GAA ATC AAT AAT GAT CTC GGT AAT CAG CTA AGC
 50 V E I N N D L G N Q L T
 140125 TTA CTG TAT CAT TGT AAA TCA AAA GAC GAT GAT TTA
 62 L L Y H C K S K D D L
 140089 GGT AAC CGG ACT CTG CAA CCA GGT GAG TCG TGG TCT
 50 74 G N R T L Q P G E S W S
 140053 TTT AGT TTC GGG CGT CAA TTC TTT GGA AGG ACG TTG
 86 F S F G R Q F F G R T L
 140017 TAT TTT TGT AGT TTT AGT TGG CCA AAT GAA TCG CAT
 98 Y F C S F S W P N E S H
 55 139981 TCG TTC GAT ATA TAT AAA GAC CAT CGA GAT AGC GGC
 110 S F D I Y K D H R D S G
 139945 GGT GAT AAC AAG TGC GAG AGC GAC AGG TGT GTG TGG
 122 G D N K C E S D R C V W
 139909 AAG ATA AGA AGA AAC GGA CCT TGT AGG TTT AAC GAT
 60 134 K I R R N G P C R F N D

139873 GAA ACG AAG CAG TTT GAT CTT TGT TAT CCT TGG AAT
 146 E T K Q F D L C Y P W N
 139837 AAA TCT TTG TAT TGA CAACAATATGCTGATGTTCTGTCTTTTAC
 158 K S L Y •
 5 139793 GACTCATGGAGTTTCATTGTTTGAAACAATAATATAAAACATATAAAATT
 139743 TCTATTATCCAAGTTCCTCACTTATAATAATTTGATAATCATATCATATT
 139693 ATCATCTTAAGCATTCAATGCTACAAAGATAATACCCCAAGCTATTTTA
 139643 CATTTAAAGCTGAAACAGAGACACAATACTAACGATAAAAGTTCGTAGTA
 139593 TCTTTATGCAACCATACATACATATACACAAAGATAGACAGGTAGTGTCC
 10 139543 TAATAATTCTACTTGGGTGAGGTATGAACAGCAGCAACAGTAGATACCAT
 139493 TGTATCCATACCACACATATTATGAGGCCCTCTGCAGATTTTGTAGTAAC
 139443 CATGCTCTCCCCACATCGCTCCCCACGAGTTCTTGATAATCCAA

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100

FIGURE 6





Figure 7

G564 promoter: Gain of function constructs



⊕ = Gus Activity in Suspensor

⊖ = No Gus Activity in Suspensor

▨ = Region Required for Gus Activity in the Suspensor

Figure 8

Web Signal Scan Program

Database searched: PLACE

URL: <http://www.dna.affrc.go.jp/hidocs/PLACE/>

This is the sequence you submitted

>G564 promoter (-921 to -662), 450 bases, 3D1A0BF4 checksum.

```
TGAAAAGTGAAGAAAACCATGTATGAACCAAAATGGCAGACAATCAA
AAAAAGTTTTCAGCAAAAATTTCTTCAAATTTATAACATTTTCATGTT
GTGTTTGTTTTCAAGGCTAGAAAAACGAAGATTACTATTGGTAAAGAAA
AGCGAAGAAAACCAACATAATAAAACAAAATGGCAGCAGACAATCAAGAAA
AGTTTTCACACAAAACCTTTTTTCAAATTTACTATGTTTATTTCGAAAT
TAGAAAAACGAAGAGTTATTATTAGTAATGAAAAGCGAAGAAAACACAGT
AATAAAAAACAAAATGGCAGCAGACAATAAAAAAGTTTTCAGCAAAAATTT
TCTTGGTGGCAGAAAAGTTATATATATTAATTAATTAATTTTCATTACT
TTTTTCCCTTTTATTTTAAAGTAAATTATTAATTTTCATTAAAA
```

Notation: H = A, C, or T

R = A or G

K = G or T

W = A or T

RESULTS OF YOUR SIGNAL SCAN SEARCH REQUEST

/tmp/signalseqdone.9437: 450 base pairs

Signal Database File:

Factor or Site Name		Loc.(Str.)	Signal Sequence	SITE #
-300ELEMENT	site	1 (+)	TGHAARK	S000122
2SSEEDPROTBANAP	site	101 (-)	CAAACAC	S000143
ACGTABOX	site	296 (+)	TACGTA	S000130
ACGTABOX	site	296 (-)	TACGTA	S000130
AP3SV40	site	159 (-)	TGTGGWWW	S000169
CAATBOX1	site	44 (+)	CAAT	S000028
CAATBOX1	site	189 (+)	CAAT	S000028
CAATBOX1	site	323 (+)	CAAT	S000028
CAATBOX1	site	138 (+)	CAAT	S000028
CANNNNAPA	site	101 (-)	CNAACAC	S000148
CCAATBOX1	site	138 (-)	CCAAT	S000030
CEREGLUBOX2PSLE	site	55 (-)	TGAAAACCT	S000033
CEREGLUBOX2PSLE	site	201 (-)	TGAAAACCT	S000033
CEREGLUBOX2PSLE	site	333 (-)	TGAAAACCT	S000033
DOFCOREZM	site	4 (+)	AAAG	S000265
DOFCOREZM	site	53 (+)	AAAG	S000265
DOFCOREZM	site	112 (+)	AAAG	S000265
DOFCOREZM	site	149 (+)	AAAG	S000265
DOFCOREZM	site	199 (+)	AAAG	S000265
DOFCOREZM	site	282 (+)	AAAG	S000265
DOFCOREZM	site	331 (+)	AAAG	S000265
DOFCOREZM	site	364 (+)	AAAG	S000265
DOFCOREZM	site	419 (+)	AAAG	S000265
DOFCOREZM	site	216 (-)	AAAG	S000265
DOFCOREZM	site	399 (-)	AAAG	S000265
DOFCOREZM	site	408 (-)	AAAG	S000265
GT1CONSENSUS	site	120 (+)	GRWAAW	S000198
GT1CONSENSUS	site	141 (+)	GRWAAW	S000198
GT1CONSENSUS	site	196 (+)	GRWAAW	S000198
GT1CONSENSUS	site	253 (+)	GRWAAW	S000198
GT1CONSENSUS	site	69 (-)	GRWAAW	S000198
GT1CONSENSUS	site	90 (-)	GRWAAW	S000198
GT1CONSENSUS	site	347 (-)	GRWAAW	S000198
GT1CONSENSUS	site	388 (-)	GRWAAW	S000198
GT1CONSENSUS	site	436 (-)	GRWAAW	S000198

GT1CONSENSUS	site	218 (-)	GRWAARW	S000198
GT1CONSENSUS	site	401 (-)	GRWAARW	S000198
GT1CONSENSUS	site	402 (-)	GRWAARW	S000198
MAMMALENHAN	site	158 (-)	GTGGTTTK	S000121
MARTBOX	site	324 (-)	TTWTWTTT	S000067
MRE1	site	356 (-)	TGCRNCNC	S000068
NTBBFLARROLB	site	418 (-)	ACTTTA	S000273
POLASIG1	site	168 (+)	AATAAA	S000080
POLASIG1	site	301 (+)	AATAAA	S000080
POLASIG1	site	324 (+)	AATAAA	S000080
POLASIG1	site	237 (-)	AATAAA	S000080
POLASIG1	site	411 (-)	AATAAA	S000080
POLASIG3	site	268 (-)	AATAAT	S000088
POLASIG3	site	427 (-)	AATAAT	S000088
POLASIG3	site	430 (-)	AATAAT	S000088
POLASIG3	site	433 (-)	AATAAT	S000088
POLLEN1LELAT52	site	11 (+)	AGAAA	S000245
POLLEN1LELAT52	site	119 (+)	AGAAA	S000245
POLLEN1LELAT52	site	156 (+)	AGAAA	S000245
POLLEN1LELAT52	site	195 (+)	AGAAA	S000245
POLLEN1LELAT52	site	252 (+)	AGAAA	S000245
POLLEN1LELAT52	site	289 (+)	AGAAA	S000245
POLLEN1LELAT52	site	362 (+)	AGAAA	S000245
POLLEN1LELAT52	site	71 (-)	AGAAA	S000245
POLLEN1LELAT52	site	349 (-)	AGAAA	S000245
PYRIMIDINEBOXHV	site	400 (+)	TTTTTTC	S000298
RAVIAAT	site	97 (-)	CAACA	S000314
ROOTMOTIFTAPOX1	site	374 (+)	ATATT	S000098
SEF4MOTIFGM7S	site	170 (-)	TTTTTTR	S000103
SF8SFIBSP8BIB	site	134 (+)	TACTATT	S000184
TATABOX2	site	81 (-)	TATAAAT	S000109
TATABOX3	site	375 (+)	TATTAAT	S000110
TATABOX4	site	368 (-)	TATATAA	S000111
TATABOX5	site	238 (+)	TTATTT	S000203
TATABOX5	site	412 (+)	TTATTT	S000203
TATABOX5	site	434 (+)	TTATTT	S000203

For more information about the SignalScan Program, please contact Dr Dan S. Prestidge Tele: (612) 625-3744 Advanced Biosciences Computing Center, E-mail: dandp@biosci.umn.edu 1479 Gortner Ave. University of Minnesota St. Paul, MN 55108 The TFD data is at the gopher site, gopher://genome-gopher.stanford.edu. For more information about the WebSignalScan service, please contact Meena Saktharkar, meena@biomed.nus.sg, BioInformatics centre, NUS .

Database Searched: PlantCARE
URL: <http://sphinx.rug.ac.be:8080/PlantCARE/>

Sequence submitted:

>G564 promoter (-921 to -662) 11/21/00

+ GAAAAGTGAA GAAAACCATG TAATGAAAC AAAATGGCAC GACAATCAAA AAAAGTTTTC ACGCAAAATT
+ TTCTTCAAAA TTATAACAT TTTTCATGTTG TGTTTGTTC AAAGCCTAGA AAAACGAAGA GTTACTATTG
+ GTAATGAAAA GCGAAGAAAA CCACATAATA AAAACAAAT GGCAGCAGCA TCAAGAAAAA GTTTTCACAC
+ AAAACTTTTT TCAAAATTTA CTATGTTTAT TTCGAAATTT AGAAAAACGA AGAGTTATTA TTAGTAATGA
+ AAAGCGAAGA AAACACGTA ATAAAAACA AAATGGCACG ACAATAAAAA AAGTTTTCAC GCAAAATTTT
+ CTGTGTGCGC AGAAAGTTAT ATATATTAAT TAATTAAATT TCATTACTT TTTCCCTTT TTATTTTAAA
+ GTTAAATTAT TATTATTTTC ATTTAAAA

- CTTTTCACCT CTTTGTGAC ATTACTTTT TTTACCGTG CTGTAGTTT TTTCAAAAG TCGTGTTTAA
- AAGAAGTTT AAATATGTA AAAGTACAAC ACAACAAGG TTTCGGATCT TTTGCTTCT CAATGATAAC
- CATTACTTT CGCTTCTTT GGTGATTAT TTTGTTTTC CCGTGCTGT AGTTCTTTT CAAAAGTGTG
- TTTTGAAAA AGTTTAAAT GATACAAATA AAGCTTTTAA TCTTTTGTCT TCTCAATAAT AATCATTACT
- TTTCGCTTCT TTTGATGCAT TATTTTGTG TTTACCGTGC TGTATTTTT TTCAAAAGTG CGTTTAAAA
- GAACCACGCG TCTTTCAATA TATATAATTA ATTAATTAAA AGTAAATGAA AAAAGGGAAA AATAAAATTT
- CAATTTAATA ATAATAAAG TAAATTTT

3-AFL_binding_sit

Site Name	Organism	Position	Strand	Core simil.	Matrix simil	sequence
3-AFL_binding_sit	ST	260	+	1.000	0.860	AAGAgttatt

Function:

AAGAA-motif

Site Name	Organism	Position	Strand	Core simil.	Matrix simil	sequence
AAGAA-motifAvena sativa		6	+	1.000	0.903	gtgAAGAA
AAGAA-motifAvena sativa		151	+	1.000	0.870	gcgAAGAA
AAGAA-motifAvena sativa		284	+	1.000	0.870	gcgAAGAA

Function:

ABRE

Site Name	Organism	Position	Strand	Core simil.	Matrix simil	sequence
ABRE	Hordeum vulgare	293	+	1.000	0.854	actACGTaat

Function: cis-acting element involved in the abscisic acid responsiveness

ACE

Site Name	Organism	Position	Strand	Core simil.	Matrix simil	sequence
ACE	Petroselinum crispum	293	+	1.000	0.908	actACGTAat

Function: cis-acting element involved in light responsiveness

.AE-box

Site Name	Organism	Position	Strand	Core simil.	Matrix simil	sequence
AE-box	Arabidopsis thaliana	67	-	1.000	0.852	AGAAaatt
AE-box	Arabidopsis thaliana	345	-	1.000	0.852	AGAAaatt
AE-box	Arabidopsis thaliana	361	+	1.000	0.852	AGAAagtt

Function: part of a module for light response

AT1-motif

Site Name	Organism	Position	Strand	Core simil.	Matrix simil	sequence
AT1-motif	Solanum tuberosum	409	+	1.000	0.859	ttttATTTtaaa

Function: part of a light responsive module

Box_4

Site Name	Organism	Position	Strand	Core simil.	Matrix simil	sequence
Box_4	PC	375	+	1.000	1.000	ATTAat
Box_4	PC	379	+	1.000	1.000	ATTAat
Box_4	PC	383	-	1.000	1.000	ATTAat

Function:

Box_I

Site Name	Organism	Position	Strand	Core simil.	Matrix simil	sequence
Box_I	PS	107	+	1.000	1.000	TTTCaaa
Box_I	PS	203	+	1.000	0.857	TTTCaca
Box_I	PS	219	+	1.000	1.000	TTTCaaa
Box_I	PS	240	+	1.000	0.857	TTTCgaa
Box_I	PS	241	-	1.000	0.857	TTTCgaa
Box_I	PS	249	-	1.000	0.857	TTTCtaa

Function:

Box_II

Site Name	Organism	Position	Strand	Core simil.	Matrix simil	sequence
-----------	----------	----------	--------	-------------	--------------	----------

Box_II	ST	139	+	1.000	0.889	TGGTaatga
Box_II	AT	161	+	1.000	0.954	CCACataat

Function:

CAAT-box

Site Name	Organism	Position	Strand	Core simil.	Matrix simil	sequence
CAAT-box	Hordeum vulgare	43	+	1.000	1.000	CAAT
CAAT-box	Arabidopsis thaliana	137	-	1.000	1.000	aCCAAt
CAAT-box	Hordeum vulgare	188	+	1.000	1.000	CAAT
CAAT-box	Hordeum vulgare	322	+	1.000	1.000	CAAT
CAAT-box	Arabidopsis thaliana	351	-	1.000	0.857	aCCAAG

Function: common cis-acting element in promoter and enhancer regions

ERE

Site Name	Organism	Position	Strand	Core simil.	Matrix simil	sequence
ERE	Dianthus caryophyllus	239	+	1.000	0.875	ATTTCgaa
ERE	Dianthus caryophyllus	241	-	1.000	0.875	ATTTcgaa
ERE	Dianthus caryophyllus	413	+	1.000	0.875	ATTTtaaa
ERE	Dianthus caryophyllus	441	+	1.000	0.875	ATTTaaaa
ERE	Dianthus caryophyllus	442	-	1.000	0.875	ATTTtaaa

Function: ethylene-responsive element

G-box

Site Name	Organism	Position	Strand	Core simil.	Matrix simil	sequence
G-box	Zea mays	17	+	0.842	0.870	CATGta
G-box	Zea mays	38	+	1.000	0.903	CACGac
G-box	Zea mays	94	+	0.842	0.886	CATgtt
G-box	Zea mays	183	+	1.000	0.903	CACGac
G-box	Zea mays	317	+	1.000	0.903	CACGac

Function: cis-acting regulatory element involved in light responsiveness

GC-repeat

Site Name	Organism	Position	Strand	Core simil.	Matrix simil	sequence
GC-repeat	Oryza sativa	351	-	1.000	1.000	gCACCaag

Function: ?

HSE

Site Name	Organism	Position	Strand	Core simil.	Matrix simil	sequence
HSE	Brassica oleracea	49	+	0.944	0.878	aAAAAggtt
HSE	Brassica oleracea	50	+	0.944	0.912	aAAAAgttt
HSE	Brassica oleracea	52	-	0.944	0.874	gAAAActtt
HSE	Brassica oleracea	66	-	1.000	0.978	agAAAattt
HSE	Brassica oleracea	77	-	0.833	0.868	aTAAAtttt
HSE	Brassica oleracea	87	-	1.000	0.853	tGAAAatgt
HSE	Brassica oleracea	196	+	0.944	0.912	aAAAAgttt
HSE	Brassica oleracea	198	-	0.944	0.874	gAAAActtt
HSE	Brassica oleracea	210	+	0.944	0.874	cAAAActtt
HSE	Brassica oleracea	212	-	0.944	0.912	aAAAAgttt
HSE	Brassica oleracea	213	-	0.944	0.878	aAAAAgttt
HSE	Brassica oleracea	327	+	0.944	0.878	aAAAAgttt
HSE	Brassica oleracea	328	+	0.944	0.912	aAAAAgttt
HSE	Brassica oleracea	330	-	0.944	0.874	gAAAActtt
HSE	Brassica oleracea	344	-	1.000	0.978	agAAAattt
HSE	Brassica oleracea	361	+	1.000	0.888	agAAAgtta
HSE	Brassica oleracea	385	-	1.000	0.853	tGAAAatta

Function: cis-acting element involved in heat stress responsiveness

I-box

Site Name	Organism	Position	Strand	Core simil.	Matrix simil.	sequence
I-box	Pisum sativum	93	-	0.857	0.883	aACATga
I-box	Pisum sativum	162	+	0.857	0.883	CACATaa
I-box	Solanum tuberosum	163	-	1.000	1.000	TATTATgt
I-box	Pisum sativum	237	-	0.857	0.941	gAAATaa
I-box	Pisum sativum	367	-	1.000	1.000	TATATaa
I-box	Pisum sativum	372	+	1.000	0.941	TATATa
I-box	Pisum sativum	391	-	0.857	0.941	tAAATga
I-box	Pisum sativum	411	-	0.857	0.883	aAAATaa
I-box	Pisum sativum	423	+	0.857	0.883	tAAATa
I-box	Solanum tuberosum	424	-	1.000	0.903	aATAAttt
I-box	Arabidopsis thaliana	426	-	1.000	0.863	aATAAtaat
I-box	Arabidopsis thaliana	429	-	1.000	0.863	aATAAtaat
I-box	Solanum tuberosum	431	+	1.000	0.951	tATTATtt
I-box	Pisum sativum	433	-	0.857	0.883	aAAATaa
I-box	Pisum sativum	439	-	0.857	0.941	tAAATga

Function: part of a light responsive element

P-box

Site Name	Organism	Position	Strand	Core simil.	Matrix simil.	sequence
P-box	Oryza sativa	406	+	1.000	0.857	CCTTttt

Function: gibberellin-responsive element

Prolamin_box

Site Name	Organism	Position	Strand	Core simil.	Matrix simil.	sequence
Prolamin-box	Oryza sativa	145	+	1.000	0.913	tgaAAAGc
Prolamin-box	Oryza sativa	278	+	1.000	0.913	tgaAAAGc

Function: cis-acting regulatory element associated with GCN4

TATA-box

Site Name	Organism	Position	Strand	Core simil.	Matrix simil.	sequence
TATA-box	Daucus carota	79	-	1.000	1.000	TATAaatl
TATA-box	Brassica juncea	80	-	1.000	1.000	TATAaat
TATA-box	Helianthus annuus	81	-	1.000	1.000	TATAaa
TATA-box	Brassica oleracea	82	+	1.000	0.908	tTATAac
TATA-box	Brassica napus	83	-	1.000	0.892	gtTATA
TATA-box	Oryza sativa	117	+	0.818	0.912	TAGAAAA
TATA-box	Oryza sativa	169	+	0.818	0.872	TAAAAac
TATA-box	Zea mays	248	+	0.909	0.879	TTTAGaaa
TATA-box	Oryza sativa	250	+	0.818	0.912	TAGAAAA
TATA-box	Oryza sativa	302	+	0.818	0.912	TAAAAaa
TATA-box	Oryza sativa	325	+	0.818	0.912	TAAAAaa
TATA-box	Daucus carota	364	-	1.000	0.863	TATAactt
TATA-box	Brassica juncea	365	-	1.000	0.857	TATAact
TATA-box	Zea mays	366	-	1.000	0.879	TATAtaac

TATA-box	Oryza sativa	367	-	1.000	0.956	TATATAa
TATA-box	Oryza sativa	368	+	1.000	0.929	TATAtat
TATA-box	Oryza sativa	369	-	1.000	0.929	TATAtat
TATA-box	Solanum tuberosum	370	-	1.000	1.000	TATATA
TATA-box	Glycine max	372	+	1.000	0.891	TATAtt
TATA-box	Oryza sativa	407	-	0.818	0.872	TAAAAag
TATA-box	Zea mays	413	-	0.909	0.879	TTTAAaat
TATA-box	Zea mays	442	+	0.909	0.879	TTTAAaat

Function: core promoter element around -30 of transcription start

.TC-rich_repeats

Site Name	Organism	Position	Strand	Core simil.	Matrix simil	sequence
TC-rich_repeats	NT	7	-	1.000	0.952	gTTTtcttca
TC-rich_repeats	NT	68	+	1.000	1.000	aTTTtcttca
TC-rich_repeats	NT	152	-	1.000	0.909	gTTTtcttcg
TC-rich_repeats	NT	191	-	1.000	0.885	tTTTtcttga
TC-rich_repeats	NT	248	-	1.000	0.914	tTTTtctaaa
TC-rich_repeats	NT	285	-	1.000	0.909	gTTTtcttcg
TC-rich_repeats	NT	346	+	1.000	0.915	aTTTtcttcg

Function:

WUN-motif

Site Name	Organism	Position	Strand	Core simil.	Matrix simil	sequence
WUN-motif	Brassica oleracea	18	-	1.000	0.948	tCATTacat
WUN-motif	Brassica oleracea	139	-	1.000	1.000	tCATTacca
WUN-motif	Brassica oleracea	237	+	0.857	0.948	tTATTTcga
WUN-motif	Brassica oleracea	242	-	1.000	1.000	aAATTtcga
WUN-motif	Brassica oleracea	272	-	1.000	0.948	tCATTTacta
WUN-motif	Brassica oleracea	296	-	0.857	0.948	tTATTacgt

Function: wound-responsive element

DECLARATION

As a below named inventor, I declare that:

My residence, post office address and citizenship are as stated below next to my name; I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural inventors are named below) of the subject matter which is claimed and for which a patent is sought on the invention entitled: **POLYNUCLEOTIDES USEFUL FOR MODULATING TRANSCRIPTION** the specification of which X is attached hereto or was filed on as Application No. and was amended on (if applicable).

I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above. I acknowledge the duty to disclose information which is material to patentability as defined in Title 37, Code of Federal Regulations, Section 1.56. I claim foreign priority benefits under Title 35, United States Code, Section 119 of any foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of the application on which priority is claimed.

Prior Foreign Application(s)

Country	Application No.	Date of Filing	Priority Claimed Under 35 USC 119

I hereby claim the benefit under Title 35, United States Code § 119(e) of any United States provisional application(s) listed below:

Application No.	Filing Date

I claim the benefit under Title 35, United States Code, Section 120 of any United States application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code, Section 112, I acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations, Section 1.56 which occurred between the filing date of the prior application and the national or PCT international filing date of this application:

Application No.	Date of Filing	Status

Full Name of Inventor 1:	Last Name: WETERINGS	First Name: KOEN	Middle Name or Initial:	
Residence & Citizenship:	City:	State/Foreign Country: Netherlands	Country of Citizenship: Netherlands	
Post Office Address:	Post Office Address: Heidevenstraat 44 6533 TS	City: Nijmegen	State/Country: Netherlands	Postal Code:
Full Name of Inventor 2:	Last Name: APUYA	First Name: NESTOR	Middle Name or Initial: R.	
Residence & Citizenship:	City: Culver City	State/Foreign Country: California	Country of Citizenship: Philippines	
Post Office Address:	Post Office Address: 5396 Dobson Way	City: Culver City	State/Country: California	Postal Code: 90230

Full Name of Inventor 3:	Last Name: GOLDBER	First Name: ROBERT	Middle Name or Initial: B.	
Residence & Citizenship:	City: Topanga	State/Foreign Country: California	Country of Citizenship: United States	
Post Office Address:	Post Office Address: 21907 Canon Drive	City: Topanga	State/Country: California	Postal Code: 90290

I further declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

Signature of Inventor 1	Signature of Inventor 2	Signature of Inventor 3
<u>Koen Weterings</u>	<u>Nestor R. Apuya</u>	<u>Robert B. Goldber</u>
Date	Date	Date

SF 1163844 v1